

Formula Sheet (STAT-319-Second Exam)

- **Sample Mean:** $\bar{X} = \frac{\sum X_k}{n}$

- **Variance:**

$$s^2 = \frac{\sum (X_i - \bar{X})^2}{n-1} = \frac{\sum x^2 - n(\bar{X})^2}{n-1}$$

- **Locating Percentiles:** $P\alpha$

$$R\alpha = \frac{\alpha}{100}(n+1) = i.d$$

$$P\alpha = X_{(i)} + d(X_{(i+1)} - X_{(i)})$$

- **Coefficient of variation:** $v = s/\bar{X}$

- **Coefficient of skewness:**

$$SK = \frac{3(\bar{X} - m)}{s}$$

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A^\complement) = 1 - P(A)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$P(A_1 \cap A_2 \dots A_k) =$$

$$P(A_1) P(A_2|A_1) P(A_3|A_1 \cap A_2) \dots P(A_k|A_1 \cap A_2 \dots A_{k-1})$$

- **Total Rule of Probability:**

$$P(A) = \sum_{i=1}^k P(A|B_i) P(B_i)$$

- **Bayes' Rule:**

$$\begin{aligned} P(A|B_r) &= \frac{P(A|B_r)P(B_r)}{P(A)} \\ &= \frac{P(A|B_r)P(B_r)}{\sum_{i=1}^k P(A|B_i)P(B_i)} \end{aligned}$$

$$E(X) = \sum xf(x) \text{ or } \int_{-\infty}^{\infty} xf(x)dx$$

$$\sigma^2 = \sum [x - \mu]^2 f(x) \text{ or } \int_{-\infty}^{\infty} [x - \mu]^2 f(x) dx$$

$$\sigma^2 = E(x^2) - [E(x)]^2$$

$$\mu_{g(x)} = E[g(x)] = \sum g(x)f(x) \text{ or } \int_{-\infty}^{\infty} g(x)f(x)dx$$

$$\sigma_{g(x)}^2 = E[g(x) - \mu_{g(x)}]^2$$

$$F(x) = P(X \leq x) = \sum_{t=-\infty}^x f(t) \text{ or } \int_{-\infty}^x f(t)dt$$

Binomial distribution

$$b(x; n, p) = \binom{n}{x} p^x q^{n-x}, \mu = np$$

$$\sigma^2 = npq$$

Poisson distribution

$$p(x; \lambda, t) = \Pr[X = x] = \frac{(\lambda t)^x e^{-\lambda t}}{x!},$$

$$E(X) = \text{Var}(X) = \lambda t$$

Geometric Distribution

$$g(x; p) = p q^{x-1}, q = 1-p, \mu = \frac{1}{p},$$

$$\sigma^2 = \frac{q}{p^2}$$

Hypergeometric Distribution:

$$h(x, n, k, N) = P(X = x) = \frac{\binom{K}{x} \binom{N-K}{n-x}}{\binom{N}{n}}$$

$$\mu = E(X) = \frac{nK}{N},$$

$$\sigma^2 = \frac{nK}{N} \left(1 - \frac{K}{N}\right) \left(\frac{N-n}{N-1}\right)$$

The gamma function

$$\Gamma(\alpha) = \int_0^{\infty} x^{\alpha-1} e^{-x} dx, \alpha > 0$$

$$\Gamma(n) = (n-1)\Gamma(n-1)$$

if n is positive integer, $\Gamma(n) = (n-1)!$

The Exponential Distribution:

$$f(x) = \begin{cases} \frac{1}{\beta} e^{-x/\beta}, & x > 0 \\ 0, & \text{otherwise} \end{cases}$$

$$\mu = \beta, \sigma^2 = \beta^2$$

The gamma Distribution

$$f(x) = \begin{cases} \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-x/\beta}, & x > 0 \\ 0, & \text{ow} \end{cases}$$

$$\mu = \alpha\beta, \quad \sigma^2 = \alpha\beta^2$$

Sampling distribution

$$\mu_{\bar{X}} = \mu, \quad \sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}},$$

$$\mu_{\bar{X}_1 - \bar{X}_2} = \mu_1 - \mu_2,$$

$$\sigma_{\bar{X}_1 - \bar{X}_2}^2 = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}$$

Student t-distribution

$$t = \frac{\bar{X} - \mu}{s/\sqrt{n}}, \quad df = n - 1,$$

$$\alpha = \Pr[t_{n-1} > t_\alpha]$$

Chi-Square Distribution:

$$\chi^2 = \frac{(n-1)s^2}{\sigma^2}, \quad df = n - 1$$

F-distribution

$$F = \frac{s_1^2}{s_2^2}, \quad df_1 = n_1 - 1, df_2 = n_2 - 1$$

$$F_{\alpha, n_1, n_2} = \frac{1}{F_{1-\alpha, n_2, n_1}}$$