

Towards a Direct Method for the Analyticity of the Pressure for Certain Classical Unbounded Spin Systems

Assane Lo
The University of Arizona

November 12, 2006

Abstract

The aim of this paper is to study direct methods for the analyticity of the pressure for certain classical unbounded spin models. We provide a representation of the n th-derivatives of the pressure as function of some thermodynamic parameter t in terms of the Witten Laplacian on one-forms. The technique involves the formula for the covariance introduced by B. Helffer and J. Sjostrand.

1 Introduction

As already mentioned in [66], The methods for investigating phase transitions for certain physical systems took an interesting direction when powerful and sophisticated PDE techniques are introduced. The methods are generally based on the analysis of suitable differential operators

$$\mathbf{W}_\Phi^{(0)} = \left(-\Delta + \frac{|\nabla\Phi|^2}{4} - \frac{\Delta\Phi}{2} \right)$$

and

$$\mathbf{W}_\Phi^{(1)} = -\Delta + \frac{|\nabla\Phi|^2}{4} - \frac{\Delta\Phi}{2} + \mathbf{Hess}\Phi,$$

which are in some sense deformations of the standard Laplace Beltrami operator. These operators commonly called Witten Laplacians were first introduced by Edward Witten, [18] in 1982 in the context of Morse theory for the study of topological invariants of compact Riemannian manifolds. In 1994, Bernard Helffer and Jöhanne Sjostrand [8] introduced two elliptic differential operators

$$A_\Phi^{(0)} := -\Delta + \nabla\Phi \cdot \nabla$$

and

$$A_\Phi^{(1)} := -\Delta + \nabla\Phi \cdot \nabla + \mathbf{Hess}\Phi$$

sometimes called Helffer-Sjostrand operators serving to get direct method for the study of integrals and operators in high dimensions of the type that appear in Statistical Mechanics and Euclidean field theory. In 1996, Jöhanne Sjostrand [13] observed that these so called Helffer-Sjostrand operators are in fact equivalent to Witten's Laplacians. Since then, there has been significant advances in the use of these Laplacians to study the thermodynamic behavior of quantities related to the Gibbs measure $Z^{-1}e^{-\Phi}dx$.

Numerous techniques have been developed in the study of integrals associated to the equilibrium Gibbs state for certain unbounded spins systems. One of the most striking result is an exact formula for the covariance of two functions in terms of the Witten Laplacian on one forms leading to sophisticated methods for estimating the correlation functions of a random field. As mentioned in [10], this formula is in some sense a stronger and more flexible version of the Brascamp-Lieb inequality [1]. The formula may be written as follow:

$$\mathbf{cov}(f, g) = \int \left(A_{\Phi}^{(1)-1} \nabla f \cdot \nabla g \right) e^{-\Phi(x)} dx. \quad (1)$$

To understand the idea behind the formula mentioned above, let us denote by $\langle f \rangle$ the mean value of f with respect to the measure

$$e^{-\Phi(x)} dx,$$

the covariance of two functions f and g is defined by

$$\mathbf{cov}(f, g) = \langle (f - \langle f \rangle)(g - \langle g \rangle) \rangle \quad (2)$$

If one wants to have an expression of the covariance in the form

$$\mathbf{cov}(f, g) = \langle \nabla g \cdot \mathbf{w} \rangle_{L^2(\mathbb{R}^n, \mathbb{R}^n; e^{-\Phi} dx)}, \quad (3)$$

for a suitable vector field \mathbf{w} , we get, after observing that $\nabla g = \nabla(g - \langle g \rangle)$,

$$\mathbf{cov}(f, g) = \int (g - \langle g \rangle) (\nabla \Phi - \nabla) \cdot \mathbf{w} e^{-\Phi(x)} dx, \quad (4)$$

this leads to the question of solving the equation

$$f - \langle f \rangle = (\nabla \Phi - \nabla) \cdot \mathbf{w}. \quad (5)$$

Now trying to solve this above equation with $\mathbf{w} = \nabla u$, we obtain the equation

$$\left. \begin{aligned} f - \langle f \rangle &= A_{\Phi}^{(0)} u \\ \langle u \rangle &= 0 \end{aligned} \right\} \quad (6)$$

Assuming for now the existence of a smooth solution, we get by differentiation of this above equation

$$\nabla f = A_{\Phi}^{(1)} \nabla u \quad (7)$$

and the formula is now easy to see.

We attempt in these notes to study a direct method for the analyticity of the pressure for certain classical unbounded spin systems. It is central in statistical mechanics to study the differentiability or even the analyticity of the pressure with respect to some distinguished thermodynamic parameters such as temperature, chemical potential or external field. In fact the analytic behavior of the pressure is the classical thermodynamic indicator for the absence or existence of phase transition. The most famous result on the analyticity of the pressure is the circle theorem of Lee and Yang [28]. This theorem asserts the following. Consider a $\{-1, 1\}$ -valued spin system with ferromagnetic pair interaction and external field h and regard the quantity $z = e^h$ as a complex parameter, then all zeroes of all partition functions (with free boundary condition), considered as functions of z lie in the complex unit circle. This theorem readily implies that the pressure is an analytic function of h in the region $h > 0$ and $h < 0$. Heilmann [29] showed that the assumption of pair interaction is necessary. A transparent approach to the circle theorem was found by Asano [30] and developed further by Ruelle [31],[32], Slawny [33], and Gruber et al [34]. Griffiths [35] and Griffiths-Simon [36] found a method of extending the Lee-Yang theorem to real-valued spin systems with a particular type of a priory measure. Newman [37] proved the Lee-Yang theorem for every a priory measure which satisfies this theorem in the particular case of no interaction. Dunlop [38],[39] studied the zeroes of the partition functions for the plane rotor model. A general Lee-Yang theorem for multicomponent systems was finally proved by Lieb and Sokal [40]. For further references see Glimm and Jaffe [41].

The Lee-Yang theorem theorem and its variants depend on the ferromagnetic character of the interaction. There are various other way of proving the infinite differentiability or the analyticity of the pressure for (ferromagnetic and non ferromagnetic) systems at high temperatures, or at low temperatures, or at large external fields. Most of these take advantage of a sufficiently rapid decay of correlations and /or cluster expansion methods. Here is a small sample of relevant references. Bricmont, Lebowitz and Pfister [42], Dobroshin [43], Dobroshin and Sholsman [44],[45], Duneau et al [46],[47],[48], Glimm and Jaffe [41],[49], Israel [50], Kotecky and Preiss [51], Kunz [52], Lebowitz [53],[54], Malyshev [55], Malychev and Milnos [56] and Prakash [57]. M. Kac and J.M. Luttinger [58] obtained a formula for the pressure in terms of irreducible distribution functions. In this present study, we propose a new way of analyzing the analyticity of the

pressure for certain unbounded models through a representation by means of the Witten Laplacians of the remainder in the taylor expansion. The methods known up to now rely on complicated indirect arguments.

2 Towards the analyticity of the Pressure

Let Λ be a finite domain in \mathbb{Z}^d ($d \geq 1$) and consider the Hamiltonian of the phase space given by,

$$\Phi(x) = \Phi_\Lambda(x) = \frac{x^2}{2} + \Psi(x), \quad x \in \mathbb{R}^\Lambda. \quad (8)$$

where

$$|\partial^\alpha \nabla \Psi| \leq C_\alpha, \quad \forall \alpha \in \mathbb{N}^{|\Lambda|}, \quad (9)$$

$$\mathbf{Hess} \Phi(x) \geq \delta_o, \quad 0 < \delta_o < 1 \quad (10)$$

Let g is a smooth function on \mathbb{R}^Γ with lattice support $S_g = \Gamma$. We identified with \tilde{g} defined on \mathbb{R}^Λ by

$$\tilde{g}(x) = g(x_\Gamma) \quad \text{where } x = (x_i)_{i \in \Lambda} \quad \text{and } x_\Gamma = (x_i)_{i \in \Gamma} \quad (11)$$

and satisfying

$$|D^\alpha \nabla g| \leq C_\alpha \quad \forall \alpha \in \mathbb{N}^{|\Gamma|} \quad (12)$$

Under the additional assumptions that Ψ is compactly supported in \mathbb{R}^Λ and g is compactly supported in \mathbb{R}^Γ , it was proved in [66] (see also [8]) that the equation

$$\begin{cases} -\Delta f + \nabla \Phi \cdot \nabla f = g - \langle g \rangle \\ \langle f \rangle_{L^2(\mu)} = 0 \end{cases}$$

has a unique smooth solution satisfying $\nabla^k f(x) \rightarrow 0$ as $|x| \rightarrow \infty$ for every $k \geq 1$. Recall also that ∇f is a solution of the system

$$(-\Delta + \nabla \Phi \cdot \nabla) \nabla f + \mathbf{Hess} \Phi \nabla f = \nabla g \quad \text{in } \mathbb{R}^\Lambda \quad (13)$$

As in [66] and [8], these assumptions will be relaxed later on.

Let

$$\Phi_\Lambda^t(x) = \Phi(x) - tg(x) \quad (14)$$

where $x = (x_i)_{i \in \Lambda}$, and assume additionally that g satisfies

$$\mathbf{Hess} g \leq C. \quad (15)$$

We consider the following perturbation

$$\theta_\Lambda(t) = \log \left[\int dx e^{-\Phi_\Lambda^t(x)} \right] \quad (16)$$

Denote by

$$Z_t = \int dx e^{-\Phi_\Lambda^t(x)} \quad (17)$$

and

$$\langle \cdot \rangle_{t,\Lambda} = \frac{\int \cdot dx e^{-\Phi_\Lambda^t(x)}}{Z_t}. \quad (18)$$

3 Parameter dependency of the solution

From the assumptions made on Φ and g , it is easy to see that there exists $T > 0$ such that for every $t \in [0, T)$, $\Phi_\Lambda^t(x)$ satisfies all the assumptions required for the solvability, regularity and asymptotic behavior of the solution $f(t)$ associated with the potential $\Phi_\Lambda^t(x)$. Thus, each $t \in [0, T)$ is associated with a unique C^∞ -solution, $f(t)$ of the equation

$$\begin{cases} A_{\Phi_\Lambda^t}^{(0)} f(t) = g - \langle g \rangle_{L^2(\mu)} \\ \langle f(t) \rangle_{L^2(\mu)} = 0 \end{cases}$$

Hence,

$$A_{\Phi_\Lambda^t}^{(1)} \mathbf{v}(t) = \nabla g \quad (19)$$

where $\mathbf{v}(t) = \nabla f(t)$. Notice that the map

$$t \longmapsto \mathbf{v}(t)$$

is well defined and

$$\{\mathbf{v}(t) : t \in [0, T)\}$$

is a family of smooth solutions on \mathbb{R}^Λ satisfying

$$\partial^\alpha \mathbf{v}(t) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad \forall \alpha \in \mathbb{N}^{|\Lambda|} \quad \text{and for each } t \in [0, T)$$

and corresponding to the family of potential

$$\{\Phi_\Lambda^t : t \in [0, T)\} \quad (20)$$

Let us now verify that \mathbf{v} is a smooth function of $t \in (0, T)$. We need to prove that for each $t \in (0, T)$, the limit

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathbf{v}(t + \varepsilon) - \mathbf{v}(t)}{\varepsilon}$$

exists. Let

$$\mathbf{v}^\varepsilon(t) = \frac{\mathbf{v}(t + \varepsilon) - \mathbf{v}(t)}{\varepsilon}$$

We use a technique based on regularity estimates to get a uniform control of $\mathbf{v}^\varepsilon(t)$ with respect to ε .

With ε small enough, we have

$$\begin{aligned} 0 &= -\Delta \left[\frac{\mathbf{v}(t + \varepsilon) - \mathbf{v}(t)}{\varepsilon} \right] + \frac{\nabla \Phi^{t+\varepsilon} \cdot \nabla \mathbf{v}(t + \varepsilon) - \nabla \Phi^t \cdot \nabla \mathbf{v}(t)}{\varepsilon} \\ &\quad + \frac{\mathbf{Hess} \Phi^{t+\varepsilon} \mathbf{v}(t + \varepsilon) - \mathbf{Hess} \Phi^t \mathbf{v}(t)}{\varepsilon} \end{aligned}$$

Equivalently,

$$\begin{aligned}
& -\Delta \left[\frac{\mathbf{v}(t+\varepsilon) - \mathbf{v}(t)}{\varepsilon} \right] + \frac{\nabla\Phi^{t+\varepsilon} \cdot \nabla [\mathbf{v}(t+\varepsilon) - \mathbf{v}(t)]}{\varepsilon} \\
& + \mathbf{Hess}\Phi^{t+\varepsilon} \left(\frac{\mathbf{v}(t+\varepsilon) - \mathbf{v}(t)}{\varepsilon} \right) \\
= & - \left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t) - \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla \mathbf{v}(t)
\end{aligned}$$

and

$$\begin{aligned}
& -\Delta \mathbf{v}^\varepsilon(t) + \nabla\Phi^{t+\varepsilon} \cdot \nabla \mathbf{v}^\varepsilon(t) + \mathbf{Hess}\Phi^{t+\varepsilon} \mathbf{v}^\varepsilon(t) \\
= & - \left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t) - \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla \mathbf{v}(t)
\end{aligned}$$

Let $\mathbf{w}(t)$ be the unique C^∞ -solution of the system

$$-\Delta \mathbf{w}(t) + \nabla\Phi^t \cdot \nabla \mathbf{w}(t) + \mathbf{Hess}\Phi^t \mathbf{w}(t) = \mathbf{Hess}g\mathbf{v}(t) - \nabla g \cdot \nabla \mathbf{v}(t) \quad (21)$$

Combining the last two systems above, we get

$$\begin{aligned}
& -\Delta [\mathbf{w}(t) - \mathbf{v}^\varepsilon(t)] + \nabla\Phi^t \cdot \nabla [\mathbf{w}(t) - \mathbf{v}^\varepsilon(t)] + \mathbf{Hess}\Phi^t [\mathbf{w}(t) - \mathbf{v}^\varepsilon(t)] \\
= & \mathbf{Hess}g\mathbf{v}(t) - \nabla g \cdot \nabla \mathbf{v}(t) + \left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t) \\
& + \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla \mathbf{v}(t) + (\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t) \cdot \nabla \mathbf{v}^\varepsilon(t) \\
& + (\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t) \mathbf{v}^\varepsilon(t)
\end{aligned} \quad (22)$$

Now using the unitary transformation U_{Φ^t} , we get

$$\begin{aligned}
& \left(-\Delta + \frac{|\nabla\Phi^t|^2}{4} - \frac{\Delta\Phi^t}{2} \right) (\mathbf{w}(t) - \mathbf{v}^\varepsilon(t)) e^{-\Phi^t/2} \\
& + \mathbf{Hess}\Phi^t (\mathbf{w}(t) - \mathbf{v}^\varepsilon(t)) e^{-\Phi^t/2} \\
= & o_\varepsilon(1) e^{-\Phi^t/2} + [(\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t) \cdot \nabla \mathbf{v}^\varepsilon(t) \\
& + (\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t) \mathbf{v}^\varepsilon(t)] e^{-\Phi^t/2}
\end{aligned} \quad (23)$$

Next, we propose to estimate the last two terms of the right hand side of this equation.

Again using the unitary transformation $U_{\Phi^{t+\varepsilon}}$, we reduce the system

$$\begin{aligned}
& -\Delta \mathbf{v}^\varepsilon(t) + \nabla\Phi^{t+\varepsilon} \cdot \nabla \mathbf{v}^\varepsilon(t) + \mathbf{Hess}\Phi^{t+\varepsilon} \mathbf{v}^\varepsilon(t) \\
= & - \left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t) \\
& - \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla \mathbf{v}(t)
\end{aligned} \quad (24)$$

into

$$\begin{aligned}
& \left(-\Delta + \frac{|\nabla\Phi^{t+\varepsilon}|^2}{4} - \frac{\Delta\Phi^{t+\varepsilon}}{2} \right) \mathbf{V}^\varepsilon + \mathbf{Hess}\Phi^{t+\varepsilon}\mathbf{V}^\varepsilon = \\
& - \left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t)e^{-\Phi^{t+\varepsilon}/2} \\
& - \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla\mathbf{v}(t)e^{-\Phi^{t+\varepsilon}/2}
\end{aligned} \tag{25}$$

where $\mathbf{V}^\varepsilon = \mathbf{v}^\varepsilon(t)e^{-\Phi^{t+\varepsilon}/2}$. Taking scalar product with \mathbf{V}^ε on both sides of this last equality and integrating, we get

$$\begin{aligned}
& \left\| \left(\partial_x + \frac{\nabla\Phi^{t+\varepsilon}}{2} \right) \mathbf{V}^\varepsilon \right\|_{L^2}^2 + \int \mathbf{Hess}\Phi^{t+\varepsilon}\mathbf{V}^\varepsilon \cdot \mathbf{V}^\varepsilon dx = \\
& - \int \left[\left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t)e^{-\Phi^{t+\varepsilon}/2} \right] \cdot \mathbf{V}^\varepsilon dx \\
& - \left[\int \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla\mathbf{v}(t)e^{-\Phi^{t+\varepsilon}/2} \right] \cdot \mathbf{V}^\varepsilon dx
\end{aligned} \tag{26}$$

Now using the uniform strict convexity on the left hand side and Cauchy-Schwartz on the right hand side, we obtain

$$\|\mathbf{V}^\varepsilon\|_{B^0} \leq C \quad \text{for small enough } \varepsilon. \tag{27}$$

We then deduce that

$$\left(-\Delta + \frac{|\nabla\Phi^{t+\varepsilon}|^2}{4} \right) \mathbf{V}^\varepsilon = \tilde{q}_\varepsilon \tag{28}$$

where

$$\begin{aligned}
\tilde{q}_\varepsilon = & - \left(\frac{\mathbf{Hess}\Phi^{t+\varepsilon} - \mathbf{Hess}\Phi^t}{\varepsilon} \right) \mathbf{v}(t)e^{-\Phi^{t+\varepsilon}/2} - \left(\frac{\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t}{\varepsilon} \right) \cdot \nabla\mathbf{v}(t)e^{-\Phi^{t+\varepsilon}/2} \\
& + \frac{\Delta\Phi^{t+\varepsilon}}{2} \mathbf{V}^\varepsilon - \mathbf{Hess}\Phi^{t+\varepsilon}\mathbf{V}^\varepsilon
\end{aligned} \tag{29}$$

is bounded in B^0 uniformly with respect to ε for ε small enough. .

Now taking scalar product with \mathbf{V}^ε on both sides of (28) and integrating by parts, we obtain

$$\|\nabla\mathbf{V}^\varepsilon\|_{L^2}^2 + \left\| \frac{|\nabla\Phi^{t+\varepsilon}|}{2} \mathbf{V}^\varepsilon \right\|_{L^2}^2 \leq \|\tilde{q}_\varepsilon\|_{L^2} \|\mathbf{V}^\varepsilon\|_{L^2} \tag{30}$$

It follows that \mathbf{V}^ε is uniformly bounded with respect to ε in $B_{\Phi^{t+\varepsilon}}^1$ for ε small enough.

Next, observe that

$$\left(-\Delta + \frac{|\nabla\Phi^t|^2}{4} \right) \mathbf{V}^\varepsilon = \hat{q}_\varepsilon \tag{31}$$

where

$$\hat{q}_\varepsilon = \tilde{q}_\varepsilon - \frac{|\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t|^2}{4}\mathbf{V}^\varepsilon + \frac{(\nabla\Phi^{t+\varepsilon} - \nabla\Phi^t) \cdot \nabla\Phi^t}{2}\mathbf{V}^\varepsilon \quad (32)$$

is uniformly bounded in B^0 with respect to ε for small enough ε . Using regularity, it follows that for small enough ε , \mathbf{V}^ε is uniformly bounded in $B_{\Phi^t}^2$ with respect to ε . This implies that \hat{q}_ε is uniformly bounded in $B_{\Phi^t}^1$ for ε small enough. Again, we can continue by a bootstrap argument to consequently get that for ε small enough, \mathbf{V}^ε is uniformly bounded in $B_{\Phi^t}^k$ for any k .

It is then clear that for small enough ε , the right hand sides of (23) is $\mathcal{O}(\varepsilon)$ in B^0 and consequently, using the same argument as above, we get that $(\mathbf{w}(t) - \mathbf{v}^\varepsilon(t))e^{-\Phi^t/2}$ is $\mathcal{O}(\varepsilon)$ in $B_{\Phi^t}^2$; again iterating the regularity argument, we obtain that for small enough ε , $(\mathbf{w}(t) - \mathbf{v}^\varepsilon(t))e^{-\Phi^t/2}$ is $\mathcal{O}(\varepsilon)$ in $B_{\Phi^t}^k$ for every k . We have proved:

Proposition 1 *Under the above on Φ and g , there exists $T > 0$ so that for each $t \in (0, T)$, $\mathbf{v}^\varepsilon(t)$ converges to $\mathbf{w}(t)$ in C^∞ .*

Remark 2 *The proposition establishes that $\mathbf{v}(t)$ is differentiable in t and $\frac{d}{dt}\mathbf{v}(t)$ is given by the unique C^∞ -solution $\mathbf{w}(t)$ of the system*

$$-\Delta\mathbf{w}(t) + \nabla\Phi^t \cdot \nabla\mathbf{w}(t) + \text{Hess}\Phi^t\mathbf{w}(t) = \text{Hess}g\mathbf{v}(t) - \nabla g \cdot \nabla\mathbf{v}(t). \quad (33)$$

Iterating this argument, we easily get that, $\mathbf{v}(t)$ is smooth in $t \in (0, T)$

Now we are ready for the following:

4 Formula for $\theta^{(n)}(t)$

For an arbitrary suitable function $f(t) = f(t, w)$

$$\frac{\partial}{\partial t} \langle f(t) \rangle_{t, \Lambda} = \langle f'(t) \rangle_{t, \Lambda} + \text{cov}(f, g). \quad (34)$$

Hence,

$$\frac{\partial}{\partial t} \langle f(t) \rangle_{t, \Lambda} = \langle f'(t) \rangle_{t, \Lambda} + \langle A_{\Phi^t}^{(1)^{-1}}(\nabla f) \cdot \nabla g \rangle_{t, \Lambda}. \quad (35)$$

Let

$$A_g f := A_{\Phi^t}^{(1)^{-1}}(\nabla f) \cdot \nabla g \quad (36)$$

Thus,

$$\frac{\partial}{\partial t} \langle f(t) \rangle_{t, \Lambda} = \left\langle \left(\frac{\partial}{\partial t} + A_g \right) f \right\rangle_{t, \Lambda}. \quad (37)$$

The linear operator $\frac{\partial}{\partial t} + A_g$ will be denoted by H_g .

$$\begin{aligned}\theta'_\Lambda(t) &= \langle g \rangle_{t,\Lambda} \\ &= \left\langle \left(\frac{\partial}{\partial t} + A_g \right)^0 g \right\rangle_{t,\Lambda} \\ &= \langle H_g^0 g \rangle_{t,\Lambda}\end{aligned}$$

$$\begin{aligned}\theta''_\Lambda(t) &= \frac{\partial}{\partial t} \langle g \rangle_{t,\Lambda} \\ &= \langle A_{\Phi^t}^{(1)-1} (\nabla g) \cdot \nabla g \rangle_{t,\Lambda} \\ &= \left\langle \left(\frac{\partial}{\partial t} + A_g \right) g \right\rangle_{t,\Lambda}\end{aligned}$$

$$\begin{aligned}\theta'''_\Lambda(t) &= \frac{\partial}{\partial t} \langle A_{\Phi^t}^{(1)-1} (\nabla g) \cdot \nabla g \rangle_{t,\Lambda} \\ &= \left\langle \frac{\partial}{\partial t} \left(A_{\Phi^t}^{(1)-1} (\nabla g) \cdot \nabla g \right) \right\rangle_{t,\Lambda} \\ &+ \left\langle \left(A_{\Phi^t}^{(1)-1} \nabla \left(A_{\Phi^t}^{(1)-1} (\nabla g) \cdot \nabla g \right) \right) \cdot \nabla g \right\rangle_{t,\Lambda} \\ &= \left\langle \left(\frac{\partial}{\partial t} + A_g \right)^2 g \right\rangle_{t,\Lambda}\end{aligned}$$

By induction it is easy to see that

$$\begin{aligned}\theta_\Lambda^{(n)}(t) &= \left\langle \left(\frac{\partial}{\partial t} + A_g \right)^{n-1} g \right\rangle_{t,\Lambda} \quad (\forall n \geq 1) \\ &= \langle H_g^{(n-1)} g \rangle_{t,\Lambda}\end{aligned}$$

Next, we propose to find a simpler formula for $\theta_\Lambda^{(n)}(t)$ that only involves A_g .

$$\begin{aligned}H_g g &= A_{\Phi^t}^{(1)-1} (\nabla g) \cdot \nabla g \\ &= A_g g\end{aligned}$$

$$H_g^2 g = \frac{\partial}{\partial t} \nabla f \cdot \nabla g + \left(A_{\Phi^t}^{(1)-1} \nabla \left(A_{\Phi^t}^{(1)-1} (\nabla g) \cdot \nabla g \right) \right) \cdot \nabla g \quad (38)$$

where f satisfies the equation

$$\nabla f = A_{\Phi^t}^{(1)-1} (\nabla g) \quad (39)$$

With $\mathbf{v}(t) = \nabla f$, as before, we get

$$\frac{\partial}{\partial t} \nabla f \cdot \nabla g = A_{\Phi^t}^{(1)-1} (\mathbf{Hess} g \mathbf{v}(t) - \nabla g \cdot \nabla \mathbf{v}(t)) \cdot \nabla g$$

and H_g^2 becomes

$$\begin{aligned} H_g^2 g &= A_{\Phi^t}^{(1)^{-1}} \left[(\mathbf{Hess} g \mathbf{v}(t) - \nabla g \cdot \nabla \mathbf{v}(t)) + \nabla \left(A_{\Phi^t}^{(1)^{-1}} (\nabla g) \cdot \nabla g \right) \right] \cdot \nabla g \\ &= A_{\Phi^t}^{(1)^{-1}} 2 \nabla (A_g g) \cdot \nabla g \\ &= 2A_g^2 g \end{aligned}$$

Proposition 3 *If*

$$\theta_\Lambda(t) = \log \left[\int dx e^{-\Phi^t(x)} \right]$$

where

$$\Phi^t(x) = \Phi_\Lambda(x) - tg(x)$$

is as above then $\theta_\Lambda^{(n)}(t)$, the n th- derivative of $\theta_\Lambda(t)$ is given by the formula

$$\theta_\Lambda'(t) = \langle g \rangle_{t,\Lambda},$$

and for $n \geq 1$

$$\theta_\Lambda^{(n)}(t) = (n-1)! \langle A_g^{n-1} g \rangle_{t,\Lambda},$$

Proof. We have already established that

$$\theta_\Lambda^{(n)}(t) = \langle H_g^{n-1} g \rangle_{t,\Lambda} \quad \text{for } n \geq 1$$

It then only remains to prove that

$$H_g^{n-1} g = (n-1)! A_g^{n-1} g \quad \text{for } n \geq 1$$

The result is already established above for $n = 1, 2, 3, \dots$. By induction, assume that

$$H_g^{n-1} g = (n-1)! A_g^{n-1} g.$$

if n is replaced by $\tilde{n} \leq n$.

$$\begin{aligned} H_g^n g &= \left(\frac{\partial}{\partial t} + A_g \right) ((n-1)! A_g^{n-1} g) \\ &= (n-1)! \left(\frac{\partial}{\partial t} A_g^{n-1} g + A_g^n g \right) \end{aligned}$$

Now

$$\begin{aligned} A_g^{n-1} g &= \left[A_{\Phi^t}^{(1)^{-1}} \nabla (A_g^{n-2} g) \right] \cdot \nabla g \\ &= \nabla \varphi_n \cdot \nabla g \end{aligned}$$

where

$$\nabla \varphi_n = \left[A_{\Phi^t}^{(1)^{-1}} \nabla (A_g^{n-2} g) \right]$$

We obtain,

$$\frac{\partial}{\partial t} \nabla \varphi_n = A_{\Phi^t}^{(1)^{-1}} \left(\frac{\partial}{\partial t} \nabla A_g^{n-2} g + \mathbf{Hess} g \nabla \varphi_n - \nabla g \cdot \nabla (\nabla \varphi_n) \right).$$

We then have

$$\begin{aligned} \frac{\partial}{\partial t} A_g^{n-1} g &= \frac{\partial}{\partial t} \nabla \varphi_n \cdot \nabla g \\ &= \left[A_{\Phi^t}^{(1)^{-1}} \left(\frac{\partial}{\partial t} \nabla A_g^{n-2} g + \mathbf{Hess} g \nabla \varphi_n - \nabla g \cdot \nabla (\nabla \varphi_n) \right) \right] \cdot \nabla g \\ &= \left[A_{\Phi^t}^{(1)^{-1}} \left(\frac{\partial}{\partial t} \nabla A_g^{n-2} g + \nabla (\nabla \varphi_n \cdot \nabla g) \right) \right] \cdot \nabla g \\ &= A_g \left[\frac{\partial}{\partial t} A_g^{n-2} g + A_g (A_g^{n-2} g) \right] \\ &= A_g H_g (A_g^{n-2} g). \\ &= A_g H_g \left(\frac{1}{(n-2)!} H_g^{(n-2)} g \right) \quad (\text{from the induction hypothesis}) \\ &= \frac{1}{(n-2)!} A_g H_g^{(n-1)} g \\ &= \frac{1}{(n-2)!} A_g ((n-1)! A_g^{n-1} g) \quad (\text{still by the induction hypothesis}) \\ &= (n-1) A_g^n g \end{aligned}$$

Thus,

$$\begin{aligned} H_g^n g &= (n-1)! (n-1+1) A_g^n g \\ &= n! A_g^n g \end{aligned}$$

■

Proposition 4 *If $g(0) = 0$, then the formula*

$$\theta_{\Lambda}^{(n)}(t) = (n-1)! \langle A_g^{n-1} g \rangle_{t, \Lambda}, \quad n \geq 2$$

still holds if we no longer require Ψ and g to be compactly supported in \mathbb{R}^Λ .

Proof. As in [8], consider the family cutoff functions

$$\chi = \chi_\varepsilon \tag{40}$$

($\varepsilon \in [0, 1]$) in $\mathcal{C}_o^\infty(\mathbb{R})$ with value in $[0, 1]$ such that

$$\begin{cases} \chi = 1 & \text{for } |t| \leq \varepsilon^{-1} \\ |\chi^{(k)}(t)| \leq C_k \frac{\varepsilon}{|t|^k} & \text{for } k \in \mathbb{N} \end{cases}$$

We could take for instance

$$\chi_\varepsilon(t) = f(\varepsilon \ln |t|)$$

for a suitable f .

We then introduce

$$\Psi_\varepsilon(x) = \chi_\varepsilon(|x|)\Psi, \quad x \in \mathbb{R}^\Lambda \quad (41)$$

and

$$g_\varepsilon(x) = \chi_\varepsilon(|x|)g \quad x \in \mathbb{R}^\Gamma \quad (42)$$

One can check that both $\Psi_\varepsilon(x)$ and $g_\varepsilon(x)$ satisfies the assumptions made above on Ψ and g . Now consider the equation

$$-\Delta f_\varepsilon + \nabla \Phi_\varepsilon^t \cdot \nabla f_\varepsilon = g_\varepsilon - \langle g_\varepsilon \rangle_{t,\Lambda}, \quad (43)$$

which implies

$$(-\Delta + \nabla \Phi_\varepsilon^t \cdot \nabla) \otimes \mathbf{v}_\varepsilon + \mathbf{Hess} \Phi_\varepsilon^t \mathbf{v}_\varepsilon = \nabla g_\varepsilon \quad (44)$$

where

$$\mathbf{v}_\varepsilon = \nabla f_\varepsilon$$

It is proved in [8] that $\mathbf{v}_\varepsilon = A_{\Phi^t}^{(1)^{-1}} \nabla g_\varepsilon$ converges in C^∞ to $A_{\Phi^t}^{(1)^{-1}} \nabla g$ as $\varepsilon \rightarrow 0$.

■

Remark 5 *If we denote by R_n the remainder of the Taylor series expansion of the pressure $P_\Lambda(t)$, given by*

$$P_\Lambda(t) = \frac{\theta_\Lambda(t)}{|\Lambda|}$$

we have

$$\begin{aligned} R_n &= \frac{P_\Lambda^{(n+1)}(t_o)}{(n+1)!} \\ &= \frac{\langle A_g^n g \rangle_{t,\Lambda}}{(n+1)|\Lambda|} \Big|_{t=t_o} \end{aligned}$$

If Φ and g are such that $\langle A_g^n g \rangle_{t,\Lambda}$ is uniformly bounded with respect to n and does not grow faster than $|\Lambda|$, we automatically get the analyticity of the pressure in the thermodynamic limit.

5 Some consequences of the formula for n th-derivative of the Pressure.

In the following, we shall additionally assume that

$$\begin{aligned} \nabla g(0) &= 0, \quad \text{and} \\ \nabla \Phi_\Lambda^t(0) &= 0 \quad \text{for all } t \in [0, T]. \end{aligned}$$

When $n = 1$, we recall that $A_g^0 g = g$,

$$\theta'_\Lambda(t) = \langle g \rangle_{t,\Lambda}$$

and if

$$\mathbf{v}(t) = \nabla f = A_{\Phi^t}^{(1)^{-1}} \nabla g,$$

then we have

$$(-\Delta + \nabla \Phi_\Lambda^t \cdot \nabla) \otimes \mathbf{v}(t) + \mathbf{Hess} \Phi_\Lambda^t \mathbf{v}(t) = \nabla g$$

and as in [8] $\mathbf{v}(t)$ is a solution of the equation

$$g = \langle g \rangle_{t,\Lambda} + \mathbf{v}(t) \cdot \nabla \Phi_\Lambda^t - \text{div} \mathbf{v}(t). \quad (45)$$

Using the assumptions above, we have

$$\begin{aligned} \theta'_\Lambda(t) &= \langle g \rangle_{t,\Lambda} \\ &= \text{div} \mathbf{v}(t)(0). \end{aligned}$$

Similarly, the formula

$$\theta_\Lambda^{(n)}(t) = (n-1)! \langle A_g^{n-1} g \rangle_{t,\Lambda},$$

implies that

$$\theta_\Lambda^{(n)}(t) = (n-1)! \text{div} \mathbf{v}_n(t)(0),$$

where

$$\mathbf{v}_n(t) = A_{\Phi^t}^{(1)^{-1}} \nabla (A_g^{n-1} g)$$

Remark 6 *The idea of representing $\theta'_\Lambda(t)$ in terms of $\text{div} \mathbf{v}(t)(0)$ is due to Helffer and Sjostrand [8] in the context of proving the exponential convergence of the thermodynamic limit in the one dimensional case.*

We conclude these notes by a discussion about the potential contribution of this results towards solving the two dimensional dipole gas problem. The dipole gas and other gases of particles interacting through Coulomb forces are very important statistical systems. In particular, for dipole gas, the lack of screening is well known [59], and the analyticity of the pressure in the high temperature and low activity region has been proved in an indirect way, by means of renormalization group methods (see [60] and [61]).

A direct proof of the analyticity of the pressure based on estimating the coefficients of the Mayer (Taylor) series is still an open problem. The close relationship between this model and the Coulomb gas in the Kostelitz-Thouless phase ($\beta > 8\pi$), go along with the non-existence of any proof for the analyticity of the pressure in the Coulomb gas. Indirect arguments are attempted in [62],[63] and

[64]. We believe that after a suitable regularization of the Coulomb potential at short distances to assure stability, we can fit the problem into the framework of the model discussed above and get an estimate of the coefficients of the Mayer series through our formula for the n th derivative of the pressure.

Acknowledgement: I would like to thank my advisor Haru Pinson for all the fruitful discussions and the help he has provided in the writing of these notes.

References

- [1] Brascamp. H and J and Lieb. E. H, *On extensions of the Brunn-Minkowski and Prekopa-Leindler theorems including inequalities for log concave functions, and with application to the diffusion equation*, *J. Funct. Analysis*, 22 (1976), 366-389.
- [2] Bodineau. T and Helffer. B, *Correlations, spectral gap and logSobolev inequalities for unbounded spins systems*, *Proc. UAB Conf. March 16-20 1999, AMS/IP stud. adv. math 16 (2000)*, 51-66.
- [3] Evans. L. C, *Partial Differential Equations* (AMS, 1998), by L. C. Evans.
- [4] Helffer. B, *Introduction to the semiclassical analysis for the schrodinger operator and applications*, *Lecture Notes in Math*, 1336 (1988).
- [5] Helffer. B, *Around a stationary phase theorem in large dimension*. *J. Funct. Anal.* 119 (1994), no. 1, 217-252.
- [6] [He3] B. Helffer, *Semiclassical analysis, Witten laplacians and statistical mechanics series on partial differential equations and applications-Vol.1 - World Scientific (2002)*.
- [7] Helffer. B, *Remarks on decay of correlations and Witten laplacians. II, analysis of the dependence on the interaction*. *Rev. Math. Phys*, 11 (1999), no. 3, 321-336
- [8] Helffer. B and Sjöstrand. J, *On the correlation for Kac-like models in the convex case*. *J. of Stat. phys*, 74 Nos.1/2, 1994.
- [9] Helffer. B and Sjöstrand. J, *Semiclassical expansions of the thermodynamic limit for a Schrödinger equation. The one well case*. *Méthodes semi-classiques, Vol. 2 (Nantes, 1991)*. *Astérisque No. 210 (1992)*, 7-8, 135-181.
- [10] Johnsen, Jon: *On the spectral properties of Witten-Laplacians, their range projections and Brascamp-Lieb's inequality*. *Integral Equations Operator Theory* 36 (2000), no. 3, 288-324.
- [11] Kneib and Jean-Marie Mignot, *Fulbert Équation de Schmoluchowski généralisée. (French) [generalized Smoluchowski equation]* *Ann. Mat. Pura Appl. (4)* 167 (1994), 257-298.

- [12] Naddaf. A and Spencer. T, *On homogenization and scaling limit of gradient perturbations of a massless free field*, *Comm. Math. Physics* 183 (1997), 55-84.
- [13] Sjöstrand. J, *Correlation asymptotics and Witten laplacians*, *Algebra and Analysis* 8, no. 1 (1996), 160-191.
- [14] Sjöstrand. J, *Exponential convergence of the first eigenvalue divided by the dimension, for certain sequences of Schrödinger operators*. *Méthodes semi-classiques, Vol. 2 (Nantes, 1991)*. *Astérisque No. 210 (1992)*, 10, 303-326.
- [15] Sjöstrand, J, *Potential wells in high dimensions. II. More about the one well case*. *Ann. Inst. H. Poincaré Phys. Théor.* 58, no. 1 (1993), 43-53.
- [16] Sjöstrand. J, *Potential wells in high dimensions. I*. *Ann. Inst. H. Poincaré Phys. Théor.* 58, no. 1 (1993), 1-41.
- [17] Yosida. K, *Functional analysis, springer classics in mathematics by Kosaku Yosida*.
- [18] Witten. E, *Supersymmetry and Morse theory*, *J. of Diff. Geom.* 17, (1982), 661-692.
- [19] Cartier. P, *Inégalités de corrélation en mécanique statistique*, *Séminaire Bourbaki 25ème année, 1972-1973, No 431*.
- [20] Kac. M, *Mathematical mechanism of phase transitions*(Gordon and Breach, New York, 1966).
- [21] Troianiello. G. M, *Elliptic Differential Equations and Obstacle Problems* (Plenum Press, New York 1987).
- [22] Berezin. F. A and Shubin. M. A, *The Schrödinger Equation* (Kluwer Academic Publisher, 1991).
- [23] Dobrushin. R. L, *The description of random field by means of conditional probabilities and conditions of its regularity*. *Theor.Prob.Appl.* 13, (1968), 197-224.
- [24] Dobrushin. R. L, *Gibbsian random fields for lattice systems with pairwise interactions*. *Funct. Anal. Appl.* 2 (1968), 292-301.
- [25] Dobrushin. R. L, *The problem of uniqueness of a Gibbs random field and the problem of phase transition*. *Funct. Anal. Appl.* 2 (1968), 302-312.
- [26] Bach. V, Jecko. T and Sjostrand. J, *Correlation asymptotics of classical lattice spin systems with nonconvex Hamilton function at low temperature*. *Ann. Henri Poincare* (2000), 59-100.
- [27] Bach. V and Møller. J. S, *Correlation at low temperature, exponential decay*. *Jour. funct. anal* 203 (2003), 93-148.

- [28] Yang. C. N and Lee. T.D, *Statistical theory of equations of state and phase transition I. Theory of condensation. Phys.Rev. 87 (1952), 404-409.*
- [29] Heilmann. O. J, *Zeros of the grand partition function for a lattice gas. J.Math.Phys. 11 (1970), 2701-2703.*
- [30] Asano. T, *Theorem on the partition functions of the Heisenberg ferromagnets. J. Phys. Soc. Jap. 29 (1970), 350-359.*
- [31] Ruelle. D, *An Extension of lee-Yang circle theorem. Phys. Rev. Letters, 26 (1971), 303-304.*
- [32] Ruelle. D, *Some remarks on the location of zeroes of the partition function for lattice systems. Commun. Math. Phys 31, (1973), 265-277.*
- [33] Slawny. J, *Analyticity and uniqueness for spin 1/2 classical ferromagnetic lattice systems at low temperature Commun. Math. Phys. 34 (1973), 271-296.*
- [34] Gruber. C, Hintermann. A, and Merlini. D, *Analyticity and uniqueness of the invariant equilibrium state for general spin 1/2 classical lattice spin systems. Commun. Math. Phys. 40 (1975), 83-95.*
- [35] Griffiths. R. B, *Rigorous results for Ising ferromagnets of arbitrary spin. J. Math. Phys. 10 (1969), 1559-1565.*
- [36] Simon. B and Griffiths. R. B, *The $(\Phi^4)_2$ Field theory as a classical Ising model. Commun. Math. Phys. 33, (1973), 145-164.*
- [37] Newman. C. M, *Zeros of the partition function for generalized Ising systems. Commun. Pure. Appl. Math. 27, (1974), 143-159.*
- [38] Dunlop. F, *Zeros of the partition function and gaussian inequalities for the plane rotator model. J. Stat. Phys. 21 (1979), 561-572.*
- [39] Dunlop. F, *Analyticity of the pressure for Heisenberg and plane rotor models. Commun. Math. Phys. 69 (1979), 81-88.*
- [40] Lieb. E and Sokal. A. D, *A general Lee-Yang theorem for one-component and multicomponent ferromagnets. Commun. Math. Phys. 80 (1981), 153-179.*
- [41] Glimm. J and Jaffe. A, *Quantum Physics. A functional integral point of view. New York ect. Springer (1981)*
- [42] Bricmont. J, Lebowitz. J. L and Pfister. C. E, *Low temperature expansion for continuous spin Ising models. Commun. Math. Phys. 78 (1980), 117-135.*
- [43] Dobrushin. R. L, *Induction on volume and no Cluster expansion. In: M. Mebkhout and R. Seneor (eds), VIII. Internat. Congress on Mathematical Physics, Marseille 1986, Singapore: World Scientific, pp. 73-91.*

- [44] Dobrushin. R. L and Sholsmann. S. B, *Completely analytical Gibbs fields.* In: J. Fritz, A.Jaffe, and D.Szász (eds) *Statistical Mechanics and Dynamical Systems*, Boston ect. Birkhäuser, (1985), pp. 371-403.
- [45] Dobrushin. R.L and Sholsmann. S. B, *Completely analytical interactions: constructive description.* *J. Stat. Phys.* 46 (1987), 983-1014.
- [46] Duneau. M, Iagolnitzer. D and Souillard. B, *Decrease properties of truncated correlation functions and analyticity properties for classical lattice and continuous systems.* *Commun. Math. phys.* 31 (1973), 191-208.
- [47] Duneau. M and Iagolnitzer. D and Souillard. B, *Strong cluster properties for classical systems with finite range interaction* *Commun. Math. Phys.* 35 (1974), 307-320.
- [48] Duneau. M and Iagolnitzer. D and Souillard. B, *Decay of correlations for infinite range interactions.* *J. Math. Phys.* 16 (1975), 1662-1666.
- [49] Glimm. J and Jaffe. A, *Expansion in Statistical Physics.* *Commun. Pure. Appl. Math.* 38 (1985), 613-630.
- [50] Israel. R. B, *High temperature analyticity in classical lattice systems.* *Commun. Math. Phys.* 50 (1976), 245-257.
- [51] Kotecký. R and Preiss. D, *Cluster expansions for abstract polymers models.* *Commun. Math. Phys.* 103, (1986), 491-498.
- [52] Kunz. H, *Analyticity and clustering properties of unbounded spin systems.* *Commun. Math. Phys.* 59 (1978), 53-69.
- [53] Lebowitz. J. L, *Bounds on the correlations and analyticity properties of Ising spin systems.* *Commun. Math. Phys.* 28 (1972), 313-321.
- [54] Lebowit., J. L, *Uniqueness, analyticity and decay properties of correlations in equilibrium systems.* In: H. Araki (ed) *International Symposium on Mathematical Problems in Theoretical Physics.* LNPH. 80 (1975), pp. 68-80.
- [55] Malyshev. V. A, *Cluster expansions in lattice models of statistical physics and the quantum theory of fields.* *Russian Math Surveys.* 35,2 (1980), 3-53.
- [56] Malyshev. V. A and Milnos. R. A, *Gibbs Random Fields: The method of cluster expansions (In Russian)* Moscow: Nauka (1985).
- [57] Prakash. C, *High temperature differentiability of lattice Gibbs states by Dobrushin uniqueness techniques.* *J. Stat.Phys,* 31 (1983), 169-228.
- [58] Jost. Jürgen, *Riemannian Geometry and Geometric Analysis.* 4th ed Berlin : Springer, c2005.

- [59] Park. Y. M, *Lack of screening in the continuous dipole systems*, *Comm. Math. Phys.* 70 (1979), 161-167.
- [60] Gawedzki. K and Kupiainen. A, *Block spin renormalization group for dipole gas and $(\nabla\phi)^4$* , *Ann. Phys.* (1983), 147-198.
- [61] Brydges. D and Yau. H. T, *Grad ϕ perturbations of massless gaussian fields*, *Comm. Math. Phys.* (1990), 129-351.
- [62] Fröhlich. J and Spencer. T, *On the statistical mechanics of classical Coulomb and dipole gases*, *J. Stat. Phys.* 24 (1981), 617-701.
- [63] Fröhlich. J and Park. Y. M, *Correlation inequalities in the thermodynamic limit for classical and quantum systems*. *Comm. Math. Phys.* 59 (1990), 235-266.
- [64] Marchetti. D. H and Klein. A, *Power law fall-off in the two dimensional Coulomb gases at inverse temperature $\beta > 8\pi$* , *J.Stat.Phys.* 64 (1991), 135.
- [65] Berezin. F. A and Shubin. M. A, *The Schrödinger Equation* (Kluwer Academic Publisher, 1991).
- [66] Lo. Assane, *Witten laplacian methods for the decay of correlations*. Preprint (2006).