

Self-Embeddings of Computable Trees

Stephen Binns, Bjørn Kjos-Hanssen, Manuel Lerman,
James H. Schmerl, and Reed Solomon

Abstract We divide the class of infinite computable trees into three types. For the first and second types, $0'$ computes a nontrivial self-embedding while for the third type $0''$ computes a nontrivial self-embedding. These results are optimal and we obtain partial results concerning the complexity of nontrivial self-embeddings of infinite computable trees considered up to isomorphism. We show that every infinite computable tree must have either an infinite computable chain or an infinite Π_1^0 antichain. This result is optimal and has connections to the program of reverse mathematics.

1 Introduction

In this article, we examine self-embeddings of countable trees from the perspective of computable algebra. The following definition of a tree is more restrictive than some other definitions in the literature but it is more general than the notion of computable tree used in the context of Π_1^0 classes.

Definition 1.1 A *tree* is a partial order (T, \preceq) with a least element (called the *root* of T and denoted by λ) such that for all $n \in T$, the set $\{m \in T \mid m \preceq n\}$ is a finite linearly ordered set. The elements of T are referred to as *nodes*. If $n \prec m$ and there are no elements strictly between n and m in T , we say that m is a *successor* of n . If n has no successor, then it is called a *leaf* and if n has more than one successor, then it is called a *branching node*. If (T_0, \preceq_0) and (T_1, \preceq_1) are trees, then an *embedding* from T_0 to T_1 is an injective function $f : T_0 \rightarrow T_1$ such that $n \preceq_0 m$ if and only if $f(n) \preceq_1 f(m)$. We write $f : T_0 \hookrightarrow T_1$ to denote that f is an embedding of T_0 into T_1 .

Our concern is with countable trees, so we assume $|T| \leq \omega$ for the rest of this article. The *branching function* $br : T \rightarrow \omega \cup \{\infty\}$ of T is the function which maps $n \in T$

Received April 21, 2006; accepted June 8, 2007; printed December 3, 2007
2000 Mathematics Subject Classification: Primary, 03D45; Secondary, 03C57
Keywords: effective algebra, reverse mathematics, self-embedding, tree
© 2007 by University of Notre Dame 10.1215/00294527-2007-001

to the number of successors of n . Notice that n is a leaf if and only if $\text{br}(n) = 0$. T is *finitely branching* if the range of the branching function is contained in ω and T is *binary branching* if the range of the branching function is contained in $\{0, 1, 2\}$.

Classically, any countable tree T is isomorphic to a subtree of $\omega^{<\omega}$. A tree (T, \preceq) is *computable* if $T \subseteq \omega$ is a computable set and \preceq is a computable relation defined on T^2 . (It does not hurt to assume that a computable tree is coded in such a way that $T = \omega$.) For example, if T is a computable nonempty subset of $2^{<\omega}$ or $\omega^{<\omega}$ which is closed under initial segments and \preceq is the initial segment relation, then (T, \preceq) is a computable tree. In these cases, the successor relation is computable and in the case when $T \subseteq 2^{<\omega}$, the leaf relation and the branching function are computable. However, in general, it is not the case that the successor relation, the leaf relation, or the branching function is computable for a computable tree or that a computable tree is computably isomorphic to a computable subtree of $\omega^{<\omega}$.

A large amount of work has been done on Π_1^0 classes, which are sets of infinite paths through computable subtrees of $2^{<\omega}$ and $\omega^{<\omega}$. (See Cenzer [1] and Cenzer and Remmel [2] for surveys of this work.) In addition, work has been done on the possible degrees of isomorphism types of trees by Richter [12] and on computable categoricity of trees by R. Miller [11] and by Lempp, McCoy, R. Miller, and Solomon [10]. One of the main tools for working with trees in [10], [11], and [12] and in our current work is Kruskal's Lemma.

Lemma 1.2 (Kruskal [9]) *Let $\{T_i \mid i \in \omega\}$ be a countable collection of finite trees. There exists $k \in \omega$ such that for all $i > k$, there are infinitely many $j > i$ for which T_i embeds into T_j .*

The main motivation for the present work comes from the effective analysis of the Dushnik-Miller Theorem [6]. This theorem states that any countably infinite linear order has a nontrivial self-embedding. (A self-embedding of a linear order is called nontrivial if it moves at least one, and hence infinitely many, points.) Downey and Lempp [5] analyzed the classical proof of the Dushnik-Miller Theorem and observed that if L is a computable infinite linear order then $0''$ computes a nontrivial self-embedding of L . It remains an open question whether there is such an order which requires $0''$ to compute a nontrivial self-embedding, but they showed that there is a computable linear order L for which any nontrivial self-embedding computes $0'$.

Downey, Jockusch, and J. S. Miller [4] carried this analysis further and showed that for computable discrete linear orders,¹ being PA over $0'$ is enough to compute a nontrivial self-embedding and conversely that there is a computable discrete linear order for which every nontrivial self-embedding is PA over $0'$.² They also proved that there is an infinite nondiscrete computable linear order for which $0'$ cannot compute a nontrivial self-embedding.

Our goal is to carry out a similar analysis for nontrivial self-embeddings of computable trees. In the context of trees, it is useful to take a slightly more restrictive definition of nontrivial. We say that a self-embedding $f : T \rightarrow T$ is *nontrivial* if it is not an onto map. (It follows from the proof of the Dushnik-Miller Theorem that every countably infinite linear order has a self-embedding which is nontrivial in this sense as well.) We say that a self-embedding f is *weakly nontrivial* if it is not the identity map. In the context of trees (as opposed to linear orders), this condition does not imply that f must move infinitely many nodes. However, frequently one can build trees for which any weakly nontrivial self-embedding moves infinitely many points.

We will state our results for nontrivial self-embeddings and occasionally point out cases in which they can also be applied to weakly nontrivial self-embeddings.

It is known that every countably infinite tree has a nontrivial self-embedding and such an embedding can be obtained by a simple application of Kruskal's Lemma. (For example, see Ross [13]. This existence result also holds for more general definitions of trees using extensions of Kruskal's Lemma such as the main theorem from Corominas [3].) In Section 2, we analyze such a proof and show that $0''$ computes a nontrivial self-embedding for any computable infinite tree T . This proof naturally breaks into three cases depending on the structure of T and we use this distinction to define three classes of trees.

If n is a node in T , we let $T(n)$ denote the subtree $\{m \mid n \preceq m\}$ with the inherited partial order. We say that n is an *infinite node* if $T(n)$ is infinite and we say that n is an *ω -node* if n has infinitely many successors. A *path* in T is an infinite maximal linearly ordered subset of T . We say that a path $P \subseteq T$ is *isolated* if there is a node $n \in P$ such that P is the only path containing n . T is called a *type 1 tree* if it contains a maximal infinite node. That is, T contains a node n such that $T(n)$ is infinite but for all $m \succ n$, $T(m)$ is finite. T is called a *type 2 tree* if it does not have a maximal infinite node but does have an isolated path. T is called a *type 3 tree* if it is infinite but has no maximal infinite node and no isolated paths. (R. Miller [11] used the same classification of height ω trees as well as proof techniques similar to several of those used here.) We show that if T is an infinite computable tree of type 1 or type 2, then $0'$ can compute a nontrivial self-embedding of T and that if T is an infinite computable type 3 tree, then $0''$ can compute a nontrivial self-embedding of T .

There are at least two questions that one might ask concerning the optimality of these results. First, is there a computable type 1 (or type 2) tree T for which every nontrivial self-embedding computes $0'$? Second, is there a computable type 1 (or type 2) tree T such that for every (classically) isomorphic computable tree $S \cong T$ and for every nontrivial self-embedding f of S , f computes $0'$? (There are similar questions concerning the connection between $0''$ and nontrivial self-embeddings of type 3 trees.) When answering the first question, one is allowed to use facts about the particular computable coding (or presentation) of T , while in the second question, one must work with the isomorphism type of T and not with the particular coding. We examine both questions with respect to each type of trees.

In Section 3, we show that there is a computable type 1 tree for which every nontrivial self-embedding computes $0'$ and there is a computable type 1 tree for which no (classically) isomorphic computable tree has a computable nontrivial self-embedding. In Section 4, we show the same results for computable type 2 trees. In Section 5, we show that there is a computable type 3 tree for which every nontrivial self-embedding computes $0''$ and there is a computable type 3 tree such that no (classically) isomorphic computable tree has a $0'$ -computable nontrivial self-embedding. (We actually show something slightly stronger but we leave the technical statement of the result until Section 5.)

These results show that the bounds of $0'$ and $0''$ are optimal in the sense of specific computable trees but we only obtain partial results in terms of the isomorphism types. It remains an open question whether there exists a type 1 or 2 computable tree for which $0'$ is necessary to compute a nontrivial self-embedding in every isomorphic computable copy and whether there exists a type 3 computable tree for

which $0''$ is necessary to compute a nontrivial self-embedding in every isomorphic computable copy.

In Section 6, we turn to a slightly different question. The Chain/Anti-Chain Principle states that every infinite partial order has either an infinite chain or an infinite antichain. (This principle is a simple consequence of Ramsey's Theorem for pairs.) Herrmann [7] investigated this result from the perspective of computable combinatorics and proved that every infinite computable partial order has either an infinite Δ_2^0 chain or an infinite Π_2^0 antichain. Furthermore, he proved that these bounds were optimal by constructing an infinite computable partial order which has no infinite Σ_2^0 chains or antichains.

We examine how these results can be improved in the context of trees as a special type of partial order. We show that every infinite computable tree has either an infinite computable chain or an infinite Π_1^0 antichain. Furthermore, we show that these bounds are optimal by constructing an infinite computable tree which has no infinite Σ_1^0 chains or antichains. Our construction is easily modified to work in the context of models of the subsystem WKL_0 of second-order arithmetic. Thus we show that WKL_0 is not strong enough to prove the Chain/Anti-Chain Principle for binary branching trees.

Our computability theoretic notation is standard and follows Soare [15]. In particular, we use φ_e to denote the e th partial computable function, we use K to denote the halting set (or any other complete computably enumerable set), and we use $X[n]$ to denote $\{m \in X \mid m < n\}$ for any set X . The relation \leq denotes a tree order, \leq denotes the standard order on ω , and \leq_T denotes Turing reducibility.

2 Nontrivial Self-Embeddings

In this section we show that $0''$ suffices to compute a nontrivial self-embedding of any infinite computable tree and that for certain special cases, $0'$ is sufficient. For any tree T , let $S(m, n)$ denote the successor relation on T (that m is a successor of n) and let $\text{br} : T \rightarrow \omega \cup \{\infty\}$ denote the branching function on T . As mentioned in the introduction, the successor relation and the branching function need not be computable even if T is computable.

Lemma 2.1 *Every computable tree T embeds into $2^{<\omega}$ and the embedding is computable in $0'$.*

Proof Given an arbitrary $n \in T$ we compute the image of n as follows. $0'$ computes the successor relation of T (which is explicitly Π_1^0) so we can use $0'$ to compute the sequence

$$\lambda = n_0 < n_1 < \dots < n_k = n,$$

where $S(n_{i+1}, n_i)$ for all i . The image of n will then be

$$1^{n_0} * 0 * 1^{n_1} * 0 * \dots * 1^{n_k} * 0.$$

(where 1^m denotes the string of m ones and $*$ denotes concatenation). It is straightforward to confirm that this gives an embedding. (Recall that an embedding does not need to be closed under initial segments.) \square

We defined a path in T to be an infinite maximal linearly ordered subset of T . Often we specify a path by giving an infinite sequence of elements $x_0 < x_1 < x_2 < \dots$ such that $S(x_{i+1}, x_i)$. (This method of specifying a path corresponds to the notion of

path used in the study of Π_1^0 classes in which the successor relation is computable.) From such a sequence, we can compute the path $P = \{y \mid \exists i(y \preceq x_i)\}$ by calculating the elements x_i in order until we find an x_i such that either $y \preceq x_i$ or y is incomparable with x_i . This procedure cannot necessarily be reversed; in general, we cannot compute such a sequence from a given path P because the successor relation need not be computable from P . However, we can compute a sequence of elements $y_0 < y_1 < \dots$ which is cofinal in P without requiring that $S(y_{i+1}, y_i)$ holds. Furthermore, from such a sequence we can also compute the associated path.

Lemma 2.2 *If T is an infinite tree and it has no maximal infinite node then it has a path.*

Proof Suppose T is an infinite tree and has no maximal infinite node. If T has no ω -node then it is finitely branching and has a path by König's Lemma. Otherwise, T must have an ω -node n_0 and a successor n_1 of n_0 such that $T(n_1)$ is infinite (otherwise n_0 would be a maximal infinite node). If $T(n_1)$ has no ω -nodes, then $T(n_1)$ has a path (and so does T) by König's Lemma. Otherwise, $T(n_1)$ must have an ω -node n_2 which has a successor n_3 such that $T(n_3)$ is infinite. Iterating in this way, we either arrive at an infinite finitely branching tree $T(n_i)$ which has a path, or we obtain an infinite sequence $n_0 < n_1 < n_2 < \dots$ which defines a path. \square

Theorem 2.3 *Every infinite computable tree has a nontrivial self-embedding computable in $0''$.*

The proof splits (nonuniformly) into three cases depending on whether the infinite computable tree is of type 1, 2, or 3. By Lemma 2.2, if T is infinite but not of type 1, then T must have a path, so either T has an isolated path (and is a type 2 tree) or T has no isolated path (and is a type 3 tree). Therefore, every infinite tree is either type 1, 2, or 3. The next three lemmas cover these cases and show that for computable trees of type 1 or 2, $0'$ computes a nontrivial self-embedding.

Lemma 2.4 *Every computable type 1 tree has a nontrivial self-embedding computable in $0'$.*

Proof Let T be a computable type 1 tree with maximal infinite node n . Because $T(n)$ is infinite but $T(m)$ is finite for all $m \succ n$, n must be an ω -node. Let $n_0 < n_1 < \dots$ be the successors of n . (These are computable from $0'$.) By Kruskal's Lemma, there is a k such that

$$\forall m \geq k \exists^\infty s \geq m (T(n_m) \hookrightarrow T(n_s)).$$

Fix such a k . We define a $0'$ -computable embedding φ such that for all m , $\varphi(m) \neq m$ if and only if $m \geq n_l$ for some $l \geq k$. Furthermore, for all $l \geq k$, $\varphi(n_l) = n_j$ for some $j > l$.

To define φ , we set $\varphi(m) = m$ for all $m \in T$ such that $n \not\preceq m$ or $m = n$ or $n_i \preceq m$ for some $i < k$. We define φ on the subtrees $T(n_j)$ for $k \leq j$ by induction on j . Fix $j \geq k$ and suppose that φ has been defined on $T(n_i)$ for all $i < j$. We let T_s denote the subtree formed by restricting \preceq to $\{0, 1, \dots, s\}$. Use $0'$ to find an s and t such that

1. $n_t > \max\{\varphi(n_i) : i < j\} + 1$ (the max is taken with respect to \leq),
2. $T_s(n_j)$ embeds into $T_s(n_t)$,
3. $\forall s' > s \ T_{s'}(n_j) = T_s(n_j)$.

Such an s and t exist by our choice of k and because n is a maximal infinite node and hence $T(n_i)$ is finite for each i . We extend φ to include the embedding of $T(n_j) = T_s(n_j)$ into $T_s(n_t)$. Because $\varphi(n_i) = n_i$ for $i < k$, we have that for all $j \geq k$, if $\varphi(n_j) = n_t$, then $t > k$. Therefore, n_k is not in the range of φ , so φ is nontrivial. \square

If n is a successor of the root of a tree T , then we say $T(n)$ is a *successor tree* of T . Similarly, if n is a successor of m in T , then we say that $T(n)$ is a *successor tree of* $T(m)$.

Lemma 2.5 *Every computable type 2 tree has a nontrivial self-embedding computable in $0'$.*

Proof Let T be a computable type 2 tree. By definition, T has no maximal infinite node and has an isolated path X . Fix $i \in T$ such that there is only one infinite path extending i . By Lemma 2.2, if $j \succ i$ is any node on X there is exactly one successor j' of j such that $T(j')$ is infinite, namely, the successor that is on X . Therefore, the only possible ω -nodes extending i lie on X . If $m \succ i$ is an ω -node, all but one of its successor trees are finite and we are essentially in the case of a type 1 tree. (Let $n_0 < n_1 < \dots$ be the successor nodes of m and let l be such that $T(n_l)$ is the only infinite successor tree. Apply the argument in the proof of Lemma 2.4 to the sequence of successor nodes $n_{l+1} < n_{l+2} < \dots$.) Therefore, we assume there are no ω -nodes above i .

We compute X from $0'$ by computing the sequence $i = x_0 \prec x_1 \prec x_2 \prec \dots$ such that $S(x_{i+1}, x_i)$ and each $x_i \in X$. Suppose we have calculated x_j . We use $0'$ to find a stage s such that

$$\forall n \succ x_j \exists m \in T_s(x_j) (m \neq x_j \text{ and } n \succeq m).$$

Such an s must exist as $T(x_j)$ is finitely branching by our assumption. The finite number of nodes which appear to be the successors of x_j at stage s are the actual successors of x_j . Again using $0'$, search for a successor x' of x_j and a t such that

$$\forall v \geq t [v \succ x_j \implies v \succeq x'].$$

Such x' and t must exist as x_j has exactly one path through it. Since x' is the successor of x_j on X , we set $x_{j+1} = x'$.

We have defined (from $0'$) the sequence $i = x_0 \prec x_1 \prec x_2 \prec \dots$ such that for all i , x_i is on the path X and $S(x_{i+1}, x_i)$. We now can apply Kruskal's theorem to the sequence of finite trees $T(x_i) \setminus T(x_{i+1})$ and use an argument similar to the proof of Lemma 2.4. \square

Lemma 2.6 *Every computable type 3 tree has a nontrivial self-embedding computable in $0''$.*

Proof Let T be a computable type 3 tree. By definition, T is infinite but has no maximal infinite node and no isolated path. We use $0''$ to define an embedding $\varphi : 2^{<\omega} \hookrightarrow T$ by recursion (described below) and then use $0'$ to define an embedding $\beta : T \hookrightarrow 2^{<\omega}$ as in Lemma 2.1. The composition $\alpha = \varphi \circ \beta$ is the desired nontrivial self-embedding $\alpha : T \hookrightarrow T$. (There are numerous ways to see that the self-embedding α is nontrivial. The empty sequence \emptyset , which is the root of $2^{<\omega}$, is not mapped to the root of T by φ and hence φ is not onto. Also, the map β

from Lemma 2.1 is not onto. Either of these facts is enough to conclude that α is nontrivial.)

We define the embedding $\varphi : 2^{<\omega} \rightarrow T$ by recursion using $0''$. Let m be any node other than the root of T for which $T(m)$ is infinite. By our case assumption, $T(m)$ has no maximal infinite nodes and no isolated paths. Let $\varphi(\emptyset) = m$. Assume that $\varphi(\tau)$ has been defined for all τ such that $|\tau| \leq k$, that $T(\varphi(\tau))$ is infinite and that φ gives an embedding of $2^{\leq k}$ into T . Consider each σ with $|\sigma| = k$ separately and we show how to define $\varphi(\sigma * 0)$ and $\varphi(\sigma * 1)$. Assume $\varphi(\sigma) = n$. Using $0''$ find two incomparable extensions of n , say n_0 and n_1 , such that $T(n_0)$ and $T(n_1)$ are both infinite. Because $T(n)$ is infinite and T has no isolated paths or maximal infinite nodes, such nodes n_0 and n_1 must exist. Set $\varphi(\sigma * 0) = n_0$ and $\varphi(\sigma * 1) = n_1$. It is easy to check that the inductive assumptions hold at level $k + 1$. \square

Theorem 2.3 gives an analysis of the existence of nontrivial self-embeddings in terms of the jump hierarchy. We could also ask for an analysis in terms of other computable relations on T . That is, are there natural algebraic relations on T such that we can compute a nontrivial self-embedding from these relations? For computable trees of type 1 or 2, there is a nontrivial self-embedding computable from the join of the successor relation and the branching function.

Corollary 2.7 *If T is a computable type 1 tree, then T has a nontrivial self-embedding computable from the join of the successor relation and the branching function.*

Proof We used $0'$ twice in the proof of Lemma 2.4. First, we used $0'$ to determine the successors of n . Clearly, we can determine these successors from the successor relation. Second, we used $0'$ to determine if $\forall s' > s (T_{s'}(n_j) = T_s(n_j))$. That is, we used it to find a stage by which the finite tree $T(n_j)$ had stopped growing. Since the trees $T(n_j)$ are finite, we can also determine such a stage from the successor relation together with the branching function. \square

Corollary 2.8 *If T is a computable type 2 tree, then T has a nontrivial self-embedding computable from the join of the successor relation and the branching function.*

Proof Consider the proof of Lemma 2.5. If there is an ω -node $m \succeq i$, then this proof reduces to the proof of Lemma 2.4 so we are done by Corollary 2.7. Otherwise, we used $0'$ twice in the definition of the sequence $i = x_0 \prec x_1 \prec \dots$. First, we used it to find a stage s such that all the successors of x_j had appeared by stage s . However, if we know the branching function then we can calculate the number of successors of x_j (which in this situation is finite) and we can use the successor relation to find this number of successors. Second, we used $0'$ to determine the unique infinite successor of x_j . Because the successor relation together with the branching function can tell when a finite tree has stopped growing, we can use them to make this determination as well. \square

The same type of argument does not work in the case of a computable type 3 tree. As we will show in Section 5, there is an infinite computable tree T which is finitely branching (and hence has no maximal infinite nodes), has no isolated paths and for which every nontrivial self-embedding computes $0''$. We claim that the branching

function for this tree T is computable from O' . To calculate $\text{br}(n)$, ask O' if there is a node $m \succ n$. If not, $\text{br}(n) = 0$. If so, use O' to find a successor n_0 of n . Ask O' if there is a node $m \succ n$ such that $n_0 \not\leq m$. If not, $\text{br}(n) = 1$. If so, use O' to find a second successor n_1 of n . Because T is finitely branching, this process must eventually stop with a complete set of successors n_0, \dots, n_{k-1} for n . Therefore, O' can compute both the branching function and the successor relation in T . Since every nontrivial self-embedding of T computes O' , there cannot be such a self-embedding computable from the join of the successor relation and the branching function.

3 Type 1 Trees

Recall that a computable type 1 tree is a computable tree which has a maximal infinite node. By Lemma 2.4, O' computes a nontrivial self-embedding for such trees. In this section, we show that this result is optimal in the sense that there is a computable type 1 tree for which every nontrivial self-embedding computes O' . We also show that there is a computable type 1 tree T such that for all computable trees $S \cong T$, S does not have a computable self-embedding.

The *height of a node n* in T (denoted by $\text{ht}(n)$) is the size of the set $\{m \mid m \prec n\}$. For example, the root of any tree has height 0 and the successors of the root have height 1. The *height of a finite tree T* (denoted $\text{ht}(T)$) is the maximum height of a node of T . Frequently, our examples of type 1 trees have an ω branching root λ and have $T(x)$ finite for all $x \neq \lambda$. Recall that for each x of height 1, we say that $T(x)$ is a *successor tree* of T .

Theorem 3.1 *There is a computable type 1 tree T such that any nontrivial self-embedding of T computes O' .*

Proof Our proof will show that even the weakly nontrivial self-embeddings of T compute O' . We describe the tree before explicitly constructing it. 0 will be the root of T and the set of successors of 0 will be the set of positive even numbers \mathbb{E}^+ . Each subtree $T(n)$ with $n \in \mathbb{E}^+$ will be a finite tree with no branching nodes and $\text{ht}(T(n)) \geq n$. (That is, $T(n)$ is a finite linear order of length at least n . The exact length of this order will be determined during the construction.) We construct a sequence of positive even numbers $2 = m_0 < m_1 < m_2 < \dots$ which will have the following properties:

- I. For all $i \in \omega$, $\text{ht}(T(m_i)) < \text{ht}(T(m_{i+1}))$;
- II. For all $i \in \omega$ and $p, q \in \mathbb{E}^+$, if $m_i \leq p < q < m_{i+1}$ then $\text{ht}(T(p)) > \text{ht}(T(q))$ and if $i > 0$, then $\text{ht}(T(q)) > \text{ht}(T(m_{i-1}))$;
- III. Let $K = \bigcup_{s=0}^{\infty} K_s$ be a fixed computable enumeration of a complete c.e. set. For all i , $K[i] = K_{m_i}[i]$. (Recall that for any set X , $X[i] = \{n < i \mid n \in X\}$.)

We picture T as looking like a series of strictly descending staircases. That is, for any i , the subtrees $T(m_i), T(m_i + 2), T(m_i + 4), \dots, T(m_{i+1} - 2)$ are all linear orders which are decreasing in height (but all taller than $T(m_{i-1})$). The subtree $T(m_{i+1})$ jumps up in height (to be taller than $T(m_i)$) and begins another sequence of subtrees of decreasing height (but all taller than $T(m_i)$) which continues until we reach $T(m_{i+2})$. The idea behind this tree is that if $q \in \mathbb{E}^+$ is such that $m_i \leq q < m_{i+1}$, then $T(q)$ does not embed into any subtree of the form $T(p)$ where $p \in \mathbb{E}^+$ satisfies $p < m_i$ or $q < p < m_{i+1}$. (It follows from properties I and II that in both of these cases $\text{ht}(T(q)) > \text{ht}(T(p))$.) Therefore, if δ is a weakly nontrivial self-embedding

which moves the node q , then δ must map $T(q)$ into a subtree $T(p)$ such that $p \in \mathbb{E}^+$ and either $m_i \leq p < q$ or $m_{i+1} \leq p$. In the former case, by iterating δ we must arrive at some $k \geq 1$ such that δ^k maps $T(q)$ into $T(p)$ where $p \in \mathbb{E}^+$ and $p > q$ (and so $p > m_{i+1}$ by the previous comments).

Properties I, II, and III are sufficient to guarantee that any weakly nontrivial self-embedding φ computes K . Because φ is weakly nontrivial, there must be an $n \in \omega$ such that $\varphi(n) \neq n$. Fix any such n . For all $m > 0$ let $\lfloor m \rfloor$ be the unique element of \mathbb{E}^+ such that $m \in T(\lfloor m \rfloor)$. We define a strictly increasing function ψ from φ as follows:

$$\psi(0) = \lfloor n \rfloor,$$

$$\psi(s+1) = \lfloor \varphi^k(n) \rfloor \text{ where } k = k(s+1) \text{ is the least natural number such that } \lfloor \varphi^k(n) \rfloor > \psi(s).$$

(The existence of ψ follows from the comments above about the general form of T .) Once we prove by induction that $\psi(s) \geq m_s$ for all $s \in \omega$, we will have by property III that $K[i] = K_{\psi(i)}[i]$ and hence that $\psi \geq_T K$. Since $\varphi \geq_T \psi$, we have $\varphi \geq_T K$ as claimed.

The base of the induction is $\psi(0) = \lfloor n \rfloor \geq 2 = m_0$. Suppose that $\psi(s) \geq m_s$. Let j be such that $m_j \leq \psi(s) < m_{j+1}$. Then $j \geq s$ as $\langle m_s \rangle$ is an increasing sequence. If $j > s$, then we are done as ψ is increasing, so we can assume that $m_s \leq \psi(s) < m_{s+1}$.

Property I ensures that for all l if $\psi(s) < l < m_{s+1}$, then $T(\psi(s)) \not\leftrightarrow T(l)$. Therefore, for all $t \geq k = k(s)$,

$$\lfloor \varphi^t(n) \rfloor > \psi(s) \implies \lfloor \varphi^t(n) \rfloor \geq m_{s+1}.$$

But $k(s+1) > k(s)$ and so in particular $\psi(s+1) \geq m_{s+1}$ as required.

It remains to give the construction of a computable type 1 tree T satisfying I, II, and III. We build T in stages. At stage s we build T^s and T will be $\bigcup_s T^s$. T^0 will consist of all the even nodes as above as well as an infinite/cofinite computable set of odd numbered nodes arranged so that for all $e \in \mathbb{E}^+$, $\text{ht}(T(e)) = e$. (That is, $T(e)$ is a linear chain of length e .)

We use a movable marker argument to create the sequence $\langle m_i \rangle$. As we do this we also ensure that I, II, and III are satisfied. We describe a uniformly computable sequence $\langle m_{i,s} \rangle$ with the properties

- (i) $\forall i (m_{i,0} = 2i)$,
- (ii) $\forall i, s (m_{i,s} < m_{i+1,s})$,
- (iii) $\forall i, s (m_{i,s} \leq m_{i,s+1})$,
- (iv) $\forall i (\lim_s m_{i,s} \text{ exists})$.

For each i , m_i is defined to be $\lim_s m_{i,s}$. We enumerate K one element at a time. Suppose s is a stage at which $i \in K_{s+1} \setminus K_s$ and let $k \geq i$ be the smallest number such that $m_{k,s} \geq s+1$. Then we set

$$m_{j,s+1} = \begin{cases} m_{j,s} & \text{if } j < i \\ m_{k+t,s} & \text{if } j = i+t, (t \in \omega). \end{cases}$$

At the same time it is necessary to adjust the subtrees $T^s(e)$ with $e \in \mathbb{E}^+$. We leave all successor trees in T unchanged except perhaps those $T^s(e)$ with $m_{i-1,s} = m_{i-1,s+1} \leq e < m_{k,s} = m_{i,s+1}$ (if $i = 0$ take $m_{i-1} = 2$). To the subtrees

$T(e)$ with $m_{i-1,s} \leq e < m_{i,s+1}$, we add the minimum number of nodes to the top of each subtree, retaining the property that there are no branching nodes and ensuring that properties I and II are preserved.

The argument that the tree T and the sequence $\langle m_i \rangle$ have the required properties is now just the typical movable marker argument, made explicit in the following lemmas.

Lemma 3.2 *For every j , $\lim_s m_{j,s} = m_j$ exists.*

Proof Let s be such that $K_s[j] = K[j]$. Then for all $t \geq s$, if $i \in K_{s+1} \setminus K_s$, then $i > j$, so for all $t \geq s$, $m_{j,t} = m_{j,t+1}$. \square

Lemma 3.3 *Every successor tree $T(e)$ with $e \in \mathbb{E}^+$ is finite.*

Proof Fix $e \in \mathbb{E}^+$. Let i be such that $m_i > e$. Once $m_{i,s}$ reaches its limit m_i , $T^s(e)$ will never grow again. At each stage before this limit is reached, at most finitely many elements are added to $T(e)$. Therefore, $T(e)$ is finite. \square

Lemma 3.4 *T has properties I, II, and III.*

Proof The fact that T has properties I and II follows immediately from the fact that these properties are explicitly retained at each stage in the construction and the previous two lemmas.

To see that T satisfies property III, fix any $j \in \omega$. If s is the largest stage at which $i \in K_{s+1} \setminus K_s$ for some $i \leq j$, then by the construction,

$$m_j = m_{j,s+1} \geq m_{i,s+1} = m_{k,s} \geq s + 1,$$

where k is as in the construction above. But $K_{s+1}[j] = K[j]$ by our choice of s , so $K_{m_j}[j] = K[j]$. \square

This completes the proof of Theorem 3.1. \square

The proof of Theorem 3.1 leads to several corollaries concerning the connection between the complexity of nontrivial self-embeddings of computable type 1 trees T and the complexity of natural algebraic relations on T .

Corollary 3.5 *There is a computable type 1 tree T such that T has a computable successor relation and every nontrivial self-embedding of T computes $0'$.*

Proof The tree T constructed in Theorem 3.1 has a computable successor relation because we only add nodes above the top node in $T(e)$ at any given stage. Therefore, if n, m are nodes in T at stage s , then $S(n, m)$ holds if and only if it holds at stage s . (Because each nontrivial self-embedding of T computes $0'$ and there is such an embedding computable from the join of the successor function and the branching function, the branching function for T has degree $0'$.) \square

Corollary 3.6 *There is a computable type 1 tree T such that T has a computable branching function and every nontrivial self-embedding of T computes $0'$.*

Proof This corollary follows by altering the construction in Theorem 3.1 such that any new nodes which are added to $T(e)$ at stage s are placed between the node e (the root of $T(e)$) and its current successor. By making this change, the nodes which are leaves at stage 0 remain leaves throughout the rest of the construction. Therefore,


the branching function is equal to ∞ for the root of T and is equal to 1 for all other nodes except those nodes which are leaves at stage 0. (By reasoning similar to the parenthetical remark at the end of Corollary 3.5, the successor relation has degree 0' for this tree.) \square

We next turn to the question of working with nontrivial self-embeddings for isomorphism types of computable type 1 trees.

Theorem 3.7 *There is a computable type 1 tree T such that no computable tree classically isomorphic to T has a computable nontrivial self-embedding.*

The remainder of this section is dedicated to the proof of Theorem 3.7. We will build a computable tree T such that the root of T is the only ω -node. For each successor n of the root, we call the subtree $T(n)$ a *component* of T and we will make each component finite. Because the root of T will be the only infinite node, T will have the required form.

To build T , we uniformly construct the sequence T_0, T_1, \dots of components of T . (These components should also have subscripts indicating the current stage of the construction but we suppress the stage subscript unless it is not clear from context.) Each component T_i will have height three and will consist of finitely many components each of which will be one of the following four types.



More specifically, each T_i will contain at most one component of type A, at least one component of type B, exactly i many components of type D, and finitely many components (including possibly none) of type C. Consider the form of a nontrivial self-embedding δ of T . We let λ denote the root of T and r_i denote the root of the component T_i . Because T has height four, $\delta(\lambda) = \lambda$. Because each T_i has height three, $\delta(r_i) = r_j$ for some j . Because T_i has exactly i many type D trees, $\delta(r_i) = r_j$ for some $j \geq i$. Finally, because each T_i is finite, if δ is nontrivial, then there must be some i for which $\delta(r_i) = r_j$ for $j > i$. By considering iterated images of T_i , it is clear that there must be infinitely many indices i for which δ maps T_i into T_j for some $j > i$. Therefore, any nontrivial self-embedding of (any isomorphic copy of) T must map infinitely many components into disjoint components. We will exploit this property in our proof.

Fix two effective enumerations of the partial computable functions: φ_e and f_i . (We use two different notations to distinguish between the partial computable function which we view as determining the e th tree and the partial computable function which we view as giving a potential nontrivial self-embedding of this tree.) We satisfy all requirements of the form

$$R_{(e,i)} : \varphi_e \text{ does not compute a tree isomorphic to } T$$

or f_i is not a nontrivial self-embedding of the tree computed by φ_e .

To make these requirements precise, we need to explain how we obtain a tree from φ_e . We view φ_e as defining a partial computable relation \leq_e on universe ω by setting $n \leq_e m$ if $\varphi_e(\langle n, m \rangle)$ converges to 0 and $n \not\leq_e m$ if $\varphi_e(\langle n, m \rangle)$ converges to a value

other than 0. If φ_e is total and (ω, \leq_e) is a tree, then we refer to this tree as the φ_e -tree. At a stage s , we consider the largest set X such that for all $n, m \in X$, $\varphi_{e,s}(\langle n, m \rangle)$ converges and call $(X, \varphi_{e,s})$ the φ_e -tree at stage s . (Of course, this finite structure may have already violated the axioms for a tree in which case φ_e does not compute a tree and we get an easy win.)

Because each component T_i of our tree T will contain a type B tree, we know that T has height four and each component T_i has height three. Therefore, if the φ_e -tree at stage s has height greater than four, we know it is not isomorphic to our tree T . Furthermore, in the case when the φ_e -tree is isomorphic to our tree T , we can identify the root of the φ_e -tree and the roots of each of its components as it is enumerated. When we say that S is a component of the φ_e -tree at stage s , we mean that the φ_e -tree at stage s has height four, that the least element of S is at level 1 in the φ_e -tree, that S contains a type B component, and that S contains all the nodes (currently) above its least element.

For any component S of the φ_e -tree, we can count the number of type D trees occurring in S . As the component S grows, this number can never decrease. (If we ever see a component of S which contains a component larger than a type D tree, we know the φ_e -tree is not isomorphic to our tree and we get an easy win.) Finally, if the component S has i many type D trees at stage s , then the only possible (current) image of S in T is the component T_i . As the number of type D trees in S grows, the possible image of this component in our tree changes, but if the φ_e -tree is isomorphic to T , then this number must eventually stop growing. Therefore, we eventually correctly guess the only possible image of the component S in our tree.

Before giving the full construction, we consider how to satisfy a single requirement $R_{(e,i)}$ in isolation. Assume that (ω, φ_e) is a tree of height four (so that we do not get an easy win). We wait for a stage s at which the φ_e -tree contains disjoint components U and V such that f_i is defined on all of U and embeds U into V . (By our analysis of nontrivial self-embeddings of trees isomorphic to T , if the φ_e -tree is isomorphic to T and f_i is a nontrivial self-embedding of the φ_e -tree, then such components must exist.) Once we find such components, we fix them and define two parameters. For all stages $t \geq s$, $u(t)$ is equal to the number of type D trees in U at stage t and $v(t)$ is equal to the number of type D trees in V at stage t . (We also assume that neither U nor V ever adds any additional elements to a type D tree. Such additional elements would again give us an easy win. This assumption means that $u(t)$ and $v(t)$ are increasing in t .) For any stage $t \geq s$, we know that $T_{u(t)}$ is the only possible (current) image of U in T and that $T_{v(t)}$ is the only possible (current) image of V in T .

At stage s , $u(s) < v(s)$ and f_i gives an embedding of U into V . We *set up to diagonalize* by taking the following two steps. First, if $T_{u(s)}$ does not have a type A component, then we add elements to $T_{u(s)}$ to create a type A component in $T_{u(s)}$. Second, if $T_{v(s)}$ does have a type A component, then we add an element to $T_{v(s)}$ to change this component from type A to type C. (Recall that $T_{u(s)}$ and $T_{v(s)}$ can have at most one type A component.)

At each stage $t > s$, we calculate $u(t)$ and $v(t)$ and check whether either of these parameters have changed. Assume for the moment that neither of these parameters changes at a future stage. We next check whether at stage t , U is isomorphic to $T_{u(t)}$ and V is isomorphic to $T_{v(t)}$. If not, then we go on to the next stage. If so, we check

whether f_i is defined on all of U and is an embedding on U into V . If not, go on to the next stage. If so, then we are ready to diagonalize.

In this situation, we have $U \cong T_{u(t)}$, $V \cong T_{v(t)}$ and $f_i : U \hookrightarrow V$. $T_{u(t)}$ and U each have a single type A component (because of our set up to diagonalize procedure) and we refer to these components as the *designated components* of $T_{u(t)}$ and U . We refer to the component in V which is the image of the designated component in U as the designated component in V . Because of our set up to diagonalize procedure, we know that the designated component in V is either of type B, C, or D. If the designated component of V is of type B, then we add an element to $T_{u(t)}$ to change the designated component of $T_{u(t)}$ from type A to type C. If the designated component of V is of type C or D, then we add an element to $T_{u(t)}$ to change the designated component in $T_{u(t)}$ from type A to type B.

Consider what can happen after this diagonalization step. If the opponent does not change $u(t)$ at a later stage, then the only way for the φ_e -tree to be isomorphic to T is for U to change its designated component (currently of type A) into the same type of component as the designated component in $T_{u(t)}$ (which is now either type B or C). However, we chose the new type for the designated component in $T_{u(t)}$ in such a way that f_i cannot be extended to map from this type of component into the designated component in V . Therefore, unless the opponent adds new type D components to U to change $u(t)$ or adds new type D components to V to change $v(t)$, we win requirement $R_{(e,i)}$. If the opponent does change $u(t)$ or $v(t)$, then we repeat this process of setting up to diagonalize and later diagonalizing.

To be more specific about this process and give an indication of the full construction, we remove the earlier assumption that $u(t) = u(s)$ and $v(t) = v(s)$ for all $t \geq s$. At each stage $t > s$, we calculate $u(t)$ and $v(t)$ and check whether $u(t) = u(t-1)$ and $v(t) = v(t-1)$. If $u(t-1) < u(t)$, then U has gained extra type D trees and its potential image in T has changed. In this situation in the full construction, we will take an outcome on a tree of strategies indicating that $u(t)$ may be approaching infinity in the limit. Notice that if $u(t)$ does go to infinity in the limit, then U is an infinite component of the φ_e -tree and we win $R_{(e,i)}$ because T has no infinite components. Similarly, if $u(t) = u(t-1)$ but $v(t-1) < v(t)$, then in the full construction, we will take an outcome indicating that $v(t)$ may be approaching infinity in the limit. Again, if $v(t)$ goes to infinity in the limit, then V is an infinite component of the φ_e -tree and we win $R_{(e,i)}$.

If $u(t) = u(t-1)$ and $v(t) = v(t-1)$ and $f_i : U \hookrightarrow V$ at stage t , then we set up diagonalize as above. That is, we add a type A component to $T_{u(t)}$ (if it does not already have one) and change the type A component in $T_{v(t)}$ (if it has one) to a type C component. We check if $U \cong T_{u(t)}$ and $V \cong T_{v(t)}$. If so, then we diagonalize as above.

There are four possible outcomes of this strategy to meet $R_{(e,i)}$ ordered from highest to lowest priority by $u_\infty <_L v_\infty <_L fin <_L triv$. The tree of strategies consists of all finite sequences of these outcomes ordered lexicographically using the $<_L$ order. We use the trivial outcome $triv$ before we have defined the components U and V and if the φ_e -tree has height greater than four or enumerates a component S which contains a component larger than a type D tree. (If we always take this outcome, then we win $R_{(e,i)}$ because of the form of T and the form of the nontrivial self-embeddings of any tree isomorphic to T .) We use the u_∞ outcome whenever the parameter $u(s)$ increases. (If we take this outcome infinitely often, we win $R_{(e,i)}$)

because U is an infinite component in the φ_e -tree.) We use the v_∞ outcome whenever the parameter $u(s)$ retains its value but the parameter $v(s)$ increases. (If we take this outcome infinitely often, then we win because V is an infinite component in the φ_e -tree.) We use the outcome *fin* when both parameters $u(s)$ and $v(s)$ stay the same. (If we take the u_∞ and v_∞ outcomes only finitely often and the *fin* outcome infinitely often, then we win $R_{\langle e, i \rangle}$ by the diagonalization process.)

We will use α and β to denote nodes on the tree of strategies. If $|\alpha| = \langle e, i \rangle$, then α works on requirement $R_{\langle e, i \rangle}$ and we use φ_α and f_α to denote φ_e and f_i . We denote the components chosen by α in the φ_α -tree by U_α and V_α (which also have stage number subscripts which we typically suppress) and we denote the parameters associated with α at stage s by $u(\alpha, s)$ and $v(\alpha, s)$. Each strategy will keep two other parameters $a(\alpha, s)$ and $b(\alpha, s)$ (described below) to deal with the interaction between different strategies. If a strategy α is initialized, then its components U_α and V_α and its parameters become undefined.

To see how strategies for different R requirements interact, assume that $\alpha \subsetneq \beta$. If $\alpha * \text{triv} \subseteq \beta$, then β ignores α when it acts. If $\alpha * \text{fin} \subseteq \beta$, then β assumes that the parameters $u(\alpha, s)$ and $v(\alpha, s)$ have reached their final values and β makes sure that its chosen components U_β and V_β each contain more type D trees than V_α . (That is, β tries to diagonalize using components in T which have indices greater than those used by α to diagonalize.) This restriction causes no problems for β because if the φ_β -tree is isomorphic to T and f_β is a nontrivial self-embedding of the φ_β -tree, then f_β must map infinitely many components of the φ_β -tree to components which contain strictly more type D trees.

If $\alpha * u_\infty \subseteq \beta$, then β assumes that the parameter $u(\alpha, s)$ will approach infinity in the limit. In this case, β waits to work behind α in the sense that β only works with components U_β and V_β which contain strictly fewer type D trees than $T_{u(\alpha, s)}$. If $\alpha * u_\infty$ is on the true path, then $u(\alpha, s)$ will go to infinity, so α may occasionally delay β from acting but will not prevent β from succeeding in the end.

Similarly, if $\alpha * v_\infty \subseteq \beta$, then β assumes that the parameter $u(\alpha, s)$ has reached its limit and that $v(\alpha, s)$ will approach infinity. In this case, β works in between U_α and V_α in the sense that β works with components U_β and V_β for which the number of type D trees is strictly greater than in U_α and is strictly less than in V_α . If $\alpha * v_\infty$ is on the true path, then the parameter $u(\alpha, s)$ eventually reaches a finite value (which β can work beyond in the sense described above) and $v(\alpha, s)$ does go to infinity; so α may cause β to delay acting occasionally, but will not prevent β from succeeding in the end.

To implement these restrictions, we introduce the parameters $a(\beta, s)$ and $b(\beta, s)$. When β is first eligible to act (or first eligible to act after having been initialized), $a(\beta, s)$ is defined to be large. (That is, it is defined to be larger than any number used in the construction so far.) The important feature of $a(\beta, s)$ (which we verify after the construction) is that it is greater than $u(\alpha, s)$ for all α such that $\alpha * v_\infty \subseteq \beta$ and it is greater than $v(\alpha, s)$ for all α such that $\alpha * \text{fin} \subseteq \beta$. When β searches for components U_β and V_β to use in its diagonalization, it only looks at components in the φ_β -tree which have strictly more than $a(\beta, s)$ many type D trees.

At each stage s , β defines its parameter $b(\beta, s)$ to be the minimum of $v(\alpha, s)$ for all α such that $\alpha * v_\infty \subseteq \beta$ and $u(\alpha, s)$ for all α such that $\alpha * u_\infty \subseteq \beta$. When β looks for its components U_β and V_β , it only looks at components in the φ_β -tree which have strictly less than $b(\beta, s)$ many type D trees.

Suppose β is successful at finding components U_β and V_β which satisfy these size restrictions. Because U_β has more than $a(\beta, s)$ many type D trees, β is working with components which are beyond (in the sense of the indices of potentially isomorphic components in T) those used by all α for which $\alpha * \text{fin} \subseteq \beta$. Because V_β has fewer than $b(\beta, s)$ many type D components, β is working with components which are behind (in the same sense as the previous sentence) the components used by each α such that $\alpha * u_\infty \subseteq \beta$. Finally, because U_β has more than $a(\beta, s)$ many type D trees and V_β has fewer than $b(\beta, s)$ many trees, β is working between the witness components for all α such that $\alpha * v_\infty \subseteq \beta$. Thus, these parameters succeed in forcing β to work with components of the intuitively correct size.

We now present the formal construction. At the beginning of stage s , T will contain a root node plus finite components T_0, \dots, T_{s-1} . We begin by adding component T_s consisting of a single type A component, a single type B component and s many type D components. We proceed to let strategies (beginning with the unique strategy for the requirement $R_{(0,0)}$) act as in the basic module described below. Once a strategy α with $|\alpha| = s$ acts, we end the stage and initialize all strategies of lower priority than α . If a strategy is not eligible to act at stage s but is not initialized, then its parameters retain their values.

Basic module for strategy β . When β is first eligible to act (or first eligible to act after being initialized), define $a(\beta, s)$ to be large. This parameter retains its value unless β is initialized. At every β stage, define

$$b(\beta, s) = \min(\{v(\alpha, s) \mid \alpha * v_\infty \subseteq \beta\} \cup \{u(\alpha, s) \mid \alpha * u_\infty \subseteq \beta\})$$

(we take as a convention that the minimum of the empty set is ∞). If U_β and V_β were defined at a previous β stage (since the last initialization of β), then define $u(\beta, s)$ to be the number of type D components in U_β and $v(\beta, s)$ to be the number of type D components in V_β . (Throughout this module, we assume that the φ_β -tree at stage s is a tree of height four and that we have identified its root node. We can identify nodes at level one in the φ_β -tree using type B trees and we assume that no type D component in the φ_β tree ever grows. If any of these conditions are not true, we let $\beta * \text{triv}$ act.)

Step 1. Check if there are disjoint components U_β and V_β in the φ_β -tree such that U_β has $> a(\beta, s)$ many type D components, V_β has $< b(\beta, s)$ many type D components, f_β is defined on all of U_β , and f_β is an embedding of U_β into V_β . If there are no such components, then let $\beta * \text{triv}$ act.

If there are such components, then fix such U_β and V_β . (These components do not change at future stages unless β is initialized.) Define $u(\beta, s)$ and $v(\beta, s)$ as above and *set up to diagonalize*. First, check if $T_{u(\beta, s)}$ has a type A component. If not, then add such a component to $T_{u(\beta, s)}$. If so, then $T_{u(\beta, s)}$ remains unchanged. Second, check if $T_{v(\beta, s)}$ has a type A component. If so, then we add an element to this component to make it into a type C component. If not, then $T_{v(\beta, s)}$ remains unchanged. Let $\beta * \text{fin}$ act. When β is next eligible to act, it acts in step 2 (unless it is initialized).

Step 2. Let t be the previous β stage. Break into the following three subcases. Unless β proceeds to step 3 (or is initialized), β acts in step 2 again at the next β stage.

2(a). If $u(\beta, s) > u(\beta, t)$, then let $\beta * u_\infty$ act.

- 2(b). If $u(\beta, s) = u(\beta, t)$ and $v(\beta, s) > v(\beta, t)$, then let $\beta * v_\infty$ act.
- 2(c). If $u(\beta, s) = u(\beta, t)$ and $v(\beta, s) = v(\beta, t)$, then check if $u(\beta, s) < v(\beta, s) < b(\beta, s)$. If not, then let $\beta * \text{fin}$ act. If so, then set up to diagonalize as described in the second paragraph of step 1. (U_β and V_β are already defined, so setting up to diagonalize consists in making sure that the corresponding trees $T_{u(\beta, s)}$ and $T_{v(\beta, s)}$ have the correct number of A components. Also, once we have set up to diagonalize with a particular choice of $u(\beta, s)$ and $v(\beta, s)$, future setting up procedures will not add elements to $T_{u(\beta, s)}$ and $T_{v(\beta, s)}$ unless these parameters change.) Check if $U_\beta \cong T_{u(\beta, s)}$, $V_\beta \cong T_{v(\beta, s)}$, f_β is defined on all of U_β and gives an embedding of U_β into V_β . If any of these conditions fail, then let $\beta * \text{fin}$ act. If all of these conditions hold, then proceed to step 3.

Step 3. In this step, β diagonalizes as follows. The single type A components in $T_{u(\beta, s)}$ and U_β are called the designated components of $T_{u(\beta, s)}$ and U_β (respectively) and the image of the designated component in U_β under f_β is called the designated component of V_β . If the designated component of V_β is a type B tree, then add an element to the designated component of $T_{u(\beta, s)}$ to make it into a type C tree. If the designated component of V_β is a type C or D tree, then add an element to the designated component of $T_{u(\beta, s)}$ to make it into a type B tree. Let $\beta * \text{fin}$ act. When β is next eligible to act, it acts in step 4.

Step 4. Let $t < s$ be the last β stage. Check if $u(\beta, s) = u(\beta, t)$ and $v(\beta, s) = v(\beta, t)$. If so, then let $\beta * \text{fin}$ act. If not, then return to step 3.

This completes the formal description of the construction. As usual, we say that a strategy α is on the true path if α is the leftmost strategy of length $|\alpha|$ which is eligible to act infinitely often. Because the tree of strategies is finitely branching and because strategies of length up to s get to act at stage s , the true path is infinite.

Lemma 3.8 *Let β be a strategy on the true path.*

1. β is initialized only finitely often.
2. $a(\beta, s)$ reaches a finite limit $a(\beta)$.
3. Unless $\beta * \text{triv}$ is on the true path, β eventually defines the components U_β and V_β permanently. Once these components have been permanently defined, the parameters $u(\beta, s)$ and $v(\beta, s)$ are always defined and are increasing in s . Furthermore, if $\beta * \text{triv}$ acts infinitely often, then $\beta * \text{triv}$ is on the true path.
4. If $\beta * u_\infty$ is on the true path, then $\lim_s u(\beta, s) = \infty$ and U_β has infinitely many type D components.
5. If $\beta * v_\infty$ is on the true path, then $\lim_s v(\beta, s) = \infty$ and V_β has infinitely many type D components.
6. If $\beta * \text{fin}$ is on the true path, then $\lim_s u(\beta, s) = u(\beta)$ and $\lim_s v(\beta, s) = v(\beta)$ both exist, U_β has $u(\beta)$ many type D components (in the limit) and V_β has $v(\beta)$ many type D components (in the limit).
7. $\lim_s b(\beta, s) = \infty$.

Proof The verification of these properties proceeds by induction on β and is standard. For properties 1 and 2, let s be the least β stage such that $s \geq |\beta|$ and the path in the tree of strategies is never to the left of β after stage s . Property 1 follows because β is never initialized after stage s and property 2 follows because β defines $a(\beta, s)$ at stage s and this definition can only be removed by initialization.

For property 3, assume that $\beta * \text{triv}$ is not on the true path and β is not initialized after stage s . In this case, β must eventually move from step 1 to step 2 in the basic module at some stage after s and β defines U_β and V_β at this stage. Because β is not initialized after this time, the only way that β could return to taking outcome $\beta * \text{triv}$ is if the φ_β -tree violated the axioms of a tree or had height greater than four or added an extra element to some type D component in U_β or V_β . In any of these cases, β would take outcome $\beta * \text{triv}$ at all future β stages and hence $\beta * \text{triv}$ would be on the true path. It is clear that if this situation does not occur, then the parameters $u(\beta, t)$ and $v(\beta, t)$ are defined and increasing in t at every subsequent stage. Furthermore, it follows from these comments that if β takes outcome $\beta * \text{triv}$ infinitely often, then $\beta * \text{triv}$ is on the true path.

For property 4, assume that $\beta * u_\infty$ is on the true path and let t be a stage such that β is never initialized after t and $\beta * \text{triv}$ is not eligible to act after t . The strategy $\beta * u_\infty$ is only eligible to act at β stages after t at which the parameter $u(\beta, s)$ has increased. Because this parameter measures the number of type D components in U_β , property 4 follows. The proof of property 5 is essentially the same.

For property 6, assume that $\beta * \text{fin}$ is on the true path. There must be a stage t after which β is never initialized and none of $\beta * \text{triv}$, $\beta * u_\infty$, or $\beta * v_\infty$ are ever eligible to act. Therefore, after stage t , the parameters $u(\beta, s)$ and $v(\beta, s)$ never increase and the number of type D components in U_β and V_β never increase (and none of these type D components grow).

For property 7, let s be the least stage such that β is never initialized after s . If there are no strategies α such that $\alpha * u_\infty \subseteq \beta$ or $\alpha * v_\infty \subseteq \beta$, then $b(\beta, t) = \infty$ for all $t \geq s$. Otherwise, by the induction hypothesis, for each $\alpha * u_\infty \subseteq \beta$, the value of $u(\alpha, t)$ approaches infinity and for each $\alpha * v_\infty \subseteq \beta$, the value of $v(\alpha, t)$ approaches infinity. Therefore, $b(\beta, t)$ approaches infinity. \square

Lemma 3.9 *For all strategies β and all β stages s , $a(\beta, s)$ is greater than $\max(\{u(\alpha, s) \mid \alpha * v_\infty \subseteq \beta\} \cup \{v(\alpha, s) \mid \alpha * \text{fin} \subseteq \beta\})$.*

Proof This lemma follows from three observations. First, whenever β is initialized, it defines $a(\beta, s)$ to be large at the next β stage. This large value is by definition greater than $\max(\{u(\alpha, s) \mid \alpha * v_\infty \subseteq \beta\} \cup \{v(\alpha, s) \mid \alpha * \text{fin} \subseteq \beta\})$. Second, if $\alpha * v_\infty \subseteq \beta$ and $u(\alpha, s)$ increases, then α takes outcome u_∞ and β is initialized. Third, if $\alpha * \text{fin} \subseteq \beta$ and $v(\alpha, s)$ increases, then α takes outcome v_∞ and β is initialized. \square

Lemma 3.10 *Assume that β sets up to diagonalize at stage s . No strategy $\alpha \neq \beta$ can add elements after this point to $T_{u(\beta, s)}$ (unless β is initialized or $u(\beta, s)$ increases at a later stage) or to $T_{v(\beta, s)}$ (unless β is initialized or $v(\beta, s)$ increases at a later stage).*

Proof β can set up to diagonalize in either step 1 or step 2 of the basic module. In either case, we have $a(\beta, s) < u(\beta, s) < v(\beta, s) < b(\beta, s)$ and β takes outcome $\beta * \text{fin}$ when β performs this action. To show that no $\alpha \neq \beta$ can add elements to $T_{u(\beta, s)}$ or $T_{v(\beta, s)}$ after this stage unless β is initialized or the parameters $u(\beta, s)$ or $v(\beta, s)$ change, we break into cases depending on the relative priority of α and β .

Case 1 $\alpha <_L \beta$. In this case, β is initialized at the end of the stage when α acts.

Case 2 $\beta * \text{fin} <_L \alpha$. In this case, α is initialized when β sets up to diagonalize at stage s . Therefore, $a(\alpha, t)$ for $t > s$ is greater than $v(\beta, s)$ and hence α works with trees with more type D components than $T_{v(\beta, s)}$ and cannot add new elements to either $T_{u(\beta, s)}$ or $T_{v(\beta, s)}$.

Case 3 $\beta * \text{fin} \subseteq \alpha$. By Lemma 3.9, the value of $a(\alpha, s)$ is greater than $v(\beta, s)$. As above, α works with trees with strictly more type D components than $T_{u(\beta, s)}$ and $T_{v(\beta, s)}$.

Case 4 $\alpha \subseteq \beta$. We break this case into four subcases.

Subcase 4a $\alpha * \text{triv} \subseteq \beta$. In this subcase, α does not add any elements to T without taking an outcome to the left of β and initializing β .

Subcase 4b $\alpha * \text{fin} \subseteq \beta$. By Lemma 3.9, $v(\alpha, s) < a(\beta, s)$. Hence, α works with trees with strictly fewer type D components than $T_{u(\beta, s)}$ and $T_{v(\beta, s)}$. If either the parameter $u(\alpha, s)$ or $v(\alpha, s)$ should increase at a later stage, α would take outcome $\alpha * u_\infty$ or $\alpha * v_\infty$ and β would be initialized.

Subcase 4c $\alpha * v_\infty \subseteq \beta$. By Lemma 3.9 and the definition of $b(\beta, s)$,

$$u(\alpha, s) < a(\beta, s) < u(\beta, s) < v(\beta, s) < b(\beta, s) < v(\alpha, s).$$

Therefore, U_α has strictly fewer type D components than $T_{u(\beta, s)}$ (unless $u(\alpha, s)$ later increases in which case β is initialized) and V_α has strictly more type D components than $T_{u(\beta, s)}$ and $T_{v(\beta, s)}$ (unless these parameters increase at a later stage).

Subcase 4d $\alpha * u_\infty \subseteq \beta$. By the definition of $b(\beta, s)$,

$$u(\beta, s) < v(\beta, s) < b(\beta, s) < u(\alpha, s).$$

Therefore, α works with components that contain strictly more type D trees than $T_{u(\beta, s)}$ and $T_{v(\beta, s)}$ (unless these parameters increase at a later stage). \square

Lemma 3.11 *Each component T_k is finite in the limit.*

Proof By the construction, a component T_k can only grow at stage $s > k$ if there is a strategy β such that $u(\beta, s) = k$ and β sets up to diagonalize or diagonalizes at stage s or such that $v(\beta, s) = k$ and β sets up to diagonalize at stage s . Because the parameters $a(\beta, s)$ are always chosen large, only finitely many strategies β can ever have $u(\beta, s) = k$ or $v(\beta, s) = k$ for any fixed value of k . Therefore, it suffices to show that such strategies only cause T_k to grow finitely often.

Consider the case when $v(\beta, s) = k$ and β sets up to diagonalize. In this situation, β could cause T_k to grow by adding an element to change a type A component in T_k into a type C component. By Lemma 3.9, unless β is initialized or $v(\beta, s)$ increases at a later stage, no other strategy can add an element to T_k after stage s . (Even if $u(\beta, s)$ later increases and β sets up to diagonalize again with the same value of $v(\beta, s)$, the tree T_k will not grow because we have already removed the type A component and no other strategy can have added a new type A component.)

If β is initialized after setting up to diagonalize, then it will work with components which contain strictly more type D components than T_k in the future. Therefore, in this case β causes only finitely much change to T_k .

If the value of $v(\beta, s)$ increases at a later stage, then α might reach a later stage t in which $u(\beta, t) = k$ and β sets up to diagonalize at stage t . In this situation, β will change T_k by adding a type A component. (We could also arrive at this situation without having $v(\beta, s) = k$ at some earlier stage in which case β will only add a type A component if T_k does not already have such a component.)

By Lemma 3.9, unless β is initialized or the value of $u(\beta, t)$ increases at a later stage, no strategy $\alpha \neq \beta$ can add elements to T_k . If β is later initialized or if $u(\beta, t)$ increases at a later stage, then β will work with components with strictly more type D trees than T_k and hence β will cause only finitely much growth to T_k .

If β later diagonalizes with $u(\beta, t) = k$, then β will add an element to T_k to change the type A component to either type B or type C. However, because β diagonalizes at most once with any given components, it will not add any more elements to T_k at a future stage. Therefore, β only adds finitely many elements to T_k . \square

Lemma 3.12 *All requirements $R_{\langle e, i \rangle}$ are satisfied.*

Proof Let β be the strategy on the true path such that $|\beta| = \langle e, i \rangle$. Assume for a contradiction that the φ_β -tree is isomorphic to T and that f_β is a nontrivial self-embedding of the φ_β -tree.

Let s be a stage after which β is never initialized and $a(\beta, s)$ has reached its final value. Because $\lim_s b(\beta, s) = \infty$, there must be a β stage $t > s$ and disjoint components U and V of the φ_β -tree such that the number of type D components in U is strictly greater than $a(\beta, t)$, the number of type D components in V is strictly less than $b(\beta, t)$, and f_β is an embedding of U into V . At this stage, β defines U_β and V_β permanently.

By the choice of U_β and V_β , we know that f_β is an embedding from U_β into V_β at stage t . These components may gain new type D components after stage t , but because f_β is a self-embedding of the φ_β -tree, it must eventually become defined on all of U_β as it grows and continue to be an embedding into V_β . Furthermore, it is possible that V_β gains new type D components more quickly than the parameter $b(\beta, s)$ grows. However, since $\lim_s b(\beta, s) = \infty$, we must eventually be in the situation where V_β has stopped growing and it has strictly fewer than $b(\beta, s)$ many type D components.

Therefore, we can assume without loss of generality that t is a β stage, that U_β and V_β have completely enumerated all of their type D components by stage t , that V_β has strictly fewer than $b(\beta, t)$ many type D components, and that f_β is an embedding of U_β into V_β . Notice that these assumptions imply that the parameters $u(\beta, t)$ and $v(\beta, t)$ have reached their limits and that $u(\beta, t) < v(\beta, t) < b(\beta, t)$.

At stage t , β sets up to diagonalize (in either step 1 or step 2 depending on the previous actions of β). β adds a type A component to $T_{u(\beta, t)}$ (if necessary) and changes the type A component in $T_{v(\beta, t)}$ to a type C component (if necessary). Because the φ_β -tree is isomorphic to T and because neither U_β nor V_β gains a type D component after stage t , there must be a stage $t' > t$ at which $U_\beta \cong T_{u(\beta, t')}$ and $V_\beta \cong T_{v(\beta, t')}$. At this stage, β moves to step 4 of the basic module and diagonalizes by changing the designated component in $T_{u(\beta, t')}$ so that it cannot be embedded into the designated component of V_β .

Because $U_\beta \cong T_{u(\beta, t')}$ before this additional element is added, we know that the number of each type of component in U_β and $T_{u(\beta, t')}$ (before the additional element is added) match up. Furthermore, because β is never initialized again and both $u(\beta, s)$

and $v(\beta, s)$ have reached their limits, by Lemma 3.9 we know that $T_{u(\beta, t')}$ does not change again after stage t' . Therefore, to make $U_\beta \cong T_{u(\beta, t')}$ (after the extra element is added), U_β must change its designated component to match the new type of the designated component in $T_{u(\beta, t')}$. Furthermore, because U_β is already committed by f_β to embedding the designated component of U_β into the designated component of V_β and because the new type of the designated component of $T_{u(\beta, t')}$ does not embed into the designated component of V_β , the embedding f_β cannot be extended in a way that is compatible with extending the designated component of U_β . This fact gives the desired contradiction. \square

This completes the verification that our construction succeeds.

4 Type 2 Trees

Recall that a type 2 computable tree is one which has no maximal ω -nodes and which has an isolated path. (The type 2 trees we construct below will all be finitely branching and hence will have no ω -nodes at all.) It is well known that there are such trees T which have a single path and such that this path codes $0'$. (The successor relation on T can even be computable. Among other places, such a construction is contained in the proof that König's Lemma for finitely branching trees is equivalent to ACA_0 in Simpson [14].) We construct such a tree and show that any nontrivial self-embedding of it can compute $0'$.

Lemma 4.1 *If T is a tree ordering on ω such that*

- (a) *T is finitely branching,*
- (b) *$\forall m, n (m \succeq n \implies m \geq n)$,*
- (c) *$\forall n (n \succeq \text{the immediate predecessor of } n + 1)$,*

then T has exactly one path X . Furthermore, if this path X is written as $\lambda = x_0 < x_1 < x_2 < \dots$ where x_{i+1} is the successor of x_i , then $x_n = \max\{m : \text{ht}(m) \leq n\}$.

Proof Let $x_n = \max\{m : \text{ht}(m) \leq n\}$ which exists by (a). For any fixed n , we will show by induction that $l \succeq x_n$ for all $l \geq x_n$. Thus for any n , x_n is an infinite node. By (b), x_n has height n and is the only infinite node of height n . Therefore, $x_0 < x_1 < x_2 < \dots$ gives the unique path through T .

Fixing $n \in \omega$, we show that $l \succeq x_n$ for all $l \geq x_n$ by induction on l . For the base of the induction, we have trivially that $x_n \succeq x_n$. For the induction step, suppose that $l \geq x_n$ and $l \succeq x_n$ and we show that $l + 1 \succeq x_n$. Let p be the immediate predecessor of $l + 1$ in T . By (c), $l \succeq p$ and hence p is comparable to x_n . So either $p \succeq x_n$, in which case $l + 1 \succeq x_n$ and we are done, or $p < x_n$, in which case $\text{ht}(l + 1) \leq n$ (as $\text{ht}(x_n) \leq n$) contradicting the fact that $x_n = \max\{m : \text{ht}(m) \leq n\}$ as $l + 1 > l \geq x_n$. \square

Theorem 4.2 *There is a computable type 2 tree T such that any nontrivial self-embedding of T computes $0'$.*

Proof We construct T to have exactly one infinite path X such that $\text{deg}_T(X) \geq 0'$. Fix a c.e. set K of degree $0'$.

$\langle T, \preceq \rangle$ is built computably in stages denoted $\langle T_s, \preceq_s \rangle$ with $T_s = \{0, 1, 2, \dots, s\}$ and $\preceq_s = \preceq \cap (T_s \times T_s)$ for all s . $T = \bigcup_s T_s = \omega$ and $\preceq = \bigcup_s \preceq_s$. At each stage s

we designate an element n_s of T_s to be the immediate predecessor of $s + 1$ and \leq_{s+1} is defined to be the reflexive and transitive closure of $\leq_s \cup \{(n_s, s + 1)\}$.

Let $\leq_0 = \{(0, 0)\}$ and at stage s let

$$n_s = \begin{cases} \max\{m \leq s \mid K_{s+1}[\text{ht}(m)] = K_s[\text{ht}(m)]\} & \text{if this set is nonempty,} \\ 0 & \text{otherwise.} \end{cases}$$

Lemma 4.3 *T has exactly one infinite path X and this path computes K .*

Proof Conditions (b) and (c) of Lemma 4.1 are satisfied immediately by the construction of T . Furthermore, because new nodes are always added as (current) leaves in T , the height of any node is fixed once it is placed in T . Therefore, we can speak of $\text{ht}(x)$ for any $x \in T$ without reference to a stage number.

To see that (a) is satisfied, we reason by contradiction. Assume that $x \in T$ is an ω -node. It follows that there are infinitely many stages $s_0 < s_1 < \dots$ such that $n_{s_i} = x$. At stage s_0 , x gains $s_0 + 1$ as a successor and hence x is no longer a leaf in T after stage s_0 . Consider any stage s_i for $i \geq 1$. Since $x = n_{s_i}$, we have $x \leq s_i$ and so $x < s_i$ (because s_i was added as a leaf at the previous stage and we know x is no longer a leaf in T). Therefore, x has a successor y_i such that $x < y_i \leq s_i$. By the definition of $n_{s_i} = x$, we know $K_{s_i+1}[\text{ht}(x)] = K_{s_i}[\text{ht}(x)]$ but $K_{s_i+1}[\text{ht}(y_i)] \neq K_{s_i}[\text{ht}(y_i)]$. However, $\text{ht}(y_i) = \text{ht}(x) + 1$, so $K_{s_i+1}[\text{ht}(x)] = K_{s_i}[\text{ht}(x)]$ and $K_{s_i+1}[\text{ht}(x) + 1] \neq K_{s_i}[\text{ht}(x) + 1]$ for all $i \geq 1$. Because K is a c.e. set, there can be at most one such stage s_i , giving the desired contradiction. (This argument really shows that T is binary branching.) Therefore (a) holds and T has exactly one path X .

We next show that $X \geq_T K$. Because $s + 1$ is added as a (current) leaf of T_s at stage s , the successor relation on T is computable. Therefore, from the set X , we can compute the sequence $x_0 < x_1 < \dots$ such that x_0 is the root of T , x_{i+1} is the successor of x_i , and $X = \{x_i \mid i \in \omega\}$.

By Lemma 4.1 we know that $x_n = \max\{m \mid \text{ht}(m) \leq n\}$. Furthermore, by the proof of Lemma 4.1, we know that for all $s \geq x_n$, we have $x_n \leq s$. We claim that for all n , $K_{x_n}[n] = K[n]$. Suppose not and fix n such that $K_{x_n}[n] \neq K[n]$. Let $s \geq x_n$ be such that $K_{s+1}[n] \neq K_s[n]$. Because $x_n \leq s$ and $\text{ht}(x_n) = n$, we have that $n_s < x_n$. Therefore, $x_n \not\leq s + 1$ contradicting the fact that $x_n \leq s + 1$. \square

Lemma 4.4 *Any nontrivial self-embedding φ of T computes K .*

Proof Let φ be a nontrivial self-embedding of T and let m be some node on X such that $\varphi(m) \neq m$. (That such a node exists is a consequence of X being the only path and there being no nontrivial self-embeddings of finite trees.) $\varphi(m)$ must also lie on X as T has only one infinite path. By induction one sees that for all n , $\varphi^n(m) \geq x_n$ and $\varphi^n(m) \in X$. Therefore, for all n , $\varphi^n(m) \geq x_n$, $K_{\varphi^n(m)}[n] = K_{x_n}[n] = K[n]$ and $\varphi \geq_T K$. \square

This completes the proof of Theorem 4.2. \square

The result can be improved slightly by replacing *nontrivial* with *weakly nontrivial* in 4.2. To show this we construct T' from T . T' will have, like T , a single isolated path X' that computes K . T' will be modified, however, to ensure that any weakly nontrivial self-embedding must move a node on X' .

We say a node n on T is *just off* X if n is not on X but the immediate predecessor is on X . Any weakly nontrivial self-embedding of T that fixes every node on X must be weakly nontrivial on some finite tree $T(n)$ with n just off X . Any finite tree can be properly extended to a finite tree that has no weakly nontrivial self-embedding by adding nodes extending the leaves so that no two leaves have the same height. This is what we do to ensure that $T(n)$ has no nontrivial self-embedding. Extensions may be added at different stages in the construction but we ensure that each leaf is extended only a finite amount.

We first repeat the construction of T using only the even numbers—adding node $2(s+1)$ as the immediate successor to $2n_s$ at stage s . We describe the placement of every odd number on T' .

We begin with $T'_0 = T_0$. As before at stage s we determine $2n_s$ and place $2(s+1)$ as its immediate successor. At stage s we also find all leaves extending $2n_s$ except $2(s+1)$ and we properly extend all such leaves with successive odd numbers so that any two distinct leaves have different heights. That is, we are guessing that the path X will pass through $2(s+1)$ and so all the other successors of $2n_s$ are just off X . Therefore, we want to extend the leaves above these other successors to have different lengths. T'_{s+1} is this extension of T_{s+1} . We need only show now that every leaf on T is extended only finitely and that all the odd numbers are used.

As in Lemma 4.1, x'_n (the n th element in the unique path X' in T') will be the numerically greatest even number of height less than or equal to n . x'_n is added to T' at stage $s = x'_n/2$ and for all $t \geq s$, $2n_t \geq x'_n$. So no more extensions will be added to the leaves above any node just off X whose height is less than or equal to n . Therefore, each leaf is extended finitely only a finite number of times.

To see that all the odd numbers are used we merely need to note that there are infinitely many nodes just off X' (otherwise X' would be computable) and that we have decreed that each leaf extending such a node must be properly extended.

We next turn our attention to nontrivial self-embeddings in computable trees $S \cong T$ where T is a computable type 2 tree.

Theorem 4.5 *There is a computable finitely branching tree S with exactly one infinite path (so S is a type 2 tree) such that no computable tree classically isomorphic to S has a computable nontrivial self-embedding.*

Proof Let $\langle T, \preceq_T \rangle$ be the tree constructed in Theorem 3.7, let T_i (for $i \in \omega$) be the sequence of components of T , let λ_T denote the root of T , and let λ_i denote the root of the component T_i . Let $A = \{a_i : i \in \omega\}$ be a set of distinct elements disjoint from T . We define the tree $\langle S, \preceq_S \rangle$ as follows:

1. $S = (T \setminus \{\lambda_T\}) \cup A$,
2. a_0 is the root of S ,
3. $\forall x, y \in T \setminus \{\lambda_T\} (x \preceq_T y \leftrightarrow x \preceq_S y)$, and
4. $\forall i \in \mathbb{N} (a_{i+1} \text{ and } \lambda_i \text{ are immediate successors of } a_i)$.

It is straightforward to see that there is a unique tree S satisfying 1, 2, 3, and 4 and that S has exactly the one infinite path given by $a_0 < a_1 < a_2 \cdots$.

We claim that any nontrivial self-embedding δ of S computes a nontrivial self-embedding of T , and hence by Theorem 3.7, δ must be noncomputable. To prove this claim, notice that because $a_0 < a_1 < \cdots$ is the unique path in S , we have that for each $i \in \omega$, $\delta(a_i) = a_j$ for some $j \geq i$. Because δ is nontrivial and every subtree off the unique path is finite, we can fix the least k such that $\delta(a_k) \neq a_k$. For every

$i \geq k$, $\delta(a_i) = a_j$ for some $j > i$. In other words, δ gives an embedding from T_i into T_j . It follows that δ induces a nontrivial self-embedding δ' of T given by

1. $\delta'(\lambda_T) = \lambda_T$ and
2. for all $x \in T \setminus \{\lambda_T\}$ ($\delta'(x) = \delta(x)$).

To finish the proof, let $\langle U, \leq_U \rangle$ be any computable tree classically isomorphic to S . We show that any nontrivial self-embedding δ of U computes a nontrivial self-embedding of a computable tree classically isomorphic to T . Hence by Theorem 3.7, $\delta >_T 0$.

As U is isomorphic to S , it too has exactly one infinite path B . We claim that B is computable. It is c.e. because $x \in B$ if and only if there exists a chain of height 4 above x (recall all the components of T have height 3). And it is co-c.e. because $x \notin B$ if and only if there exists a $b \in B$ such that b and x are incomparable.

We build a computable tree $\langle V, \leq_V \rangle$ isomorphic to T as follows. Let $V = (U \setminus B) \cup \{\rho\}$, where ρ is a new element which will serve as the root of V . For all $x, y \in V$ define $x \leq_V y$ if and only if $x = \rho$ or $x \leq_U y$. As B is computable, so is $\langle V, \leq_V \rangle$, and it is easy to see that V is isomorphic to T .

From δ we compute a nontrivial self-embedding δ' of V . Let $\delta'(\rho) = \rho$ and for all $x \in U \setminus B$, let $\delta'(x) = \delta(x)$. δ' is clearly computable from δ and hence $\delta >_T 0$. \square

5 Type 3 Trees

Recall that a type 3 computable tree is an infinite computable tree which has no maximal infinite node and no isolated paths. In particular, any infinite computable binary branching tree which has no isolated paths is a type 3 tree. In Theorem 2.3, we proved that each computable type 3 tree has a nontrivial self-embedding computable in $0''$. The next theorem shows that this bound is optimal.

Theorem 5.1 *There is an infinite computable binary branching tree S with no isolated paths such that any nontrivial self-embedding computes $0''$.*

Before proving Theorem 5.1, we outline the main steps of the proof. We begin by giving a particular computable approximation to $0''$ which is conducive to our coding methods. Next, we define a c.e. subtree $T \subseteq 2^{<\omega}$ such that $0''$ is coded into the branching levels of T . (A c.e. subtree $T \subseteq 2^{<\omega}$ is a c.e. set T of elements of $2^{<\omega}$ which is closed under initial segments. The tree order is given by the initial segment relation \subseteq .) We say n is a *branching level* of T if there is a string $\sigma \in T$ such that $|\sigma| = n$ and both $T(\sigma * 0)$ and $T(\sigma * 1)$ are infinite. We use a c.e. subtree T of $2^{<\omega}$ because it makes the notation easier when verifying properties such as where the branching levels occur in T and the fact that T has no isolated paths.

We show that from any nontrivial self-embedding of T we can compute a function dominating the branching levels and that any such function computes $0''$. Finally, we show how to define a computable tree $S \cong T$ for which the successor relation is computable. Because the branching levels of T are invariant under isomorphisms, we have $0''$ coded into the branching levels of S . From any nontrivial self-embedding of S , we can decode $0''$ as long as we can determine the height of each node in S . However, since the successor relation is computable in S , we can effectively determine the height of any node.

We begin by developing our computable approximation to $0''$. Fix a uniformly c.e. sequence of c.e. sets A_n for $n \in \omega$ such that $\{n \mid A_n \text{ is finite}\} \equiv_T 0''$. (For

example, we could use the standard enumeration of all c.e. sets.) Without loss of generality, we assume that in the uniform enumeration of the A_n sequence, exactly one set gets an element at each stage. Let $f(n) =$ the least s such that the sets among A_0, \dots, A_n which are finite have been completely enumerated by stage s .

Lemma 5.2 *For any function b which dominates f , $0'' \leq_T b \oplus 0'$.*

Proof Let $k \in \omega$ be such that $f(x) \leq b(x)$ for all $x \geq k$. To determine whether $n \in 0''$ for $n \geq k$, ask $0'$ whether A_n gets an element after stage $b(n)$. The answer to this question is no if and only if $n \in 0''$. \square

We want to define a computable approximation $f(n, s)$ to the function $f(n)$ so that $f(n) = \liminf_s f(n, s)$. To define $f(n, s)$, proceed as follows. If the sets A_0, \dots, A_n are all empty at stage s , then set $f(n, s) = 0$. If at least one of these sets is nonempty but none of them receives a new element at stage s , then let $f(n, s) = t$ where $t < s$ is the last stage at which one of these sets received an element.

If we are not in one of these two cases, then at stage s , exactly one set among A_0, \dots, A_n gets a new element. Let $i_{n,s} \leq n$ be such that $A_{i_{n,s}}$ gets a new element at stage s and let $t_{n,s} < s$ be the last stage at which $A_{i_{n,s}}$ received an element. (If s is the first stage at which $A_{i_{n,s}}$ gets an element, then set $t_{n,s} = 0$.) Let

$$I_{n,s} = \{j \leq n \mid A_j \text{ has received an element since } t_{n,s}\}.$$

$I_{n,s}$ represents our current guess at which sets among A_0, \dots, A_n are infinite. Let $f(n, s) = t$ where $t < s$ is the greatest stage such that there exists a $j \leq n$ for which $j \notin I_{n,s}$ and A_j gets an element at stage t . (If the sets A_j for $j \notin I_{n,s}$ are all empty or if $I_{n,s} = \{0, 1, \dots, n\}$, then set $f(n, s) = 0$.) That is, to calculate $f(n, s)$ we look at the sets A_j for $j \leq n$ which we currently think are not infinite and take the last stage at which one of these sets received a new element. The function $f(n, s)$ is a total computable function.

Lemma 5.3 *The function $f(n, s)$ satisfies the following properties.*

1. $f(n) = \liminf_s f(n, s)$.
2. For every $k > f(n)$, there is a stage s_k such that for all $t \geq s_k$ either $f(n, t) = f(n)$ or $f(n, t) > k$.

Proof Fix n and break into two cases. If A_0, \dots, A_n are all finite, then let u be the last stage at which any of these sets gets an element. Because $f(n, s) = u$ for all $s > u$, we have both property 1 and 2 in this case.

Otherwise, there is at least one set among A_0, \dots, A_n which is infinite. Let I be the set of all $i \leq n$ such that A_i is infinite and let $u_0 = f(n)$ be the last stage such that some A_j with $j \leq n$ and $j \notin I$ receives an element. Let $u_1 > u_0$ be a stage such that each A_i with $i \in I$ has received at least one element between stages u_0 and u_1 .

Consider any stage $s > u_1$ and split into two cases. First, if none of the sets A_i for $i \leq n$ receives an element at stage s , then $f(n, s) > u_0$ since $I \neq \emptyset$ and each A_i for $i \in I$ received an element after stage u_0 . Second, if one of the A_i sets for $i \in I$ does receive an element at stage s , then $t_{n,s} > u_0$ since each such set receives an element between stages u_0 and u_1 . Furthermore, $I_{n,s} \subseteq I$ since none of the sets A_j for $j \notin I$ receives an element after stage u_0 .

If $I_{n,s} = I$, then $f(n, s) = u_0 = f(n)$. If $I_{n,s} \subsetneq I$, then $f(n, s) > u_0$ since there is an A_i for which $i \in I \setminus I_{n,s}$ and this A_i received an element after stage u_0 . Therefore, for all $s > u_1$, $f(n, s) \geq u_0 = f(n)$.

Define a sequence of stages $v_0 < v_1 < v_2 < \dots$ such that $u_1 < v_0$ and at each stage v_k , $I_{n,v_k} = I$. To see that such a sequence exists, consider any stage $t > u_1$. We claim there is a stage $s > t$ such that $I_{n,s} = I$. To find s , let j_n be the index such that $j_n \leq n$ and A_{j_n} is the last set among A_i with $i \in I$ to receive a new element after stage t . Let $s > t$ be the first stage at which A_{j_n} receives a new element. Since A_{j_n} receives a new element at stage s , we have $i_{n,s} = j_n$. As $s > t$ is the first stage at which A_{j_n} receives a new element and since A_{j_n} has received an element since u_0 , we have $u_0 < t_{n,s} \leq t$. Each set A_i with $i \in I$ has now received a new element since stage t (and hence since $t_{n,s}$), so $I_{n,s} = I$ as required. Therefore, we have established the existence of the sequence $v_0 < v_1 < \dots$ with $I_{n,v_k} = I$. Since $I_{n,v_k} = I$, we have $f(n, v_k) = u_0 = f(n)$. Therefore, we have established property 1.

On the other hand, we can extend our sequence of stages $u_0 < u_1 < u_2 < \dots$ so that each A_i , $i \in I$, receives an element between stages u_k and u_{k+1} . Consider any $s > u_{k+1}$. If none of the sets A_i , $i \in I$, receive an element at s , then $f(n, s) > u_k$ since $I \neq \emptyset$ and each A_i , $i \in I$, has received an element since u_k . If some A_i , $i \in I$, does receive an element at stage s , then either $I_{n,s} = I$ (in which case $f(n, s) = u_0 = f(n)$) or $I_{n,s} \subsetneq I$ (in which case $f(n, s) > u_k$ since $A_{i_{n,s}}$ has received an element since stage u_k). Therefore, we have established property 2. \square

We define a computable function $g(n, s)$ from $f(n, s)$ that has one further property. We define $g(n, s)$ by induction on s , and for each s by induction on n . For every s , let $g(0, s) = f(0, s)$. Assume $g(i, t)$ has been defined for all $i \leq n$ and $t \leq s$, and we explain how $g(n+1, s)$ is defined. Let $k_{n,s}$ be the number of stages $t < s$ for which $g(i, t) = g(i, s)$ for all $i \leq n$ and let $m_{n,s}$ be the maximum value of $g(i, s)$ for $i \leq n$. Let $l_{n,s} = k_{n,s} + m_{n,s}$. Define $g(n+1, s) = f(n+1, l_{n,s})$.

Lemma 5.4 *The function $g(n, s)$ satisfies the following properties.*

1. $f(n) = \liminf_s g(n, s)$.
2. For every $k > f(n)$, there is a stage s_k such that for all $t \geq s_k$ either $g(n, t) = f(n)$ or $g(n, t) > k$.
3. For every n , there are infinitely many stages s at which $g(i, s) = f(i)$ for all $i \leq n$.

Proof We proceed by induction on n . By Lemma 5.3, these properties hold for $n = 0$. Assume these properties hold for $i \leq n$ and we prove them for $g(n+1, s)$. Applying property 3 to n , let $u_0 < u_1 < \dots$ list all the stages at which $g(i, s) = f(i)$ for all $i \leq n$. Let $M =$ the maximum of $f(i)$ for $i \leq n$. (Because f is nondecreasing, M is really just equal to $f(n)$.) At each stage u_k , we have $m_{n,u_k} = M$ and $k_{n,u_k} = k$, so by definition $g(n+1, u_k) = f(n+1, M+k)$. Therefore, as $k \rightarrow \infty$, $g(n+1, u_k)$ takes on all the values of $f(n+1, t)$ for $t > M$.

Let $t > M$ be a stage for which $f(n+1, t) = f(n+1)$. (By property 2 of Lemma 5.3 there are infinitely many such stages.) Let $k = t - M$. At stage u_k , we have $g(i, u_k) = f(i)$ for all $i \leq n$ by definition of u_k and we have $g(n+1, u_k) = f(n+1, M+k) = f(n+1, t) = f(n+1)$. Therefore, property 3 of this lemma holds for $n+1$.

For any $a \in \omega$, let $s_a > u_a$ be a stage such that for every $s > s_a$ and every $i \leq n$, either $g(i, s) = f(i)$ or $g(i, s) > a$. (The existence of s_a follows from property 2 of this lemma applied inductively to $i \leq n$.) Consider any $s > s_a$. By definition, $l_{n,s} = k_{n,s} + m_{n,s}$. We claim that $l_{n,s} \geq a$. There are two cases to consider. First, suppose $g(i, s) = f(i)$ for all $i \leq n$. In this case, $m_{n,s} = M$ and because $s_a > u_a$, there have been at least a many stages $t < s$ for which $g(i, t) = g(i, s) = f(i)$ for all $i \leq n$. Therefore, $k_{n,s} \geq a$, so $l_{n,s} \geq a$. Second, suppose that for some $i \leq n$ we have $g(i, s) \neq f(i)$. By the choice of s_a , $g(i, s) > a$, so $m_{n,s} > a$ and $l_{n,s} > a$. Therefore, in either case $l_{n,s} \geq a$ and so $g(n+1, s) = f(n+1, t)$ for some $t \geq a$.

The previous paragraph established that for all a , there is a stage s_a such that for all $s > s_a$, there is a $t \geq a$ for which $g(n+1, s) = f(n+1, t)$. Combining this fact with property 2 of Lemma 5.3 and with the fact that $g(n+1, u_k) = f(n+1)$ for infinitely many u_k yields properties 1 and 2 of this lemma. \square

We now put together the last two pieces of our approximating function. Let $h(n) =$ the least stage s for which $K_s[n+1] = K[n+1]$. Because $h(n)$ is a Δ_2^0 function, it has a computable approximation $h(n, s)$ such that $\lim_s h(n, s) = h(n)$. Finally, let $a(n, s)$ be the computable function defined by $a(n, s) = \max\{g(n, s), h(n, s)\}$.

Lemma 5.5 *The computable function $a(n, s)$ satisfies the following properties.*

1. $a(n) = \liminf_s a(n, s)$ exists and for all n , $a(n) \geq f(n), h(n)$.
2. For all n and for every $k > a(n)$, there is a stage s_k such that for all $t \geq s_k$, either $a(n, t) = a(n)$ or $a(n, t) > k$.
3. For every n , there are infinitely many stages s at which $a(i, s) = a(i)$ for all $i \leq n$.
4. For any function b which dominates a , $0'' \leq_T b$.

Proof Properties 1 through 3 follow from Lemma 5.4 and the fact that $h(n) = \lim_s h(n, s)$. Property 4 follows from the fact that if b dominates a , then b dominates both h and f . The fact that b dominates h gives $0' \leq_T b$. Combining this fact with Lemma 5.2 gives $0'' \leq_T b$. \square

We next define a c.e. subtree $T \subseteq 2^{<\omega}$ such that the branching levels of T dominate the function $a(n)$. The branching levels are the levels that contain branching nodes. That is, level k in T is a branching level if there is a node $\sigma \in T$ such that $|\sigma| = k$ and both $T(\sigma * 0)$ and $T(\sigma * 1)$ are infinite.

The basic idea of our construction is as follows. We make the node $0^{a(0)} \in 2^{<\omega}$ the branching node of least length in T . Therefore, we need to insure that both $0^{a(0)} * 0$ and $0^{a(0)} * 1$ have infinitely many extensions in T . Since we want the next branching level to be above $a(1)$, we make the next branching nodes equal to $0^{a(0)} * 0 * 0^{a(1)}$ and $0^{a(0)} * 1 * 0^{a(1)}$. To do this, we need to ensure that for each node σ of length 2, we have that the nodes $0^{a(0)} * \sigma(0) * 0^{a(1)} * \sigma(1)$ have infinitely many extensions in T . (Below, we will denote these nodes by τ_2^σ .) We repeat this process by making the next branching nodes have the form $0^{a(0)} * \sigma(0) * 0^{a(1)} * \sigma(1) * 0^{a(2)}$ for all $|\sigma| = 2$. In other words, for all strings σ of length 3, we need to ensure that the nodes $0^{a(0)} * \sigma(0) * 0^{a(1)} * \sigma(1) * 0^{a(2)} * \sigma(2)$ have infinitely many extensions in T .

By repeating the process in the previous paragraph, the branching levels of T will occur at levels of the form $n + \sum_{i=0}^n a(i)$ for $n \in \omega$. We use the approximation

$a(n, s)$ to define a c.e. tree with these branching levels and we use properties 2 and 3 of Lemma 5.5 to verify that these levels are branching levels and that no other levels are branching. Since $a(n) \leq n + \sum_{i=0}^n a(i)$, we have the required domination property. We then need to show how to extract information about the branching levels from any nontrivial self-embedding of our tree.

We begin with some notation. For any s , any $n \leq s$, and any string $\sigma \in 2^{<\omega}$ with $|\sigma| = n + 1$, we define

$$\tau_{n,s}^\sigma = 0^{a(0,s)} * \sigma(0) * 0^{a(1,s)} * \sigma(1) * \dots * 0^{a(n,s)} * \sigma(n).$$

For any n and any string $\sigma \in 2^{<\omega}$ such that $|\sigma| = n + 1$, we let

$$\tau_n^\sigma = 0^{a(0)} * \sigma(0) * 0^{a(1)} * \sigma(1) * \dots * 0^{a(n)} * \sigma(n).$$

For any nonempty string $\alpha \in 2^{<\omega}$, let α' denote the string obtained by removing the last element of α . Notice that

$$(\tau_{n,s}^\sigma)' = 0^{a(0,s)} * \sigma(0) * 0^{a(1,s)} * \sigma(1) * \dots * 0^{a(n,s)}$$

is a string of length $n + \sum_{i=0}^n a(i, s)$ and that $(\tau_n^\sigma)'$ is a string of length $n + \sum_{i=0}^n a(i)$.

As described above, the goal of our construction is to make each node of the form $(\tau_n^\sigma)'$ a branching node of our c.e. tree T and we accomplish this goal by making each node τ_n^σ have infinitely many extensions on T . Because we cannot effectively know which nodes are of the form τ_n^σ , we have to use the approximations $\tau_{n,s}^\sigma$. At stage s , we add at least one new node extending each string of the form $\tau_{s,s}^\sigma$. We then verify that in the limit, this process makes the branching nodes of T exactly those nodes of the form $(\tau_n^\sigma)'$.

Once we have such a tree T , we show that from any nontrivial embedding δ of T , we can effectively obtain a nontrivial embedding ι of T and a node α such that there are at least n many branching levels below $|\iota^{n+1}(\alpha)|$. By our calculation of the branching levels (described in the previous paragraph), the function $c(n) = |\iota^{n+1}(\alpha)|$ dominates the function $a(n)$, and hence by Lemma 5.5, c computes $0''$. Because c is obtained effectively from δ , we conclude that $0'' \leq_T \delta$.

The subtree $T \subseteq 2^{<\omega}$ is enumerated in stages as a sequence of finite trees $T_0 \subseteq T_1 \subseteq T_2 \subseteq \dots$. Set $T_0 = \emptyset$. To define T_{s+1} , consider each string $\sigma \in 2^{<\omega}$ which has length $s + 1$. Let α_σ be the lexicographically least element of $2^{<\omega}$ which extends $\tau_{s,s}^\sigma$ and which is not in T_s . Add α_σ and all of its initial segments to T_s . T_{s+1} is the tree formed by adding these strings when σ ranges over all elements of $2^{<\omega}$ of length $s + 1$. Our desired tree is $T = \bigcup_s T_s$.

Lemma 5.6 *For each n and each $\sigma \in 2^{<\omega}$ of length $n + 1$, the node $(\tau_n^\sigma)'$ is a branching node of T .*

Proof Let $u_0 < u_1 < \dots$ be the stages such that $n < u_0$ and $a(i, u_k) = a(i)$ for all $i \leq n$. For each such stage, $(\tau_n^\sigma)' = (\tau_{n,u_k}^\sigma)'$ and $(\tau_{n,u_k}^\sigma)' \subseteq \tau_{u_k, u_k}^\zeta$ for all strings ζ such that $\sigma' \subseteq \zeta$ and $|\zeta| = u_k + 1$. Therefore, both $T_{u_k}((\tau_n^\sigma)' * 0)$ and $T_{u_k}((\tau_n^\sigma)' * 1)$ gain extra elements at stage $u_k + 1$. Therefore, these trees are infinite and $(\tau_n^\sigma)'$ is a branching node in T . \square

Lemma 5.7 *If ζ is a branching node of T , then there is an n such that $|\zeta| = n + \sum_{i=0}^n a(i)$.*

Proof Suppose $\zeta \in T$ is such that there is an n such that

$$n + \sum_{i=0}^n a(i) < |\zeta| < n + 1 + \sum_{i=0}^{n+1} a(i). \quad (1)$$

By Lemma 5.5, let u be a stage such that for all $s > u$ and all $i \leq n + 1$, either $a(i, s) = a(i)$ or $a(i, s) > |\zeta|$. Fix any stage $s > u$.

We claim that there is a $j \leq n$ such that

$$j + \sum_{i=0}^j a(i, s) < |\zeta| < j + 1 + \sum_{i=0}^{j+1} a(i, s). \quad (2)$$

The proof of the claim breaks into two cases. If $a(i, s) = a(i)$ for all $i \leq n$, then the claim with $j = n$ follows from Equation (1). Otherwise, let $j < n$ be the least number such that $a(j + 1, s) \neq a(j + 1)$. In this case,

$$j + \sum_{i=0}^j a(i, s) = j + \sum_{i=0}^j a(i) < n + \sum_{i=0}^n a(i) < |\zeta|.$$

Because $a(j + 1, s) > |\zeta|$, we have $j + 1 + \sum_{i=0}^{j+1} a(i, s) > |\zeta|$ and hence Equation (2) holds in this case as well.

By Equation (2), at stage s there is a unique $\sigma \in 2^{<\omega}$ with length $j + 1$ such that $\tau_{j,s}^\sigma \subseteq \zeta$. Furthermore, for $a \in \{0, 1\}$ we have $\zeta \not\subseteq (\tau_{j+1,s}^{\sigma * a})'$. That is,

$$0^{a(0,s)} * \sigma(0) * \dots * 0^{a(j,s)} * \sigma(j) \subseteq \zeta \not\subseteq 0^{a(0,s)} * \sigma(0) * \dots * 0^{a(j,s)} * \sigma(j) * 0^{a(j+1,s)}.$$

It follows that at stage $s + 1$, $T_s(\zeta * 0)$ gets a new element but $T_s(\zeta * 1)$ does not. Because this property holds for any $s > u$, $T(\zeta * 1) = T_u(\zeta * 1)$ is finite; so ζ is not a branching node of T .

To finish the proof, we need to show that if $\zeta \in T$ and $|\zeta| < a(0)$, then ζ is not a branching node. The proof of this fact is similar to (but simpler than) the argument above and we leave it to the reader to verify. \square

From Lemmas 5.6 and 5.7, we obtain the following fact.

Lemma 5.8 *The n th branching level of T is given by the formula $b(n) = n + \sum_{i=0}^n a(i)$.*

Lemma 5.9 *If $T(\zeta)$ is infinite, then $\zeta \subseteq \tau_n^\sigma$ for some n and σ with $|\sigma| = n + 1$.*

Proof Suppose that $\zeta \not\subseteq \tau_n^\sigma$ for any n and σ . We show that $T(\zeta)$ is finite. First, notice that ζ must contain at least one value of 1 or else $\zeta \subseteq \tau_n^\sigma$ for sufficiently large n by choosing σ to contain all zeros.

Second, notice that if ζ does not have $0^{a(0)}$ as an initial segment, then this property follows trivially. That is, fix a stage u such that for all $s > u$, $a(0, s) \geq a(0)$. At any stage $s > u$, we add nodes only above strings $\tau_{s,s}^\sigma$ and each string $\tau_{s,s}^\sigma$ begins with $0^{a(0,s)}$. Because this string is not an initial segment of ζ , $T_s(\zeta)$ does not get a new element at stage $s + 1$. Therefore, $T_u(\zeta) = T(\zeta)$ and hence $T(\zeta)$ is finite.

It remains to consider the case when $0^{a(0)}$ is an initial segment of ζ and ζ contains at least one value of 1. Let j be the largest value such that there are strings α (with $|\alpha| = j + 1$) and μ such that

$$\zeta = 0^{a(0)} * \alpha(0) * \dots * 0^{a(j)} * \alpha(j) * \mu.$$

Fix such j , α and μ . Because $\zeta \not\subseteq \tau_n^\sigma$ for any n and σ , the string μ must contain at least one value of 1. Write $\mu = \mu_0 * 1 * \mu_1$ where μ_0 is such that $\mu_0(k) = 0$ for all $k < |\mu_0|$. Because j is chosen maximal, $|\mu_0| < a(j+1)$.

Let u be a stage such that for all $s > u$ and for all $i \leq j+1$, $a(i, s) = a(i)$ or $a(i, s) > |\zeta|$. The lemma follows from the claim that $T_u(\zeta) = T(\zeta)$. To prove this claim, fix any $s > u$ and we show that $T_s(\zeta)$ does not gain a new element at stage $s+1$. We split into two cases. First, suppose that for all $i \leq j$, $a(i, s) = a(i)$. The only way for $T_s(\zeta)$ to gain a new element at stage $s+1$ is if there is a string σ of length $s+1$ such that $\zeta \subseteq \tau_{s,s}^\sigma$. Because $a(i, s) = a(i)$ for $i \leq j$, this string σ must satisfy $\sigma(i) = \alpha(i)$ for all $i \leq j$. It follows that $0^{a(0)} * \alpha(0) * \dots * 0^{a(j)} * \alpha(j) * 0^{a(j+1,s)}$ is an initial segment of $\tau_{s,s}^\sigma$. However, regardless of whether $a(j+1, s) = a(j)$ or not, we have $|\mu_0| < a(j+1, s)$. Hence the strings $0^{a(0)} * \alpha(0) * \dots * 0^{a(j)} * \alpha(j) * 0^{a(j+1,s)}$ and ζ are incomparable. (The point is that ζ contains the value 1 right after μ_0 while the other string has value 0 in this position.) Therefore, $T_s(\zeta)$ does not get a new element in this case.

The other case is when there is an $i < j$ for which $a(i, s) \neq a(i)$. Let k denote the least such i . The argument is similar. $T_s(\zeta)$ can gain a new element only if there is a σ such that $\zeta \subseteq \tau_{s,s}^\sigma$. Because $a(i, s) = a(i)$ for all $i < k$, we have $\sigma(i) = \alpha(i)$ for $i < k$ and hence $0^{a(0)} * \alpha(0) * \dots * 0^{a(k-1)} * \alpha(k-1) * 0^{a(k,s)}$ is an initial segment of $\tau_{s,s}^\sigma$. Because $a(k, s) > |\zeta|$, this string is incomparable with ζ and hence $T_s(\zeta)$ does not get a new element in this case. \square

Lemma 5.10 *The tree T has no isolated paths.*

Proof This lemma follows immediately from Lemmas 5.9 and 5.6. \square

Lemma 5.11 *If $\delta : T \rightarrow T$ is a nontrivial self-embedding, then there is a string ζ such that $|\delta(\zeta)| > |\zeta|$.*

Proof Suppose there is no such string ζ . Because $|\zeta| \leq |\delta(\zeta)|$ for any self-embedding δ and because T is binary branching, it follows that for each n , δ restricted to the strings of length n in T is a permutation. Therefore, δ is onto and hence is not nontrivial. \square

Lemma 5.12 *If $\delta : T \rightarrow T$ is a nontrivial self-embedding then there is a $k \in \omega$ and a node ζ such that $\zeta \subsetneq \delta^k(\zeta)$.*

Proof By Lemma 5.11, let μ_0 be a node such that $|\mu_0| < |\delta(\mu_0)|$. If $\mu_0 \subseteq \delta(\mu_0)$ then $\mu_0 \subsetneq \delta(\mu_0)$ and we can let $\zeta = \mu_0$ and $k = 1$ to verify the lemma. Otherwise, assume that $\mu_0 \not\subseteq \delta(\mu_0)$. Let μ_1 be such that $|\mu_1| = |\mu_0|$ and $\mu_1 \subseteq \delta(\mu_0)$. Notice that $\mu_1 \neq \mu_0$ and $\delta(\mu_0) \neq \mu_0$.

We proceed by induction. Assume that $n \geq 1$ and we have defined a sequence of pairwise distinct nodes $\mu_0, \mu_1, \dots, \mu_n$ such that $|\mu_i| = |\mu_0|$ for all $i \leq n$ and $\mu_{i+1} \subseteq \delta(\mu_i)$ and $\mu_i \neq \delta(\mu_i)$ for all $i < n$. (The last two sentences of the previous paragraph establish the required properties when $n = 1$.)

We claim that in this situation, $\mu_n \neq \delta(\mu_n)$. Suppose that $\mu_n = \delta(\mu_n)$. Because $|\mu_n| = |\mu_{n-1}|$ and $\mu_n \neq \mu_{n-1}$, μ_n and μ_{n-1} are incomparable nodes. However, $\delta(\mu_n) = \mu_n \subseteq \delta(\mu_{n-1})$. Therefore, $\delta(\mu_n)$ and $\delta(\mu_{n-1})$ are comparable contradicting the fact that δ is a self-embedding.

Next, we let μ_{n+1} be such that $|\mu_{n+1}| = |\mu_0|$ and $\mu_{n+1} \subseteq \delta(\mu_n)$. We claim that if $\mu_{n+1} = \mu_i$ for some $i \leq n$, then the conclusion of the lemma is true. Otherwise, if $\mu_{n+1} \neq \mu_i$ for all $i \leq n$, then we add μ_{n+1} to the list of pairwise distinct nodes above and continue by induction. Because there are only finitely many nodes at level $|\mu_0|$, we must eventually find an n such that $\mu_{n+1} = \mu_i$ for some $i \leq n$. Hence, the lemma follows from the claim in this paragraph.

Suppose that $\mu_{n+1} = \mu_i$ for some $i \leq n$ and let $l \geq 1$ be such that $i = (n+1) - l$. In this situation we have $\mu_i = \mu_{n+1} \subseteq \delta^l(\mu_i)$. We claim that $\mu_i \neq \delta^l(\mu_i)$ (and hence we have established the lemma with $\xi = \mu_i$ and $k = l$). We break into three cases.

Case 1 $i = 0$. In this case, we have $\mu_0 \subseteq \delta^l(\mu_0)$. But, $|\mu_0| < |\delta(\mu_0)|$ implies $|\mu_0| < |\delta^l(\mu_0)|$ so we have $\mu_0 \subsetneq \delta^l(\mu_0)$ as required.

Case 2 $l = 1$. In this case, we have $\mu_i \subseteq \delta(\mu_i)$. Because $i \leq n$, we know $\mu_i \neq \delta(\mu_i)$, so $\mu_i \subsetneq \delta(\mu_i)$ as required.

Case 3 $i > 0$ and $l > 1$. For a contradiction, assume that $\mu_i = \delta^l(\mu_i)$. We have $\mu_i = \mu_{n+1} \subseteq \delta(\mu_n)$ and $\mu_i \subseteq \delta(\mu_{i-1})$. By our induction hypothesis, μ_n and μ_{i-1} are incomparable nodes. Furthermore, we have

$$\mu_i = \mu_{n+1} \subseteq \delta(\mu_n) \subseteq \delta^2(\mu_{n-1}) \subseteq \cdots \subseteq \delta^l(\mu_i) = \mu_i.$$

Therefore, $\delta(\mu_n) = \mu_i$ so $\delta(\mu_n)$ and $\delta(\mu_{i-1})$ are comparable nodes, violating the fact that δ is a self-embedding. \square

For any nontrivial self-embedding $\delta : T \rightarrow T$, we can fix k and ξ as in Lemma 5.12 and let $\gamma = \delta^k : T \rightarrow T$. γ is a nontrivial self-embedding of T such that $\xi \subsetneq \gamma(\xi) \subsetneq \gamma^2(\xi) \subsetneq \cdots$. It will also be useful to consider the nontrivial self-embedding $\iota : T \rightarrow T$ given by $\iota = \gamma^2$. Notice that both ι and γ are obtained from δ by finitely many parameters.

Lemma 5.13 *Let δ be any nontrivial self-embedding $\delta : T \rightarrow T$ and let γ and ι be defined from δ as above. There are nodes α and β_0 such that $\alpha \subsetneq \beta_0 \subsetneq \iota(\alpha)$ and β_0 is a branching node.*

Proof Fix ξ as in the paragraph before this lemma. Because $\xi, \gamma(\xi), \gamma^2(\xi), \dots$ traces out a path in T and because T has no isolated paths, there must be a $j \geq 1$ and a branching node β_0 such that $\gamma^j(\xi) \subseteq \beta_0 \subsetneq \gamma^{j+1}(\xi)$. Let $\alpha = \gamma^{j-1}(\xi)$. Because $\alpha = \gamma^{j-1}(\xi) \subsetneq \gamma^j(\xi) \subseteq \beta_0$, we have $\alpha \subsetneq \beta_0$. Because $\iota = \gamma^2$, we have $\beta_0 \subsetneq \gamma^{j+1}(\xi) = \gamma^2(\gamma^{j-1}(\xi)) = \iota(\alpha)$. \square

Lemma 5.14 *Let $\iota : T \rightarrow T$ be a nontrivial self-embedding for which there are nodes α and β_0 such that $\alpha \subsetneq \beta_0 \subsetneq \iota(\alpha)$ and β_0 is a branching node. Then there is a branching node β_1 such that $\iota(\alpha) \subsetneq \beta_1 \subsetneq \iota^2(\alpha)$.*

Proof Fix α and β_0 . Because $\alpha \subsetneq \beta_0 \subsetneq \iota(\alpha)$, we have

$$\begin{aligned} \alpha \subsetneq \beta_0 \subsetneq \iota(\alpha) \subsetneq \iota(\beta_0) \subsetneq \iota(\beta_0 * 0) \\ \alpha \subsetneq \beta_0 \subsetneq \iota(\alpha) \subsetneq \iota(\beta_0) \subsetneq \iota(\beta_0 * 1). \end{aligned}$$

Let β_1 be the infimum of $\iota(\beta_0 * 0)$ and $\iota(\beta_0 * 1)$. Because these two nodes are incomparable, β_1 is strictly contained in both of them. From the offset containments

above, it is clear that $\beta_0 \subsetneq \iota(\alpha) \subsetneq \beta_1$. Let $i_0 \in \{0, 1\}$ be such that $\beta_0 * i_0 \subseteq \iota(\alpha)$. Because $\iota(\beta_0 * i_0) \subseteq \iota^2(\alpha)$ and $\beta_1 \subsetneq \iota(\beta_0 * i_0)$ we have $\beta_1 \subsetneq \iota^2(\alpha)$.

Finally, because β_0 is a branching node, both $T(\iota(\beta_0 * 0))$ and $T(\iota(\beta_0 * 1))$ are infinite. Therefore, the infimum of $\iota(\beta_0 * 0)$ and $\iota(\beta_0 * 1)$ (which is β_1) is a branching node. \square

Lemma 5.15 *Let $\delta : T \rightarrow T$ be any nontrivial self-embedding. There is a nontrivial self-embedding $\iota : T \rightarrow T$ (defined from δ together with finitely many parameters) and a node α such that the sequence $c(n) = |\iota^{n+1}(\alpha)|$ dominates the branching level function $b(n)$ of T (see Lemma 5.8).*

Proof Define ι from δ as above and let α and β_0 be as in Lemma 5.13. Applying Lemma 5.14 inductively, we obtain a sequence of branching node $\beta_0 \subsetneq \beta_1 \subsetneq \beta_2 \subsetneq \dots$ such that $\iota^n(\alpha) \subsetneq \beta_n \subsetneq \iota^{n+1}(\alpha)$. (For $n = 0$, we define $\iota^0(\alpha) = \alpha$.) Therefore, there are at least n many branching levels below $|\iota^{n+1}(\alpha)|$. \square

To prove Theorem 5.1, we need to transform the c.e. subtree $T \subseteq 2^{<\omega}$ into a computable tree S . This transformation is easily done in a general setting.

Lemma 5.16 *For any c.e. subtree $\hat{T} \subseteq 2^{<\omega}$, there is a computable tree \hat{S} such that $\hat{S} \cong \hat{T}$. Furthermore, we can assume that the successor relation is computable in \hat{S} .*

Proof If \hat{T} is finite, this lemma follows trivially. Assume \hat{T} is infinite and \hat{T} is the range of the total computable 1-1 function φ_e . Let \hat{S} have domain ω and let $\leq_{\hat{S}}$ be defined by $n \leq_{\hat{S}} m \Leftrightarrow \varphi_e(n) \subseteq \varphi_e(m)$. Then φ_e is an isomorphism from $(\hat{S}, \leq_{\hat{S}})$ to (\hat{T}, \subseteq) as required. Furthermore, m is a successor of n in \hat{S} if and only if $\varphi_e(m) \setminus \varphi_e(n) \neq \emptyset$. \square

We now present the proof of Theorem 5.1. Let T be the c.e. subtree of $2^{<\omega}$ we have constructed and let $S \cong T$ be the computable tree with a computable successor relation given by Lemma 5.16. Let b denote the branching level function for S (which is the same as the branching level function for T since $S \cong T$). By Lemma 5.8, b dominates a and hence by Lemma 5.5, $0'' \leq_T b$. Fix any nontrivial self-embedding $\delta : S \rightarrow S$. By Lemma 5.15, there is a nontrivial self-embedding $\iota : S \rightarrow S$ (defined from finitely many parameters) and a node α such that the function $c(n) = \text{ht}(\iota^{n+1}(\alpha))$ dominates b (and hence by Lemma 5.5, $0'' \leq_T c$). Because we only need finitely many parameters to obtain ι from δ , we have $\iota \leq_T \delta$. Furthermore, because the successor function is computable in S , we can determine the height of any node in S . Therefore, $0'' \leq_T c \leq_T \iota \leq_T \delta$ as required. This completes the proof of Theorem 5.1.

Because the coding in the proof of Theorem 5.1 is done with the branching levels of S and these levels are invariant under isomorphisms, we also obtain a result concerning the existence of nontrivial self-embeddings of computable type 3 trees up to isomorphism.

Theorem 5.17 *There is a computable type 3 tree S such that for any computable $\hat{S} \cong S$ and any nontrivial self-embedding $\delta : \hat{S} \rightarrow \hat{S}$, $0'' \leq_T 0' \oplus \delta$. In particular, \hat{S} does not have any Δ_2^0 nontrivial self-embeddings and \hat{S} does not have any nontrivial self-embeddings which are strictly between $0'$ and $0''$.*

Proof Fix a computable tree $S \cong T$ where T is the c.e. subtree $T \subseteq 2^{<\omega}$ constructed above. Fix any $\hat{S} \cong S$ and any nontrivial self-embedding δ of \hat{S} . Let ι and c be the functions given by Lemma 5.15 for \hat{S} and δ . The only change from the proof of Theorem 5.1 is that we do not know that the successor relation in \hat{S} is computable. However, since $0' \oplus \delta$ can compute both ι and the successor relation in \hat{S} , we have $0'' \leq_T c \leq_T 0' \oplus \delta$. \square

6 Chains and Antichains

The fact that any infinite partial order must have either an infinite chain or an infinite antichain is a simple application of Ramsey's Theorem for pairs and two colors. Herrmann [7] examined the effective content of this result and proved the following theorem.

Theorem 6.1 (Herrmann [7]) *If P is an infinite computable partial order, then P has either an infinite Δ_2^0 chain or an infinite Π_2^0 antichain. In addition, there is an infinite computable partial order which has no infinite Σ_2^0 chains or antichains.*

In this section, we consider this result in the context of trees rather than general partial orders and we show that for trees these results can be improved by exactly one quantifier.

Theorem 6.2 *Let T be an infinite computable tree. T has either an infinite computable chain or an infinite Π_1^0 antichain.*

Proof If T has infinitely many leaves, then the set of leaves is an infinite Π_1^0 antichain. Otherwise, T must have a node x such that $T(x)$ is infinite and contains no leaves. In this case, let $x_0 = x$ and x_{i+1} be the $\leq_{\mathbb{N}}$ least element of T which satisfies $x_i < x_{i+1}$. The sequence x_0, x_1, \dots gives an infinite computable chain. \square

Theorem 6.3 *There is an infinite binary branching computable tree such that T has no infinite c.e. chains or antichains.*

Proof We build (T, \preceq) to meet the following requirements.

$$\begin{aligned} R_{2e} &: W_e \text{ is not an infinite chain} \\ R_{2e+1} &: W_e \text{ is not an infinite antichain} \end{aligned}$$

We build T in stages beginning with $T_0 = \{\lambda\}$. Throughout the construction, we maintain the property that each node x is either currently a leaf or else has exactly two successors. Each requirement R_i keeps a parameter r_i such that any node x added to T by a lower priority requirement after r_i is defined satisfies $r_i \preceq x$. For uniformity of notation, we set $r_{-1} = \lambda$. If a strategy is initialized, then all of its parameters become undefined. Any parameter not explicitly redefined or undefined by initialization retains its value. If a requirement ends the current stage, then it initializes all lower priority requirements. The action for R_{2e} at stage s is as follows.

1. If s is the first stage at which R_{2e} is eligible to act or if R_{2e} has been initialized since it was last eligible to act, let a be such that $r_{2e-1} \preceq a$ and a is a leaf in T_s . Add new nodes b and c to T_s as immediate successors of a . Set $r_{2e} = b$ and end the stage.

2. If r_{2e} is defined but R_{2e} has not succeeded yet, then check whether there is a node $x \in T_s$ such that $r_{2e} \preceq x$ and $x \in W_{e,s}$. If not, then let R_{2e+1} act. If so, then let z denote the immediate predecessor of x . Since z is not a leaf, it has two immediate successors. Let y denote the successor of z which is not equal to x . Redefine $r_{2e} = y$ and end the stage. We say that R_{2e} has succeeded.
3. If R_{2e} has succeeded, then let R_{2e+1} act.

The action for R_{2e+1} at stage s is as follows.

1. If s is the first stage at which R_{2e+1} is eligible to act or if R_{2e+1} has been initialized since it was last eligible to act, define $r_{2e+1} = r_{2e}$. End the stage.
2. If r_{2e+1} is defined and R_{2e+1} has not succeeded yet, check whether there is a node $x \in T_s$ such that $r_{2e+1} \preceq x$ and $x \in W_e$. If not, then let R_{2e+2} act. If so, then redefine $r_{2e+1} = x$ and end the stage. We say R_{2e+1} succeeds.
3. If R_{2e+1} has succeeded, then let R_{2e+2} act.

This argument is finite injury so each parameter reaches a limit. Because nodes are added to T only in step 1 of the R_{2e} action, T has the property that at each stage, each node is either currently a leaf or has exactly two successors.

To see that R_{2e} is met, let s be the least stage such that R_{2e} is never initialized after stage s . The parameter r_{2e} is defined at stage s and can only change values after stage s if R_{2e} changes the value in step 2 of its action.

There are two cases to consider. First, suppose there is a stage $t > s$ and a node $x \in T_t$ such that $r_{2e} \preceq x$ and $x \in W_{e,t}$. In this case, r_{2e} is redefined so that r_{2e} is incompatible with x . Because r_{2e} is not changed again and because no strategy of higher priority than R_{2e} adds elements to T after stage t , there are only finitely many elements in T which are not above this final value of r_{2e} . Therefore, x cannot be part of an infinite chain and R_{2e} is met.

Second, suppose there is no such stage t and node x . In this case, r_{2e} has reached its limit at stage s and every node added to T after stage s is added above r_{2e} . Because there are only finitely many nodes in T which are not above r_{2e} , there cannot be an infinite chain which is disjoint from $T(r_{2e})$. Therefore, R_{2e} is met.

The argument that R_{2e+1} is met is quite similar. Let s be the least stage such that R_{2e+1} is never initialized after s and let r_{2e+1} denote the value of the parameter at stage s . If there is no node $x \in W_e$ such that $r_{2e+1} \preceq x$, then R_{2e+1} never changes the value of r_{2e+1} and there are only finitely many nodes of T which are not above r_{2e+1} . Any infinite antichain must intersect $T(r_{2e+1})$, so R_{2e+1} is met.

If there is a node $x \in W_e$ such that $r_{2e+1} \preceq x$, then let x be the first such node seen by R_{2e+1} after stage s . At this point, r_{2e+1} is redefined to be equal to x , so there are only finitely many nodes in T which are not comparable to x . Hence, no set containing x can be an infinite antichain. Therefore, R_{2e+1} is won. \square

Theorem 6.4 *Let L be any low set. There is an infinite binary branching computable tree T such that T has no infinite chains or antichains computable from L .*

Proof We need to meet the following requirements.

$$R_{2e} : \phi_e^L \text{ is not an infinite chain}$$

$$R_{2e+1} : \phi_e^L \text{ is not an infinite antichain}$$

As in the proof of Theorem 6.3, we build T in stages and maintain the property that each node x is either currently a leaf or else has exactly two immediate successors.

Each requirement R_i keeps a parameter r_i as before. The main change in this construction is that the value of r_i can change more than once (but still only finitely often) after the last time R_i is initialized. As before, whenever a requirement ends a stage, it initializes all lower priority requirements.

R_{2e} keeps three parameters: r_{2e} , \hat{r}_{2e} , and x_{2e} . The r_{2e} parameter is used as before to force lower priority requirements to work above r_{2e} . The \hat{r}_{2e} parameter is used to store an “old value” of r_{2e} in case our approximations to computations from L change and we need to revert back to an earlier situation and wait for reconvergence. The x_{2e} parameter will be explained when it appears in the construction below.

When R_{2e} first acts or if R_{2e} has been initialized since its last action, it lets a be a node such that $r_{2e-1} \leq a$ and a is currently a leaf. It adds two new nodes b and c as immediate successors of a in T , defines $\hat{r}_{2e} = r_{2e} = b$, and ends the stage. At future stages s , R_{2e} requires lower priority strategies to work above r_{2e} and it tries to decide whether $\exists x \exists t (r_{2e} \leq x \wedge \varphi_{e,t}^L(x) = 1)$. This predicate is Σ_1^L , so it is computable from L' and hence from O' (since L is low). Fix the Δ_2^0 predicate $P(e, k)$ defined by

$$P(e, k) \Leftrightarrow \exists x \exists t (k \leq x \wedge \varphi_{e,t}^L(x) = 1).$$

Let $P(e, k, s)$ be a computable approximation such that $P(e, k) = \lim_s P(e, k, s)$.

At stage s , R_{2e} checks whether $P(e, \hat{r}_{2e}, s) = 1$. (Notice that $\hat{r}_{2e} = r_{2e}$ at this point, so R_{2e} is really checking whether $P(e, r_{2e}, s) = 1$.) If not, then R_{2e} has no need to diagonalize and it lets R_{2e+1} act. If so, then R_{2e} wants to find the least witness x which appears to satisfy this existential statement. We define a second Δ_2^0 predicate $Q(x, u)$ by

$$Q(x, u) \Leftrightarrow \forall t (t \geq u \rightarrow \varphi_{e,t}^L(x) = 1)$$

and fix a computable approximation $Q(x, u, s)$ such that $Q(x, u) = \lim_s Q(x, u, s)$. (The predicate $Q(x, u)$ is computable from L' , so it is Δ_2^0 because L is low.) R_{2e} looks for the least x such that $\hat{r}_{2e} = r_{2e} \leq x$ and $Q(x, s, s)$. If there is no such x then R_{2e} lets R_{2e+1} act and waits to check again at the next stage. Eventually, it must find an x and s for which $Q(x, s, s)$. (Of course, if $P(e, \hat{r}_{2e}, s)$ changes from value 1 to value 0 while R_{2e} is waiting for such an x , it ends the stage and returns to waiting for $P(e, \hat{r}_{2e}, s)$ to have value 1.)

When R_{2e} finds such an x , it defines its third parameter $x_{2e} = x$. Let z be the immediate predecessor of x_{2e} and let y be the successor z which is not equal to x_{2e} . R_{2e} sets $r_{2e} = y$ but leaves the value of \hat{r}_{2e} unchanged. That is, \hat{r}_{2e} retains the “old value” of r_{2e} . R_{2e} ends the stage (and hence initializes the lower priority strategies so that they will work above the new value of r_{2e} in the future).

If $P(e, \hat{r}_{2e})$ really holds and x_{2e} really is a correct witness for this existential statement, then we have successfully diagonalized. However, it is possible that either $P(e, \hat{r}_{2e})$ does not hold or that x_{2e} is not a correct witness. Therefore, at each future stage s , R_{2e} continues to check whether $P(e, \hat{r}_{2e}, s) = 1$. If this value ever changes to 0, then R_{2e} redefines r_{2e} to have value $r_{2e} = \hat{r}_{2e}$, cancels its parameter x_{2e} , ends the stage, and returns to waiting for $P(e, \hat{r}_{2e}, s) = 1$. If $P(e, \hat{r}_{2e}, s)$ retains its value of 1, then R_{2e} checks whether $Q(x_{2e}, s, s)$ still gives the value 1. If so, then R_{2e} continues to believe it has correctly diagonalized and lets R_{2e+1} act. If $Q(x_{2e}, s, s) = 0$ at some future stage s (while $P(e, \hat{r}_{2e}, s) = 1$), then R_{2e} cancels the parameter x_{2e} , redefines r_{2e} to have value $r_{2e} = \hat{r}_{2e}$, ends the stage, and returns to looking for the least x such that $Q(x, s, s) = 1$.

To understand why this strategy eventually succeeds, let u be a stage such that R_{2e} is never initialized after u . At stage u , R_{2e} defines r_{2e} and \hat{r}_{2e} and it will never change the value of \hat{r}_{2e} again. (The entire construction is finite injury so there is such a stage.) Because $P(e, \hat{r}_{2e})$ is a Δ_2^0 predicate, there is a $t \geq u$ such that for all $s \geq t$, $P(e, \hat{r}_{2e}, s)$ is either constantly 0 or constantly 1.

If $P(e, \hat{r}_{2e}, s)$ is eventually constantly 0, then r_{2e} will eventually be set permanently equal to \hat{r}_{2e} . From this stage on, all nodes added to T are above \hat{r}_{2e} . Because $P(e, \hat{r}_{2e})$ does not hold, φ_e^L does not place any elements from $T(\hat{r}_{2e})$ into its chain. Because there are only finitely many elements of T outside of $T(\hat{r}_{2e})$, φ_e^L cannot compute an infinite chain and R_{2e} is met.

If $P(e, \hat{r}_{2e}, s)$ is eventually constantly 1, then there is an x such that $\hat{r}_{2e} \leq x$ and x is a witness to the existential statement $P(e, \hat{r}_{2e})$. Because $Q(x, u, s)$ is a Δ_2^0 predicate and we look for the least witness x , R_{2e} eventually defines x_{2e} such that $Q(x_{2e}, s, s)$ has reached its limit of 1. Both r_{2e} and x_{2e} have reached their limits at this stage. After this stage, all elements added to T are above r_{2e} and hence are incomparable with x_{2e} . Because $\varphi_e^L(x_{2e}) = 1$, φ_e^L cannot compute an infinite chain in T so R_{2e} is met.

In either case, notice that \hat{r}_{2e} and r_{2e} reach limits and that x_{2e} either reaches a limit or there is a stage after which it is never defined. Therefore, R_{2e} only initializes lower priority strategies finitely often.

The strategy to meet requirement R_{2e+1} is similar. R_{2e+1} also keeps three parameters r_{2e+1} , \hat{r}_{2e+1} , and x_{2e+1} . When it first acts (or after it has been initialized), R_{2e+1} sets $\hat{r}_{2e+1} = r_{2e+1} = r_{2e}$ and ends the stage.

At future stages, R_{2e+1} checks whether $P(e, \hat{r}_{2e+1}, s) = 1$. If not, it lets R_{2e+2} act. If so, it looks for the least x such that $Q(x, s, s) = 1$. If there is no such x , it lets R_{2e+2} act next. If there is such an x , it sets $x_{2e+1} = x$, redefines r_{2e+1} so that $r_{2e+1} = x_{2e+1}$, and ends the stage. (As above, it leaves \hat{r}_{2e+1} unchanged to mark the “old value” of r_{2e+1} . If $P(e, \hat{r}_{2e+1}, s)$ changes values from 1 to 0 while R_{2e+1} is waiting for such an x , it ends the stage and returns to waiting for $P(e, \hat{r}_{2e+1}, s)$ to equal 1.)

Once x_{2e+1} is defined, R_{2e+1} continues to check whether $P(e, \hat{r}_{2e+1}, s) = 1$. If this value ever changes to 0, it cancels x_{2e+1} , redefines r_{2e+1} so that $r_{2e+1} = \hat{r}_{2e+1}$, ends the stage, and returns to waiting for $P(e, \hat{r}_{2e+1}, s)$ to equal 1. As long as $P(e, \hat{r}_{2e+1}, s)$ remains equal to 1, R_{2e+1} checks whether $Q(x_{2e+1}, s, s)$ continues to equal 1. As long as it does, R_{2e+1} lets R_{2e+2} act. If $Q(x_{2e+1}, s, s)$ changes values to 0, then R_{2e+1} cancels x_{2e+1} , redefines r_{2e+1} to have value $r_{2e+1} = \hat{r}_{2e+1}$, ends the stage, and returns to looking for the least x such that $Q(x, s, s) = 1$.

The analysis that R_{2e+1} eventually succeeds and that it initializes the lower priority requirements only finitely often is similar to the analysis given for R_{2e} . The details are left to the reader. \square

There is no need to restrict ourselves to a single low set L in the proof of Theorem 6.4. That is, essentially the same proof (with a little extra bookkeeping in the indices) shows that if L_i (for $i \in \mathbb{N}$) is a sequence of uniformly low, uniformly Δ_2^0 sets, then there is an infinite binary branching computable tree T such that T has no infinite chains and no infinite antichains computable from any of the L_i sets. By Jockusch and Soare [8] and Simpson [14], there is an ω -model \mathcal{M} of WKL_0 such that the second-order part of \mathcal{M} consists of all the sets in the Turing

ideal generated by a sequence $L_0 \leq_T L_1 \leq_T \dots$ of uniformly low, uniformly Δ_2^0 sets. Thus, we obtain the following corollary.

Corollary 6.5 *WKL₀ is not strong enough to prove that every infinite binary branching tree has either an infinite chain or an infinite antichain.*

Notes

1. A discrete linear order is a linear order in which every element except the least has an immediate predecessor and every element except the greatest has an immediate successor.
2. A Turing degree is called PA over $0'$ if it can compute an infinite path through any infinite $0'$ -computable subtree of $2^{<\omega}$.

References

- [1] Cenzer, D., “ π_1^0 classes in recursion theory,” pp. 37–88 in *Handbook of Computability Theory*, edited by E. R. Griffor, vol. 140 of *Studies in Logic and the Foundations of Mathematics*, North-Holland Publishing Co., Amsterdam, 1999. Zbl 0923.03001. MR 1721236. 2
- [2] Cenzer, D., and J. B. Remmel, “ Π_1^0 classes in mathematics,” pp. 623–821 in *Handbook of Recursive Mathematics, Vol. 2*, edited by Yu. L. Ershov, S. S. Goncharov, A. Nerode, and J. B. Remmel, vol. 139 of *Studies in Logic and the Foundations of Mathematics*, North-Holland, Amsterdam, 1998. Zbl 0941.03044. MR 1673586. 2
- [3] Corominas, E., “On better quasi-ordering countable trees,” *Discrete Mathematics*, vol. 53 (1985), pp. 35–53. Special volume on ordered sets and their applications (L’Arbresle, 1982). Zbl 0575.05024. MR 786475. 3
- [4] Downey, R. G., C. Jockusch, and J. S. Miller, “On self-embeddings of computable linear orderings,” *Annals of Pure and Applied Logic*, vol. 138 (2006), pp. 52–76. Zbl 1105.03036. MR 2183808. 2
- [5] Downey, R. G., and S. Lempp, “The proof-theoretic strength of the Dushnik-Miller theorem for countable linear orders,” pp. 55–57 in *Recursion Theory and Complexity (Kazan, 1997)*, edited by M. Arslanov and S. Lempp, vol. 2 of *de Gruyter Series in Logic and its Applications*, Walter de Gruyter & Co, Berlin, 1999. Zbl 0951.03053. MR 1724931. 2
- [6] Dushnik, B., and E. W. Miller, “Concerning similarity transformations of linearly ordered sets,” *Bulletin of the American Mathematical Society*, vol. 46 (1940), pp. 322–26. Zbl 0024.30102. MR 0001919. 2
- [7] Herrmann, E., “Infinite chains and antichains in computable partial orderings,” *The Journal of Symbolic Logic*, vol. 66 (2001), pp. 923–34. Zbl 1018.03038. MR 1833487. 4, 32
- [8] Jockusch, C. G., Jr., and R. I. Soare, “ Π_1^0 classes and degrees of theories,” *Transactions of the American Mathematical Society*, vol. 173 (1972), pp. 33–56. Zbl 0262.02041. MR 0316227. 35

- [9] Kruskal, J. B., “Well-quasi-ordering, the Tree Theorem, and Vazsonyi’s conjecture,” *Transactions of the American Mathematical Society*, vol. 95 (1960), pp. 210–25. Zbl 0158.27002. MR 0111704. 2
- [10] Lempp, S., C. McCoy, R. Miller, and R. Solomon, “Computable categoricity of trees of finite height,” *The Journal of Symbolic Logic*, vol. 70 (2005), pp. 151–215. Zbl 1104.03026. MR 2119128. 2
- [11] Miller, R., “The computable dimension of trees of infinite height,” *The Journal of Symbolic Logic*, vol. 70 (2005), pp. 111–41. Zbl 1098.03049. MR 2119126. 2, 3
- [12] Richter, L. J., “Degrees of structures,” *The Journal of Symbolic Logic*, vol. 46 (1981), pp. 723–31. Zbl 0512.03024. MR 641486. 2
- [13] Ross, D., “Tree self-embeddings,” *Canadian Mathematical Bulletin*, vol. 32 (1989), pp. 30–33. Zbl 0692.04006. MR 996120. 3
- [14] Simpson, S. G., *Subsystems of Second Order Arithmetic*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1999. Zbl 0909.03048. MR 1723993. 20, 35
- [15] Soare, R. I., *Recursively Enumerable Sets and Degrees. A Study of Computable Functions and Computably Generated Sets*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1987. Zbl 0667.03030. MR 882921. 4

Acknowledgments

The authors thank Carl Jockusch and Ted Slaman for helpful discussions regarding the results in the last section of this article. Solomon’s research was partially supported by an NSF Grant DMS-0400754.

Department of Mathematics and Statistics
King Fahd University of Petroleum and Minerals
Dhahran 31261
SAUDI ARABIA
binns@kfupm.edu.sa

Department of Mathematics
University of Hawai’i at Mānoa
Honolulu HI 96822
bjoern@math.hawaii.edu

Department of Mathematics
196 Auditorium Road
University of Connecticut U-3009
Storrs CT 06269-3009
lerman@math.uconn.edu
schmerl@math.uconn.edu
solomon@math.uconn.edu