

King Fahd University of Petroleum and Minerals
Department of Mathematical Sciences
Math 202 Exam I
Semester II, 2006- (052)
Dr. Faisal Fairag

Name:	KEY
-------	-----

ID:	KEY
-----	-----

Sec:	2 3
------	-----

Serial No:	KEY
------------	-----

Question #		Points
1		30
2		30
3		30
4		30
5		20
6		10
7		15
8		15
9		15
10		15
11		10
Total:		220

1. Solve: $xy' = 2xe^x - y + 6x^2$ (#13/page 73)
(Show all your work)

~~divid by x~~

$$y' = \cancel{2e^x} - \frac{y}{x} + 6x$$

$$x \frac{dy}{dx} = 2xe^x - y + 6x^2$$

$$x dy = (2xe^x - y + 6x^2) dx$$

$$(2xe^x - y + 6x^2) dx - x dy = 0 \quad \text{--- (1)}$$

Equation (1) is **exact** with $\triangle 5$

$$M(x,y) = 2xe^x - y + 6x^2, \quad N(x,y) = -x$$

$$\frac{\partial M}{\partial y} = -1 \quad \text{and} \quad \frac{\partial N}{\partial x} = -1 \Rightarrow (1) \text{ is exact}$$

$$\triangle 5 \quad f(x,y) = \int N(x,y) dy + g(x) = \int -x dy + g(x)$$

$$f(x,y) = -xy + g(x) \Rightarrow \frac{\partial f}{\partial x} = -y + g'(x)$$

$$\text{Now, } \frac{\partial f}{\partial x} = M(x,y) \Rightarrow -y + g'(x) = 2xe^x - y + 6x^2$$

$$\Rightarrow \triangle 5 \quad g'(x) = 2xe^x + 6x^2 \Rightarrow g(x) = (2x-2)e^x + 2x^3 \quad \triangle 5$$

$$\text{Hence, } f(x,y) = -xy + 2(x-1)e^x + 2x^3 \quad \triangle 5$$

Now, $\boxed{-xy + 2(x-1)e^x + 2x^3 = C}$ is the general solution $\triangle 5$

$$\text{Another: } f = \int M dx + g(y) = 2(x-1)e^x - yx + 2x^3 + g(y)$$

$$\frac{\partial f}{\partial y} = -x + g'(y) = N \Rightarrow g'(y) = 0 \Rightarrow \boxed{g(y) = 0}$$

$$\boxed{f(x,y) = -xy + 2(x-1)e^x + 2x^3} \Rightarrow \boxed{-xy + 2(x-1)e^x + 2x^3 = C}$$

2. Solve: $(3x + y)dy = (x + 3y)dx$ (#7/page 78) — (1)
 (Show all your work. Hint: it is homog of degree 1)

let $y = ux$ — (2)

$dy = u dx + x du$ — (3)

use (2) & (3) in (1)

$$(3x + ux)(u dx + x du) = (x + 3ux) dx$$

$$x(3+u)(u dx + x du) = x(1+3u) dx$$

divid by x

$$(3+u)(u dx + x du) = (1+3u) dx$$

$$\cancel{3u dx} + \underline{3x du} + u^2 dx + \underline{xu du} = dx + \cancel{3u dx}$$

$$(3x + xu) dx = (1 - u^2) dx$$

divid by $x(1-u^2)$

$$\frac{x(3+u) du}{x(1-u^2)} = \frac{(1-u^2)}{x(1-u^2)} dx$$

$$\left(\frac{3+u}{1-u^2} \right) du = \frac{1}{x} dx$$

$$\left[\frac{2}{1-u} + \frac{1}{1+u} \right] du = \frac{dx}{x}$$

integrate

$$2 \ln |1-u| + \ln |1+u| = \ln |x| + C$$

$$2 \ln \left| 1 - \frac{y}{x} \right| + \ln \left| 1 + \frac{y}{x} \right| = \ln |x| + C$$

3. Solve: $y^{\frac{1}{2}} y' + y^{\frac{3}{2}} - x^2 = 0$ (1)
(Show all your work)

multiply (1) by $y^{-\frac{1}{2}}$

$y' + y - x^2 y^{-\frac{1}{2}} = 0 \Rightarrow y' + y = x^2 y^{-\frac{1}{2}}$ (2)

(2) is Bernoulli with $n = -\frac{1}{2}$, $p(x) = 1$, $f(x) = x^2$

let $u = y^{1-n} \Rightarrow u = y^{1+\frac{1}{2}} \Rightarrow u = y^{\frac{3}{2}}$ (**)

$y = u^{\frac{2}{3}} \Rightarrow y' = \frac{2}{3} u^{-\frac{1}{3}} \cdot u'$ (*)

use (*) and (**) in (2):

$\frac{2}{3} u^{-\frac{1}{3}} \cdot u' + u^{\frac{2}{3}} = x^2 \cdot u^{-\frac{1}{3}}$ (3)

multiply (3) by $\frac{3}{2} u^{\frac{1}{3}}$

$u' + \frac{3}{2} u = x^2$ (4) is linear in u .

integrating factor = $u_1 = e^{\frac{3}{2}x} = e^{\frac{3}{2}x}$

multiply (4) by $e^{\frac{3}{2}x}$

$e^{\frac{3}{2}x} u' + \frac{3}{2} e^{\frac{3}{2}x} u = x^2 e^{\frac{3}{2}x}$

$\Rightarrow \frac{d}{dx} [e^{\frac{3}{2}x} \cdot u] = x^2 e^{\frac{3}{2}x}$

integrate

$e^{\frac{3}{2}x} \cdot u = \int x^2 e^{\frac{3}{2}x} \cdot dx$

$e^{\frac{3}{2}x} \cdot u = (\frac{2}{3}x^2 - \frac{8}{9}x + \frac{16}{27}) e^{\frac{3}{2}x} + c$

$\Rightarrow u = \frac{2}{3}x^2 - \frac{8}{9}x + \frac{16}{27} + ce^{-\frac{3}{2}x}$ use $u = y^{\frac{3}{2}}$ give

$\Rightarrow y^{\frac{3}{2}} = \frac{2}{3}x^2 - \frac{8}{9}x + \frac{16}{27} + ce^{-\frac{3}{2}x}$

Aside:

x^2	\times	$e^{\frac{3}{2}x}$
$2x$	\rightarrow	$\frac{2}{3}e^{\frac{3}{2}x}$
2	\rightarrow	$\frac{4}{9}e^{\frac{3}{2}x}$
0	\rightarrow	$\frac{8}{27}e^{\frac{3}{2}x}$

 $\int x^2 e^{\frac{3}{2}x} dx = (\frac{2}{3}x^2 - \frac{8}{9}x + \frac{16}{27}) e^{\frac{3}{2}x} + c$

4. Solve: $(2x + y + 1)y' = 1$ (#14/page 85)
(Show all your work)

$y' = \frac{1}{2x + y + 1}$ is in the form $y' = f(Ax + By + c)$

Let $u = 2x + y + 1$ $\triangle 5$

$\Rightarrow y = u - 2x - 1$ (2)

$\Rightarrow y' = u' - 2$ (3)

use (2) & (3) in (1):

$u' - 2 = \frac{1}{u}$

$\Rightarrow u' = \frac{1}{u} + 2 \Rightarrow u' = \frac{1 + 2u}{u}$

$\Rightarrow du = \left(\frac{1 + 2u}{u}\right) dx$

$\Rightarrow \left(\frac{u}{1 + 2u}\right) du = dx$ $\triangle 10$ separable

integrate
 $\int \frac{u}{1 + 2u} du = \int dx$

$\frac{1}{2}u - \frac{1}{4} \ln|1 + 2u| + C_1 = x + C_2$

$\Rightarrow 2u - \ln|1 + 2u| + 4C_1 = 4x + 4C_2$

$2u - \ln|1 + 2u| - 4x = C_3$ $\triangle 10$

Hence, $\boxed{2(2x + y + 1) - \ln|4x + 2y + 3| - 4x = C}$ $\triangle 5$

is one-parameter family of solutions

Aside:
 $\frac{1}{1 + 2u} \int \frac{u}{u + \frac{1}{2}}$
 $\frac{u}{1 + 2u} = \frac{1}{2} - \frac{1}{2} \frac{1}{1 + 2u}$
 $\int \frac{u}{1 + 2u} du = \frac{1}{2}u - \frac{1}{4} \ln|1 + 2u|$

5. Use an appropriate substitution to reduce the DE

$$y' = -x^4 + \frac{2}{x}y + y^2 \quad (1)$$

into a linear DE. Write the new DE in the following form

$$u' + p(x)u = f(x) \quad (2)$$

where $y_1 = x^2$ is a known solution of the DE. [Note: Just reduce it to linear DONOT SOLVE]

(1) is Riccati equation.

Let $y = y_1 + \frac{1}{u}$

$$y = x^2 + \frac{1}{u} \quad \text{--- (3) } \triangle 5$$

$$\Rightarrow y' = 2x - \frac{1}{u^2} u' \quad \text{--- (4) } \triangle 5$$

Use (3) & (4) in (1):

$$2x - \frac{1}{u^2} u' = -x^4 + \frac{2}{x} \left(x^2 + \frac{1}{u}\right) + \left(x^2 + \frac{1}{u}\right)^2$$

$$\underline{2x} - u^{-2} u' = \underline{-x^4} + \underline{2x} + \frac{2}{x} u^{-1} + \underline{x^4} + \underline{2x^2} u^{-1} + u^{-2}$$

$$-u^{-2} u' = \left(\frac{2}{x} + 2x^2\right) u^{-1} + u^{-2}$$

multiply by $-u^2$

$$u' = -\left(\frac{2}{x} + 2x^2\right) u - 1 \quad \triangle 10$$

$$\Rightarrow \boxed{u' + \left(\frac{2}{x} + 2x^2\right) u = -1} \quad \text{--- (5)}$$

(5) is linear DE with

$$p(x) = \frac{2}{x} + 2x^2 \quad \text{and} \quad f(x) = -1$$

6. $y = 2 \frac{1+c e^{4x}}{1-c e^{4x}}$ is a one-parameter family of solutions of the first-order DE $y' = y^2 - 4$. Which one of the following statements is TRUE.

- (a) $y = 2$ is a singular solution
 (b) $y = 2$ is a trivial solution
 (c) $y = 0$ is a trivial solution
 → (d) $y = -2$ is a singular solution
 (e) $y = 0$ is a particular solution

$y = -2$ is a solution.
 Check: LHS = $y' = \frac{d}{dx}[-2] = 0$
 RHS = $y^2 - 4 = (-2)^2 - 4 = 4 - 4 = 0$
 but no value for c so that $-2 = 2 \frac{1+c e^{4x}}{1-c e^{4x}}$
 Hence, $y = -2$ is not a member of the family.
 ⇒ $y = -2$ is singular solution.

7. The DE

$$y^2 x^3 dx + y^2 x^3 dy = xy^3 dy \quad (3)$$

is classified as

- (a) separable
 (b) linear in y
 (c) linear in x
 (d) exact
 (e) made exact
 (f) homog. of degree α
 (g) Bernoulli in y
 (h) Bernoulli in x
 (i) $y' = f(Ax + By + C)$
 (j) Riccati in y
 (k) Riccati in x
8. Find an appropriate integrating factor which make the non-exact DE

$$6xydx + (4y + 9x^2) = 0 \quad (4)$$

an exact DE.

- (a) y^6
 (b) x^2
 (c) y^{-2}
 (d) $12x$
 → (e) y^2

$$M = 6xy, \quad N = 4y + 9x^2$$

$$M_y = 6x, \quad N_x = 18x$$

observe that: $\frac{N_x - M_y}{M} = \frac{18x - 6x}{6xy} = \frac{12x}{6xy} = \frac{2}{y}$
 = function of y alone.

$$\text{integrating factor} = \mu(y) = e^{\int \frac{N_x - M_y}{M} dy}$$

$$= e^{\int \frac{2}{y} dy} = e^{\ln y^2} = y^2$$

9. If $y(x)$ is the solution of the IVP

$$x^2 y' = y(1-x), \quad y(-1) = -1 \quad (5)$$

Then $y(2) =$

[Note: equation (5) is separable]

- (a) $\frac{1}{2}e^{-3/2}$
 (b) $-\frac{1}{2}e^{-1/2}$
 (c) $\frac{1}{2}e^{3/2}$
 (d) 0
 (e) $\frac{1}{2}e^{-1/2}$

10. Determine a region of the xy plane for which the differential equation

$$y' = \frac{y^2 + 4}{x^2 - 4} \quad (6)$$

would have a unique solution.

- (a) $(-4, 4)$
 (b) $(0, +\infty)$
 (c) $(-\infty, 0)$
 → (d) $(4, +\infty)$
 (e) $(-\infty, 4)$

$$f(x, y) = \frac{y^2 + 4}{x^2 - 4} = \left(\frac{1}{x^2 - 4}\right)y^2 + \frac{4}{x^2 - 4}$$

$$\frac{\partial f}{\partial y} = \frac{2y}{x^2 - 4}$$

$$\frac{-2}{\phi} \quad \frac{2}{\oplus}$$

$f(x, y)$ and $\frac{\partial f}{\partial y}$ are discontinuous at $x = \pm 2$.
 The interval $(4, +\infty)$ does not contain $x = \pm 2$.

→ (5) can be written as $\frac{dy}{y} = \frac{1-x}{x^2} dx$

separable

integrate $\Rightarrow c_1 + \ln|y| = -\frac{1}{x} - \ln|x| + c_2$

$$\Rightarrow \ln|y| + \ln|x| = c_3 - \frac{1}{x} \Rightarrow \ln|xy| = c_3 - \frac{1}{x}$$

$$\Rightarrow |xy| = e^{c_3 - \frac{1}{x}} \Rightarrow |xy| = e^{c_3} \cdot e^{-\frac{1}{x}} \Rightarrow xy = \pm e^{c_3} \cdot e^{-\frac{1}{x}}$$

$$\Rightarrow xy = c e^{-\frac{1}{x}} \Rightarrow y = \frac{c e^{-\frac{1}{x}}}{x} \quad (*)$$

Now, $y(-1) = -1 \Rightarrow -1 = \frac{c e^{-\frac{1}{-1}}}{-1} \Rightarrow -1 = \frac{c e^{-1}}{-1} \Rightarrow c = \frac{1}{e} = e^{-1}$

(*) and $c = e^{-1} \Rightarrow y = \frac{e^{-1} \cdot e^{-\frac{1}{x}}}{x} \Rightarrow y = \frac{e^{-(1+\frac{1}{x})}}{x}$ is the sol of the IVP

Now, $y(2) = \frac{e^{-(1+\frac{1}{2})}}{2} = \frac{e^{-\frac{3}{2}}}{2} = \frac{1}{2} e^{-3/2}$

Aside:
 $\frac{1-x}{x^2} = \frac{1}{x^2} - \frac{1}{x}$
 $\int \frac{1-x}{x^2} dx = -\frac{1}{x} - \ln|x| + c_2$

