

King Fahd University of Petroleum and Minerals  
Department of Mathematical Sciences  
Math 301 Exam II  
Semester I, 2006- (061)  
Dr. Faisal Fairag

Serial NO:	KEY	
ID:		
Name:	KEY	FORM <b>A</b>

Q		Points
1		8
2		8
3		8
4		8
5		8
6		8
7		8
8		16
9		16
10		16
11		16
12		16
Total		130

Say a prayer & Good luck 😊

FORM-A

FORM-B

dacabab

cbababa

1) If  $f(t) = e^{2t}(t-1)^2$  and  $F(s) = L[f(t)]$ , then  $F(1) =$

$$F(s) = \frac{1}{(s-2)^3} + \frac{2}{(s-3)^2} + \frac{1}{(s-4)^2}$$

(a)  $\frac{18}{7}$

(b) 0

(c)  $\frac{139}{108}$   
B

(d)  $\frac{-7}{18}$   
A

(e) none

2) If  $f(t) = L^{-1}\left\{\frac{1}{s^2+2s+5}\right\}$ , then  $f(0) =$

(a)  $\frac{1}{2}$   
A

(b) 0  
B

(c)  $\frac{1}{5}$

(d)  $\frac{1}{\pi^2+5}$

(e) none

3) If  $f(t) = L^{-1}\left\{\frac{e^{-2s}}{s^2(s-1)}\right\}$ , then  $f(3) =$

(a)  $e^2-3$   
B

(b) 0

(c)  $e-2$   
A

(d) 1

(e) none

4) If  $F(s) = L[te^{-3t} \cos 3t]$ , then  $F(-3) =$

(a)  $\frac{-1}{9}$   
A

(b) 0  
B

(c)  $\frac{1}{9}$

(d)  $e^{-9}$

(e) none

5) If  $y(t)$  is the solution of the initial value problem :  $y' + y = \delta(t-1)$ ,  $y(0) = 2$   
Then  $y(1) =$

(a) 2  
B

(b)  $\frac{2+e}{2}$   
A

(c) 0

(d) 1

(e) none

6) If  $h(t) = t * \cos t$ , then  $h(\pi) =$

(a) 2  
A

(b) 1  
B

(c) 0

(d) -1

(e) none

7) If  $f(t) = L^{-1}\left\{\frac{1}{s^2} \cosh s\right\}$ , then  $f(2) =$

(a) 10  
B

(b) 5  
A

(c) 1

(d) 0

(e) none

8) Evaluate the given Laplace transform without evaluating the integral  $L\left\{t \int_0^t \tau e^{-\tau} d\tau\right\}$

Let  $g(t) = \int_0^t \tau e^{-\tau} d\tau$

by Theorem 4.8

$$L\{t g(t)\} = -\frac{d}{ds} G(s) \quad \text{--- (1)}$$

$$G(s) = L\{g(t)\} = L\left\{\int_0^t \tau e^{-\tau} d\tau\right\}$$

$$= \frac{F(s)}{s} \quad \text{by eq (7)/pp 218}$$

where  $f(t) = t e^{-t} \Rightarrow F(s) = -\frac{d}{ds} L\{e^{-t}\}$   
 $= -\frac{d}{ds} \frac{1}{s+1} = \frac{1}{(s+1)^2}$

Now,  $G(s) = \frac{F(s)}{s} = \frac{1}{s(s+1)^2}$  --- (2)

(1) and (2) give

$$L\{t g(t)\} = -\frac{d}{ds} \left[ \frac{1}{s(s+1)^2} \right] = -\left[ -\frac{1}{s^2} \frac{1}{(s+1)^2} - 2 \frac{1}{(s+1)^3} \frac{1}{s} \right]$$

$$= \frac{2s+1}{s^2(s+1)^3} \quad \text{--- (4)}$$

- 9) Use the Laplace transform to solve the given initial value problem  
 $y'' + 4y = \sin t U(t - 2\pi)$ ,  $y(0) = 1$ ,  $y'(0) = 0$

Take  $\mathcal{L}$

$$\mathcal{L}[y''] + 4\mathcal{L}[y] = \mathcal{L}[\sin t U(t - 2\pi)]$$

$$[s^2 Y(s) - sy(0) - y'(0)] + 4Y(s) = e^{-2\pi s} \frac{1}{s^2 + 1}$$

$$s^2 Y(s) - s + 4Y(s) = \frac{e^{-2\pi s}}{s^2 + 1}$$

$$Y(s) = \frac{e^{-2\pi s}}{(s^2 + 1)(s^2 + 4)} + \frac{s}{s^2 + 4} \quad \triangleright$$

By partial fraction  $\frac{1}{(s^2 + 1)(s^2 + 4)} = \frac{1/3}{s^2 + 1} - \frac{1/6}{s^2 + 4} \quad \triangleright$

$$Y(s) = \frac{1}{3} \underbrace{e^{-2\pi s} \frac{1}{s^2 + 1}}_{\textcircled{1}} - \frac{1}{6} \underbrace{e^{-2\pi s} \frac{1}{s^2 + 4}}_{\textcircled{2}} + \underbrace{\frac{s}{s^2 + 4}}_{\textcircled{3}} \quad (*)$$

$$\mathcal{L}^{-1} \left\{ e^{-2\pi s} \frac{1}{s^2 + 1} \right\} = u(t - 2\pi) f_1(t - 2\pi) \quad \text{when } F_1(s) = \frac{1}{s^2 + 1}$$

$$= u(t - 2\pi) \sin(t - 2\pi) \quad \text{--- } \textcircled{1} \triangleright \textcircled{3}$$

$$\mathcal{L}^{-1} \left\{ e^{-2\pi s} \frac{1}{s^2 + 4} \right\} = u(t - 2\pi) f_2(t - 2\pi) \quad \text{when } F_2(s) = \frac{1}{s^2 + 4}$$

$$= u(t - 2\pi) \sin 2(t - 2\pi) \quad \text{--- } \textcircled{2} \triangleright \textcircled{3}$$

$$\mathcal{L}^{-1} \left\{ \frac{s}{s^2 + 4} \right\} = \cos 2t \quad \text{--- } \textcircled{3} \triangleright \textcircled{3}$$

(\*) ,  $\textcircled{1}$  ,  $\textcircled{2}$  and  $\textcircled{3}$  give

$$y(t) = \cos 2t + \left[ \frac{1}{3} \sin(t - 2\pi) - \frac{1}{6} \sin 2(t - 2\pi) \right] u(t - 2\pi) \quad \triangleright \textcircled{3}$$

10) Use the Laplace transform to solve

$$y'(t) = \cos t + \int_0^t y(\tau) \cos(t-\tau) d\tau, \quad y(0) = 1$$

Taking the  $\mathcal{L}$  of the equation

$$s\mathcal{L}[y] - y(0) = \mathcal{L}[\cos t] + \mathcal{L}\left\{\int_0^t y(\tau) \cos(t-\tau) d\tau\right\}$$

$$sY(s) - 1 = \frac{s}{s^2+1} + \mathcal{L}\{y(t) * g(t)\}$$

$$\text{where } g(t) = \cos t \Rightarrow G(s) = \frac{s}{s^2+1}$$

$$sY(s) - 1 = \frac{s}{s^2+1} + Y(s) \cdot G(s) \quad \triangle$$

$$sY(s) - 1 = \frac{s}{s^2+1} + Y(s) \cdot \frac{s}{s^2+1}$$

$$Y(s) \left[ s - \frac{s}{s^2+1} \right] = 1 + \frac{s}{s^2+1}$$

$$Y(s) \left[ \frac{s^3}{s^2+1} \right] = 1 + \frac{s}{s^2+1} \Rightarrow Y(s) = \frac{s^2+1}{s^3} + \frac{1}{s^2}$$

$$Y(s) = \frac{1}{s} + \frac{1}{s^3} + \frac{1}{s^2} \quad \triangle$$

$$y(t) = \mathcal{L}^{-1}\left[\frac{1}{s}\right] + \frac{1}{2}\mathcal{L}^{-1}\left[\frac{2}{s^3}\right] + \mathcal{L}^{-1}\left[\frac{1}{s^2}\right]$$

$$y(t) = 1 + \frac{1}{2}t^2 + t \quad \triangle$$

**11)** If  $f(t) = \mathcal{L}^{-1} \left[ \frac{64s}{(s^2+4)^2} \right]$ , then find  $f(0)$

**Solution (1)** From Example 4 in page 217,

$$\mathcal{L}^{-1} \left[ \frac{2k}{(s^2+k^2)^2} \right] = (\sin kt - kt \cos kt) / 2k^3$$

with  $k=2 \Rightarrow \mathcal{L}^{-1} \left[ \frac{16}{(s^2+4)^2} \right] = \sin 2t - 2t \cos 2t \quad \text{--- (1)}$

$$\begin{aligned} \mathcal{L} [tf(t)] &= -\frac{d}{ds} \left\{ \mathcal{L} [f(t)] \right\} = -\frac{d}{ds} \left[ \frac{16}{(s^2+4)^2} \right] \\ &= - \left[ (16)(-2)(s^2+4)^{-3} (2s) \right] = \frac{64s}{(s^2+4)^3} \end{aligned}$$

So,  $\mathcal{L} [t \sin 2t - 2t^2 \cos 2t] = \frac{64s}{(s^2+4)^3}$

$\Rightarrow f(t) = \mathcal{L}^{-1} \left[ \frac{64s}{(s^2+4)^3} \right] = t \sin 2t - 2t^2 \cos 2t \Rightarrow f(0) = 0$

**Solution (2)** We will use the result  $\mathcal{L}^{-1} [F(s)G(s)] = f * g$

Let  $F(s) = \frac{16}{(s^2+4)^2}$  and  $G(s) = \frac{4s}{(s^2+4)} \Rightarrow g(t) = 4 \cos 2t$

and  $f(t) = \sin 2t - 2t \cos 2t$  (From example 4/pp 217 or (1))

Hence,  $\mathcal{L}^{-1} \left[ \frac{64s}{(s^2+4)^3} \right] = f * g = \int_0^t f(\tau) g(t-\tau) d\tau$

$$= 4 \int_0^t [\sin 2\tau - 2\tau \cos 2\tau] \cos 2(t-\tau) d\tau$$

$$= 4 \left[ \underbrace{\int_0^t \sin 2\tau \cos 2(t-\tau) d\tau}_{I_1} - \underbrace{\int_0^t 2\tau \cos 2\tau \cos 2(t-\tau) d\tau}_{I_2} \right]$$

\* Use  $\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha+\beta) + \sin(\alpha-\beta)]$  to write  $I_1$  as

$$I_1 = \frac{1}{2} \int_0^t [\sin 2t + \sin(4\tau - 2t)] d\tau = \frac{1}{2} t \sin 2t + 0 = \frac{1}{2} t \sin 2t$$

\* Use  $\cos \alpha \cos \beta = \frac{1}{2} [\cos(\alpha+\beta) + \cos(\alpha-\beta)]$  to write  $I_2$  as

$$\begin{aligned} I_2 &= - \int_0^t \tau [\cos 2t + \cos(4\tau - 2t)] d\tau \\ &= - \int_0^t \tau \cos 2t d\tau - \int_0^t \tau \cos(4\tau - 2t) d\tau = \left[ \frac{\tau^2}{2} \cos 2t \right]_0^t - \left[ \frac{\tau \sin(4\tau - 2t)}{4} + \frac{\tau \cos(4\tau - 2t)}{16} \right]_0^t \\ &= -\frac{t^2}{2} \cos 2t - \frac{1}{4} t \sin 2t \Rightarrow f(t) = 4 [I_1 + I_2] = t \sin 2t - 2t^2 \cos 2t \\ &\Rightarrow f(0) = 0 \end{aligned}$$

- 12) Let  $F(x, y, z) = xi + 2zj + yk$  represent the flow of a liquid. Find the flux of  $F$  through the surface  $S$  given by that portion of the cylinder  $y^2 + z^2 = 4$  in the first octant bounded by  $x=0, x=3, y=0, z=0$  oriented upward.

$$\text{Flux} = \iint_S F \cdot n \, ds$$


Step 1: Finding n

The surface is  $g(x, y, z) = y^2 + z^2 - 4$

$$\nabla g = (2y)j + (2z)k$$

$$\|\nabla g\| = \sqrt{4y^2 + 4z^2} = 2\sqrt{y^2 + z^2}$$

$$n = \frac{\nabla g}{\|\nabla g\|} = \frac{y}{\sqrt{y^2 + z^2}} j + \frac{z}{\sqrt{y^2 + z^2}} k \quad \triangle 4$$



Step 2:  $ds = \sqrt{1 + f_x^2 + f_y^2} dA$

The surface is  $z = \sqrt{4 - y^2} = f(x, y)$  ( $R$  is the projection on  $xy$ -plane)

$$f_x = 0, \quad f_y = \frac{-y}{\sqrt{4 - y^2}}$$

$$ds = \sqrt{1 + \frac{y^2}{4 - y^2}} dA = \sqrt{\frac{4}{4 - y^2}} = \frac{2}{\sqrt{4 - y^2}} dA$$

$$ds = \frac{2}{\sqrt{4 - y^2}} dA \quad \triangle 5$$

$$\text{Now, } F \cdot n = \frac{2zy}{\sqrt{y^2 + z^2}} + \frac{yz}{\sqrt{y^2 + z^2}} = \frac{3yz}{\sqrt{y^2 + z^2}}$$

$$\text{Flux} = \iint_S F \cdot n \, ds = \iint_R \frac{3y\sqrt{4 - y^2}}{\sqrt{y^2 + (4 - y^2)}} \cdot \frac{2}{\sqrt{4 - y^2}} dA$$

$$= \iint_R 3y \, dA = \int_0^3 \int_0^2 3y \, dy \, dx = \int_0^3 \left. \frac{3}{2} y^2 \right|_0^2 dx = \int_0^3 6 \, dx = 18 \quad \triangle 7$$

The End 😊