

## Questions from Old Exams

### 1 Section 10.1

1. Find the echelon form of the matrix  $\begin{bmatrix} 1 & -3 & 2 & -4 \\ 2 & 0 & -2 & 4 \\ 0 & 4 & 2 & 11 \end{bmatrix}$ .

2. Consider the augmented matrix of a linear system  $\begin{bmatrix} 1 & -2 & -2 & M & -1 \\ 1 & 1 & 1 & M & 2 \\ 1 & 2 & 2 & M & 1 \end{bmatrix}$ .

Which one of the following statements is TRUE?

- (a) The system is independent.
- (b) The system is dependent.
- (c) The system has the solution  $\{(2, 1, \frac{1}{2})\}$ .
- (d) The system has the solution  $\{(5, -1, -1)\}$ .
- (e) The system has no solution.

3. Which one of the following statements is TRUE about the linear system of

equations which has the augmented matrix  $\begin{bmatrix} 1 & 2 & -1 & 1 \\ 2 & 4 & -2 & 0 \\ 1 & 2 & (c-1)^2 & c+1 \end{bmatrix}$ .

- (a) The system is consistent if  $c = 0$ , with infinitely many solutions.
- (b) The system is consistent for all  $c \neq 0$ , with exactly one solution.
- (c) The system can be made consistent for suitable choice of  $c$ .
- (d) The system is inconsistent for all values of  $c$ .
- (e) The system is consistent for  $c > 0$ .

4. If the augmented matrix of a system of linear equations is  $\begin{bmatrix} 1 & 2 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 1 & 2 & 3 & -2 \\ 0 & 0 & 0 & 3 & -3 \end{bmatrix}$ ,

then find the solution set of the system.

5. If the augmented matrix of a system of linear equations is  $\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}$

then find the solution set of the system.

6. If the augmented matrix of a system of linear equations is  $\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 4 & 0 & 0 & 1 \\ 0 & 0 & 2 & 1 & 0 \\ 0 & 0 & 4 & 2 & 2 \end{bmatrix}$

then

- (a) the system has infinitely many solutions.
- (b) the system has a unique solution.
- (c) the matrix can not be the augmented matrix of a  $4 \times 4$  system.
- (d) the system has two solutions.
- (e) the system has no solution.

7. Which one of the following represents an inconsistent system?

- (a)  $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 3 \end{bmatrix}$
- (b)  $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$
- (c)  $\begin{bmatrix} 2 & 0 & 3 & 1 \\ 0 & 1 & 2 & 3 \\ 0 & 2 & 4 & 1 \end{bmatrix}$
- (d)  $\begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$
- (e)  $\begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \\ 1 & 1 & 1 & 0 \end{bmatrix}$

8. If the augmented matrix of a system of linear equations is  $\begin{bmatrix} -1 & 1 & 0 & -1 \\ 0 & 1 & -1 & 6 \\ 1 & 0 & 1 & -1 \end{bmatrix}$ ,

then find the solution set of the system.

9. If the system  $\begin{bmatrix} 1 & 1 & 1 & 2 \\ 3 & 2 & 4 & 5 \\ 2 & 1 & 1 & 6 \end{bmatrix}$  is written as  $\begin{bmatrix} 1 & m & n & 2 \\ 0 & 1 & k & 1 \\ 0 & 0 & 1 & -\frac{3}{2} \end{bmatrix}$ , then find  $mnk$ .

10. Find the solution set of  $\begin{bmatrix} 1 & 1 & 1 & -3 \\ 2 & -1 & 1 & 1 \\ 4 & 1 & 3 & 5 \end{bmatrix}$ .

11. If  $(a, b, c)$  is the solution for  $\begin{bmatrix} 1 & 2 & -1 & 5 \\ 2 & -1 & 3 & 0 \\ 1 & 1 & 1 & 2 \end{bmatrix}$ , then find the value of  $3a + 4b + c$ .

12. Given the system  $\begin{bmatrix} 1 & -2 & 4 & 2 \\ 0 & 1 & 3 & -1 \\ 0 & 2 & 6 & A \end{bmatrix}$ . Which one of the following is FALSE?

- (a) The system is inconsistent for all  $A \neq 2$ .
- (b) The system is consistent with infinitely many solutions for  $A = -2$ .
- (c) The system has no unique solution for any real number  $A$ .
- (d) The system can be made consistent or inconsistent for a suitable choice of  $A$ .
- (e) The system is consistent for any real  $A$ .

## 2 Section 10.2

1. If  $C = AB$  where  $A = \begin{bmatrix} 1 & -1 & 0 & 1 \\ 2 & 3 & -1 & 4 \\ -1 & 2 & 1 & 3 \\ 0 & 1 & -1 & 0 \end{bmatrix}$ ,  $B = \begin{bmatrix} 2 & 1 & 0 & 1 \\ 1 & -1 & 2 & -1 \\ 0 & 1 & 1 & 1 \\ 1 & -1 & 0 & 2 \end{bmatrix}$ , then find  $C_{23}$ , the third row and second column of  $C$ .
2. Given the matrices  $A = \begin{bmatrix} 3 & 2 & 0 \\ 3 & 5 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 5 & 0 \\ -3 & 1 \\ 0 & -1 \end{bmatrix}$ , and  $C = \begin{bmatrix} \frac{3}{2} & 1 \\ 0 & \frac{3}{2} \end{bmatrix}$ , then find the matrix  $AB - 2C$ .
3. If  $A = \begin{bmatrix} 0 & -2 & 7 \\ 5 & 4 & 3 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 1 \\ -1 & 5 \\ 6 & 0 \end{bmatrix}$ ,  $C = \begin{bmatrix} 40 & -10 \\ 28 & 23 \end{bmatrix}$ , and  $D = AB - C$ , then find the element in the second row and second column of the matrix  $D$ .
4. Given  $A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 2 & 0 \\ x & 2 \end{bmatrix}$ , and  $C = \begin{bmatrix} 0 & 0 \\ 6 & 0 \end{bmatrix}$ . If  $AB = 2A^2 - C$ , then find  $x$ .
5. Let  $A = \begin{bmatrix} 2 & -3 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} -2 & 3 \\ 0 & 1 \end{bmatrix}$ . If  $X$  is a  $2 \times 2$  matrix such that  $X = 2A - B$ , then
  - (a)  $X = -3B$
  - (b)  $X = 2A$
  - (c)  $X = -2B$
  - (d)  $X = 2B$
  - (e)  $X = -3B$
6. If  $A = \begin{bmatrix} 1 & 4 \\ 0 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 & 2 \\ 3 & 4 & 0 \end{bmatrix}$ , then find the element  $a_{32}$  of  $A$ .

7. Let  $A = \begin{bmatrix} -1 & 2 \\ 3 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$ , and  $C = \begin{bmatrix} a & \frac{1}{2} \\ 3 & b \end{bmatrix}$ . If  $AB = 2C$ , then find  $a$  and  $b$ .
8. If  $A = \begin{bmatrix} 1 & 1 \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix}$  and  $B = \begin{bmatrix} \frac{1}{2} & 1 \\ \frac{1}{2} & 0 \end{bmatrix}$ , then
- $A + B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
  - $A - B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
  - $AB = BA$
  - $AB = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
  - $AB = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
9. If  $A = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 1 & -2 \\ 0 & -2 & 0 \end{bmatrix}$ , then find the element in the second row and third column of  $(A^2 - A)$ .
10. If  $C = \begin{bmatrix} 2 & 2 & 1 & 1 \\ 0 & 1 & 0 & -1 \\ 1 & 0 & -1 & 0 \end{bmatrix}$  and  $D = \begin{bmatrix} 1 & 2 & -1 & 0 \\ 0 & 0 & 2 & 1 \\ 4 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \end{bmatrix}$ , then find the element in the second row and third column of  $CD$ .
11. If  $C$  is  $4 \times 3$ ,  $A$  and  $B$  are  $3 \times 4$ , then find the size of  $C \cdot (2A + 3B)$ .
12. If  $A$  and  $B$  are two matrices of size  $4 \times 3$ , then find the size of  $B^T \cdot (2A + 3B)$ .
13. If  $A = \begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 1 \\ 4 & 5 \end{bmatrix}$ , and  $C = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}$ , then find  $A^T - 2B + C^2$ .
14. If  $A = \begin{bmatrix} -1 & 0 \\ 3 & 1 \\ 0 & -2 \end{bmatrix}$ , and  $B = \begin{bmatrix} 2 & 1 \\ 1 & 0 \\ -1 & 2 \end{bmatrix}$ , then find  $(A + B) \cdot B^T$ .
15. Let  $A$  and  $B$  be square matrices of the same order and  $A^T$  is the transpose of  $A$ . Which one of the following is not always true?
- $(A^T)^T = A$
  - $(A + B)^T = A^T + B^T$

- (c)  $(A + B)^2 = A^2 + 2AB + B^2$ .  
 (d)  $(AB)^T = B^T A^T$   
 (e)  $c(A + B) = cA + cB$ , where  $c$  is a real number.

16. If  $A = \begin{bmatrix} 3 & 2 \\ x & 0 \\ -2 & -1 \end{bmatrix}$ ,  $B = \begin{bmatrix} -1 & 2 \\ 3 & 0 \end{bmatrix}$  and  $B^T A^T = 2 \begin{bmatrix} \frac{3}{2} & \frac{1}{2} & \frac{x}{2} \\ y & -1 & -2 \end{bmatrix}$ ,  
 then find  $x$  and  $y$ .

### 3 Section 10.3

1. If  $AX = B$  is the matrix equation which represents the system  $\begin{cases} 3x + 2y = 1 \\ 2x + y = 6 \end{cases}$ ,  
 then find  $X$ .
2. Given the matrices  $A = \begin{bmatrix} 2 & -1 \\ 4 & -3 \end{bmatrix}$ ,  $B = \begin{bmatrix} 2 & 1 \\ 3 & -5 \end{bmatrix}$ ,  $C = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$ ,  $X = \begin{bmatrix} x \\ y \end{bmatrix}$ . If  $(A - B)X = C$ , then  $X =$
- (a)  $\begin{bmatrix} 2 & 2 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 4 \\ 2 \end{bmatrix}$   
 (b)  $\begin{bmatrix} 2 & 2 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix}$   
 (c)  $\begin{bmatrix} 0 & -1 \\ \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 2 \end{bmatrix}$   
 (d)  $\begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix}$   
 (e)  $\begin{bmatrix} 4 \\ 2 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix}$
3. If  $A^{-1} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  is the inverse of  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ , then find  $a$  and  $c$ .
4. If the matrix  $\begin{bmatrix} 2 & 4 & 3 \\ 0 & 1 & -1 \\ 3 & 5 & 7 \end{bmatrix}$  is the multiplication inverse of  $\begin{bmatrix} 4 & -13t & -7t \\ x & 5t & yt \\ -1 & 2t & 2t \end{bmatrix}$ ,  
 then find  $x$ ,  $y$ , and  $t$ .
5. If  $a \neq 0$  and  $A = \begin{bmatrix} a & 0 & 0 \\ 0 & 2 & -4 \\ 0 & 1 & -2 \end{bmatrix}$ , then
- (a)  $AA^{-1} = I$   
 (b)  $A^{-1}A = I$

$$(c) A^{-1} = \begin{bmatrix} a^{-1} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & 1 \end{bmatrix}$$

(d)  $A^{-1}$  does not exist.

$$(e) A^{-1} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

6. If  $A$  and  $B$  are matrices of order  $n \times n$ , then

(a) if  $A^{-1}$  exists, then  $ABA^{-1} = B$  where  $B \neq I$ .

(b)  $A^{-1}$  and  $B^{-1}$  are  $(n+1) \times (n+1)$  matrices.

(c) if  $AB = O$ , then either  $A = O$  or  $B = O$ , where  $O$  is an  $n \times n$  zero matrix.

(d) if  $A^{-1}$  exists, then  $(AA^{-1})$  is the  $n \times n$  identity matrix.

(e)  $(A+2B)(A-2B) = A^2 - 4B^2$ .

7. If  $A^{-1} = \begin{bmatrix} 2 & -1 & 1 \\ -3 & 0 & 1 \\ 0 & 2 & 2 \end{bmatrix}$  and  $B^{-1} = \begin{bmatrix} 0 & -1 & 0 \\ 3 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$ , then find the element in the second row and third column of  $(AB)^{-1}$ .

8. If  $A = \begin{bmatrix} 2 & -1 \\ 3 & -2 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 3 \\ 2 & 2 \end{bmatrix}$ , and  $O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ , then

(a)  $A^{-1} = -A$ ,  $B^{-1}$  does not exist.

(b)  $A^{-1} = A$ ,  $B^{-1}$  does not exist.

(c)  $A^{-1} = A$ ,  $B^{-1} = O$ .

(d)  $A^{-1}$  does not exist,  $B^{-1} = O$ .

(e)  $A^{-1} = \begin{bmatrix} \frac{2}{7} & \frac{1}{7} \\ \frac{-3}{7} & \frac{-2}{7} \end{bmatrix}$ ,  $B^{-1}$  does not exist.

9. If  $A^{-1} = \begin{bmatrix} 2 & 0 & 1 \\ 3 & 0 & x \\ -6 & 1 & -4 \end{bmatrix}$  is the inverse of  $\begin{bmatrix} 2 & -1 & 0 \\ 0 & 2 & 1 \\ -3 & 2 & 0 \end{bmatrix}$ , then find  $x$ .

10. Given the system  $\begin{cases} 3x - 2y = 4 \\ x + 3y = 5 \end{cases}$ . Find the element in the first row and first column of the inverse of the coefficient matrix.

11. If  $A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ -1 & 1 & 1 \end{bmatrix}$  and  $A^{-1} = \begin{bmatrix} x & 3x & -2x \\ 2x & 0 & 2x \\ -x & 3x & 2x \end{bmatrix}$ , which of the following is TRUE?

- (a)  $x < -\frac{1}{2}$   
 (b)  $\frac{1}{2} < x < 1$   
 (c)  $x > 1$   
 (d)  $-\frac{1}{2} < x < \frac{1}{2}$   
 (e)  $1 < x < 2$
12. If  $A$  and  $B$  are  $n \times n$  matrices and  $A^{-1}$  and  $B^{-1}$  exist, then which one of the following is not always true?
- (a)  $(AB)^T = B^T A^T$   
 (b)  $A^{-1}$  is  $n \times n$ .  
 (c)  $AA^{-1} = I$   
 (d)  $(AB)^{-1} = A^{-1}B^{-1}$   
 (e)  $(A+B)^2 = A^2 + B^2 + AB + BA$
13. If  $\begin{bmatrix} -2 & -3 \\ -3 & -4 \end{bmatrix}$  is the inverse of  $\begin{bmatrix} 4 & n \\ m & 2 \end{bmatrix}$ , then find  $m$  and  $n$ .
14. If the matrix equation  $A^3 = I$  is true and  $A^{-1}$  exists, then  $A^{-1} =$   
 1)  $A^2$     2)  $A$     3)  $A^3$     4)  $I$     5)  $A^6$
15. If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$ , then find the element in the second row and the third column of  $A^{-1}$ .
16. Given the matrix equation  $AXC = B$ . If  $A^{-1}$  and  $C^{-1}$  exist, then  $X =$   
 (a)  $A^{-1}BC^{-1}$   
 (b)  $BA^{-1}C^{-1}$   
 (c)  $A^{-1}C^{-1}B$   
 (d)  $BC^{-1}A^{-1}$   
 (e)  $C^{-1}BA^{-1}$
17. The solution set of the matrix equation  $\begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is given by:
- (a)  $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$   
 (b)  $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$   
 (c)  $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -\frac{2}{7} & \frac{3}{7} \\ \frac{1}{7} & \frac{-2}{7} \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$

$$(d) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \begin{bmatrix} -\frac{2}{7} & \frac{3}{7} \\ \frac{1}{7} & \frac{-2}{7} \end{bmatrix}$$

$$(e) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 1 \\ \frac{1}{3} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

18. If  $A = \begin{bmatrix} 1 & 1 & 4 \\ 2 & 3 & 6 \\ -1 & -1 & 2 \end{bmatrix}$ , then find the sum of the elements in the second row of  $A^{-1}$ .

19. Suppose that  $A = \begin{bmatrix} 3 & 2 \\ 2 & 2 \end{bmatrix}$ ,  $B = \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix}$ , and  $X = \begin{bmatrix} x \\ y \end{bmatrix}$ . If  $AX = B$ , then the matrix  $X$  is equal to:

$$(a) \begin{bmatrix} 2 & 2 \\ -2 & -3 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix}$$

$$(b) \begin{bmatrix} -2 & 2 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix}$$

$$(c) \begin{bmatrix} 11 \\ 5 \end{bmatrix} \begin{bmatrix} 2 & -2 \\ -2 & 3 \end{bmatrix}$$

$$(d) \begin{bmatrix} 2 & -2 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 11 \\ 5 \end{bmatrix}$$

$$(e) \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & \frac{3}{2} \end{bmatrix}$$

20. Which of the following is TRUE for square matrices  $A$  and  $B$  which are the same size?

(a) If  $AB = O$ , then  $A = O$  or  $B = O$ .

(b)  $(A + B)^2 = A^2 + 2AB + B^2$

(c)  $(A - B)(A + B) = A^2 - B^2$

(d)  $(AB)^{-1} = B^{-1}A^{-1}$

(e)  $A(BC) = (BA)C$

21. If  $A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ , then find the element in row 2 column 3 of  $A^{-1}$ .

22. If the matrix  $M$  and its inverse are given by  $M = \begin{bmatrix} 2 & 2 & -1 \\ 0 & 3 & -1 \\ -1 & -2 & 1 \end{bmatrix}$ ,

$M^{-1} = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ x & y & z \end{bmatrix}$ , then find  $x + y + z$ .



## 4 Section 10.4

1. If  $A$  and  $B$  are two matrices of order 4 such that  $|A| = 4$  and  $|B| = 5$ , then find the value of  $|AB| - 5|B^{-1}|$ .

2. Find the value of the determinant 
$$\begin{vmatrix} 2 & 0 & 0 & 1 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 1 & 0 & 0 & 1 \end{vmatrix}.$$

3. Find the minor and the cofactor of the element 0 in the matrix 
$$\begin{bmatrix} -3 & 2 & 1 \\ -5 & 6 & 0 \\ -2 & -1 & 3 \end{bmatrix}.$$

4. If  $A$  and  $B$  are two matrices of order  $3 \times 3$  and  $|A| = 4$  and  $|B| = 5$ , then find the value of  $2|A| - |2B^{-1}|$ .

5. Find the value of the determinant 
$$\begin{vmatrix} 4 & -1 & 3 \\ 3 & 1 & 2 \\ 1 & -1 & 1 \end{vmatrix}.$$

6. If  $M = \begin{bmatrix} 5 & 6 \\ 0 & 4 \end{bmatrix}$  and  $I$  is the  $2 \times 2$  identity matrix, then find the sum of the values of  $x$  which satisfy  $\det(M - xI) = 0$ .

7. If  $Z$  is a  $5 \times 5$  matrix and  $|Z| = 3$ , then find  $|2Z^{-1}|$ .

8. Find the value of the determinant 
$$\begin{vmatrix} -1 & 2 & 2 & 3 \\ 0 & 2 & 3 & 4 \\ 0 & 2 & 6 & 6 \\ 2 & -4 & -4 & -2 \end{vmatrix}.$$

9. If  $A$  is a square matrix with inverse  $A^{-1}$  and transpose  $A^T$ , then which one of the following is always TRUE/

- (a)  $|A^T| = |A|$
- (b)  $|AA^{-1}| = 1$
- (c)  $|AA^T| = 1$
- (d)  $|A^{-1}| = |A|$
- (e)  $|A^{-1}| = |A^T|$

10. If  $\begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ 2 & 3 & 4 \end{vmatrix} = 3$ , then find  $\begin{vmatrix} 2 & 3 & 4 \\ x-4 & y-6 & z-8 \\ -2 & -2 & -2 \end{vmatrix}.$

11. Find the solution set of  $\begin{vmatrix} x & x & 0 \\ 2 & 1+x & 2 \\ -1 & 0 & x \end{vmatrix} = 0.$

12. Find the value of  $\begin{vmatrix} a & b & 0 & 0 \\ c & d & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & c & d \end{vmatrix}$ .

13. If  $A = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$ ,  $B = \begin{vmatrix} c & -2b & -3a \\ f & -2c & -3d \\ i & -2h & -3g \end{vmatrix}$  and  $|A| = 2$ , then find  $|B|$ .

14. Find the determinant  $\begin{vmatrix} x & 2 & 4 \\ 1 & 3 & 0 \\ 1 - 2x & -1 & -8 \end{vmatrix}$ .

15. If a  $3 \times 3$  matrix  $A$  with elements  $a_{ij}$  has  $a_{11} = -1$ ,  $a_{21} = 3$ , and  $a_{31} = 4$ , and the minors of  $a_{11}$ ,  $a_{21}$ , and  $a_{31}$  are 5,  $-2$ , 3 respectively, then find  $|A|$ .

16. Find the determinant  $\begin{vmatrix} \sin \theta & -\cos \theta \\ -\sin 2\theta & \cos 2\theta \end{vmatrix}$ .

17. Find the cofactor of  $x$  in  $\begin{vmatrix} -3 & 0 & -1 & 0 \\ 2 & 4 & 6 & 2 \\ 0 & x & -2 & 4 \\ 1 & 3 & 1 & 0 \end{vmatrix}$ .

18. If  $A = \begin{bmatrix} 1 & 2 & 3 \\ a & b & c \\ x & y & z \end{bmatrix}$ ,  $B = \begin{bmatrix} -1 & -2 & -6 \\ 3a & 3b & 6c \\ x & y & 2z \end{bmatrix}$ , and  $C = \begin{bmatrix} 2 & 4 & 6 \\ 2x & 2y & 2z \\ 2a & 2b & 2c \end{bmatrix}$ ,

then

- (a)  $B = -6A$ ,  $C = 2A$
- (b)  $|B| = -6|A|$ ,  $|C| = -8|A|$
- (c)  $|B| = -6|A|$ ,  $|C| = -2|A|$
- (d)  $|B| = 6|A|$ ,  $|C| = 8|A|$
- (e)  $|B| = -8|A|$ ,  $|C| = -2|A|$

19. If  $A$  is a  $3 \times 3$  matrix, then find  $|2A|$  in terms of  $|A|$ .

20. Find the cofactor of the element in the third row and second column of  $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 4 \\ 1 & 2 & 6 \end{bmatrix}$ .

21. If  $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix}$ , then find  $|A|$ .

22. If  $B = \begin{bmatrix} 4 & 2 & -1 \end{bmatrix}$  and  $A = \begin{bmatrix} 1 \\ -3 \\ 1 \end{bmatrix}$ , then find  $|A^T B^T|$ .
23. Find the solution set of  $\begin{vmatrix} 1 & 0 & 0 \\ 0 & x & 1 \\ 0 & 1 & x \end{vmatrix} = 0$ .
24. Let  $A$  and  $B$  be  $3 \times 3$  matrices. Which one of the following is FALSE?
- $(AB)^{-1} = B^{-1}A^{-1}$ .
  - $(|A| + 1)^2 = |A|^2 + 2|A| + 1$ .
  - $|A^T| = |A|$
  - $|A^{-1}| = |A|$ .
  - $|3A| = 27|A|$ .
25. If  $A = \begin{bmatrix} 0 & 1 & 2 \\ 3 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix}$ , then find  $M_{21}$  and  $C_{13}$ .
26. Find the sum of all values of  $x$  for which  $\begin{vmatrix} -1 & 3 & 0 \\ 0 & 2 & x \\ 1 & -x & 1 \end{vmatrix} = 0$ .
27. If  $\begin{vmatrix} 3 & x & u \\ 3 & y & v \\ 3 & z & w \end{vmatrix} = 1$ , then find  $\begin{vmatrix} x & z & y \\ 2 & 2 & 2 \\ u & w & v \end{vmatrix}$ .
28. If  $A$  is  $5 \times 5$  and  $|A| = 4$ , then find the value of  $2|A| + |2A^{-1}|$ .
29. If  $A = \begin{bmatrix} 3 & -1 \\ 2 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$ , then find  $|(AB)^T|$ .
30. If  $\begin{vmatrix} 1 & 0 & 1 \\ 0 & \sin \theta & \cos \theta \\ \sec \theta & -\cos \theta & \sin \theta \end{vmatrix} = 0$ ,  $0 \leq \theta \leq \pi$ , then find  $\theta$ .
31. The determinant  $\begin{vmatrix} 1 & 2 & 3 \\ a & b & c \\ 2+a & 4+b & 6+c \end{vmatrix}$  is
- equal to 0 only if  $a = b = c = 0$ .
  - equal to 0 only if  $a = -b$  and  $b = -c$ .
  - never equal to 0.
  - always equal to 0.
  - equal to zero only if  $a = -2$ ,  $b = -4$ , and  $c = -6$ .

32. If  $A$  is a  $4 \times 4$  matrix and  $|A| = \frac{3}{2}$ , then find  $\frac{1}{4}|-2A|$ .

33. If  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$ ,  $I$  is the  $2 \times 2$  identity matrix, then find  $|A - 3I|$ .

34. Find the minor  $M_{23}$  of the element  $x$  in the matrix  $\begin{bmatrix} \cos 2\theta & -\sin 2\theta & 1 \\ 1 & 1 & x \\ -\sin \theta & \cos \theta & 0 \end{bmatrix}$ .