

1. If  $\lim_{x \rightarrow 0} \frac{3^{ax} - 1}{x} = 1$ , then the value of  $a$  equals

a)  $\frac{1}{\ln 3}$

b)  $\ln 3$

c) 3

d)  $-\ln 3$

e)  $\frac{1}{3}$

2. If  $f(x) = \begin{cases} \frac{\sin(\cos x - 1)}{x} & \text{if } x \neq 0 \\ a & \text{if } x = 0 \end{cases}$ . The value of  $a$  which makes the function  $f(x)$  continuous every where is

a) 0

b)  $\pi$

c)  $-\pi$

d)  $-1$

e)  $\frac{\pi}{2}$

3. If  $\cosh x = \frac{5}{3}$  and  $x < 0$ , then  $3 \sinh x - 5 \tanh x$  is

- a) 0
- b)  $-8$
- c) 8
- d)  $-6$
- e) 6

4. If there are two tangent lines from the point  $P(0, -1)$  that touch the graph of  $f(x) = x^2$  at  $x = a$  and  $x = b$ , then  $f'(a) + f'(b) =$

- a) 0
- b) 2
- c) 4
- d) 8
- e) 16

5. Using the graph of  $f(x) = \tan x$ , the maximum value of  $\delta$  such that  $|f(x)| < 0.1$  whenever  $|x| < \delta$  is equal to

- a)  $\tan^{-1} 0.1$
- b)  $\tan^{-1}(-0.1)$
- c)  $\tan^{-1} 0.2$
- d)  $2 \tan^{-1} 0.1$
- e) 0

6. If  $G(x) = 60\sqrt{x} - 162\sqrt[3]{x}$ , then the slope of the tangent line to the graph of  $y = G'(x)$  at  $x = 1$  is

- a) 21
- b) -18
- c) 61
- d) -13
- e) 28

7. If  $h(x) = \frac{\sec x}{g(x)}$  with  $h'(\pi) = 2$  and  $g(\pi) = \sqrt{2}$ , then  $g'(\pi)$  is

- a) 4
- b)  $4 + \sqrt{2}$
- c)  $-2\sqrt{2}$
- d) 5
- e)  $2\sqrt{2}$

8. If  $f(x) = \sec x$ , then  $f''\left(\frac{\pi}{4}\right) =$

- a)  $3\sqrt{2}$
- b) 1
- c)  $\frac{12}{13}$
- d)  $\frac{5}{4}\sqrt{3}$
- e) 9

9. The slope of tangent line to the graph of the equation  $3x^{\frac{4}{3}} + xy + 3y^{\frac{4}{3}} = 59$  at the point  $(1, 8)$  is

- a)  $-\frac{4}{3}$
- b)  $\frac{19}{8}$
- c)  $-\frac{20}{9}$
- d)  $\frac{16}{7}$
- e)  $\frac{20}{9}$

10. The slope of the tangent line to the graph of  $y = (\ln x)^x$  at the point  $(e, 1)$  is

- a) 1
- b) 2
- c)  $e$
- d) 0
- e)  $-1$

11. If  $f(x) = \ln \left| \frac{(x^2 + 4)^{5/2}}{(x + \sqrt{x})^{3/2}} \right|$ , then  $f'(1) =$

a)  $-\frac{1}{8}$

b)  $\frac{5}{9}$

c)  $\frac{7}{8}$

d)  $-\frac{4}{9}$

e)  $-\frac{13}{8}$

12. A particle is moving along the curve  $y = \sqrt{x}$ . As the particle passes through the point  $(4, 2)$ , its  $x$ -coordinate increases at a rate of 3 cm/s. The rate of change of the distance from the particle to the origin at that instant is

a)  $\frac{27}{4\sqrt{5}}$

b) 9

c)  $\frac{2}{\sqrt{5}}$

d)  $\frac{2}{3\sqrt{7}}$

e)  $\frac{4}{\sqrt{11}}$

13. Using differentials or linear approximation the number  $\sqrt[3]{26}$  is estimated as

- a)  $\frac{80}{27}$
- b)  $\frac{82}{27}$
- c) 2.99
- d) 3.1
- e)  $\frac{85}{27}$

14. The value of  $\frac{d}{dx}(\sinh^{-1}(\operatorname{csch}x))$  at  $x = \ln 3$  is

- a)  $-\frac{3}{4}$
- b)  $-\frac{5}{4}$
- c)  $\frac{7}{6}$
- d)  $\frac{11}{6}$
- e)  $\frac{7}{4}$

15. The position function of a particle moving along a straight line is  $s(t) = 8t - 3t^2$  for  $t$  in  $[1, 2]$ , where  $t$  is measured in seconds and  $s$  in meters. The particle is speeding up when

a)  $\frac{4}{3} < t < 2$

b)  $1 < t < \frac{4}{3}$

c)  $1 < t < 2$

d)  $1 < t < \frac{3}{2}$

e)  $\frac{3}{2} < t < 2$

16. The radius of a circular disk is measured to be 5 cm with a maximum error in measurement of 0.1 cm. Using differentials, the maximum error in calculating circumference of the circular disk is

a)  $\frac{\pi}{5}$  cm

b)  $\pi$  cm

c)  $\frac{\pi}{10}$  cm

d)  $\frac{\pi}{2}$  cm

e)  $\frac{\pi}{50}$  cm



17. The sum of the absolute maximum value and the absolute minimum value of the function  $f(x) = 2 \sin x + \cos 2x$  on the interval  $[0, \frac{\pi}{2}]$  is

- a)  $\frac{5}{2}$
- b) 2
- c)  $\frac{3}{2}$
- d) 3
- e)  $\frac{7}{2}$

18. The sum of all critical numbers of the function  $f(x) = \frac{(x-4)^2}{\sqrt[3]{x+1}}$  is

- a) 2
- b) 1
- c) 4
- d) -2
- e) -1

19. If  $f(5) = -\frac{5}{2}$  and  $f'(x) \geq -\frac{1}{2}$  for  $3 \leq x \leq 5$ , then the largest possible value of  $f(3)$  is

a)  $-\frac{3}{2}$

b)  $\frac{1}{2}$

c)  $-\frac{1}{4}$

d)  $-\frac{2}{5}$

e) 0

20. If  $c$  is a number satisfying the conclusion of the Mean Value Theorem when applied to  $f(x) = \tan^{-1} x$  on  $[0, 1]$ , then  $\pi c^2 =$

a)  $4 - \pi$

b) 4

c)  $\pi + 1$

d)  $2\pi$

e)  $\pi - 2$

21. The function  $f(x) = \frac{x}{x^2 + 1}$  is increasing on

- a)  $(-1, 1)$
- b)  $(-\infty, -1) \cup (1, \infty)$
- c)  $(-\infty, 0)$
- d)  $(0, \infty)$
- e)  $(-\infty, \infty)$

22. If the function  $f(x) = x^3 + 2ax^2 - 3bx + 1$  has an inflection point at  $(1, 2)$ , then  $2a + b^3$  equals

- a)  $-4$
- b)  $-2$
- c)  $2$
- d)  $3$
- e)  $-1$

23. The value of the limit  $\lim_{x \rightarrow 0} (1 - \sin x)^{\frac{1}{x}}$  equals

- a)  $\frac{1}{e}$
- b)  $e$
- c) 1
- d)  $\frac{1}{\sqrt{e}}$
- e) 0

24. The slant asymptote of  $y = \frac{2x^3 + 3x^2 + 20}{x^2 + 1}$  is

- a)  $y = 2x + 3$
- b)  $y = 2x - 3$
- c)  $y = 2x + 1$
- d)  $y = 2x - 1$
- e)  $y = 2x$

25. If  $(a, b)$  is a point on the ellipse  $4x^2 + y^2 = 4$  which is farthest away from the point  $(1, 0)$ , then  $b^2 =$

a)  $\frac{32}{9}$

b)  $\frac{1}{9}$

c) 11

d) 9

e)  $\frac{29}{11}$

26. Let  $x_1 = 1$  and  $x_2 = 1.1$  be the first and second Newton's approximations of a zero of the differentiable function  $f$ . The value of  $\frac{f(1)}{f'(1)}$  is

a)  $-0.1$

b)  $0.1$

c)  $-10$

d)  $-1.1$

e)  $1.1$

27. Find the most general antiderivative of the function  $f(x) = \frac{2 \cos^2 x - 1}{\cos^2 x}$

- a)  $2x - \tan x + c$
- b)  $2x - \sec x \tan x + c$
- c)  $2x + c$
- d)  $x + c$
- e)  $2x - \sec x + c$

28. The greatest area of the rectangle that has its base on the  $x$ -axis and is inscribed in the parabola  $y = 9 - x^2$  is equal to

- a)  $12\sqrt{3}$
- b)  $6\sqrt{3}$
- c) 9
- d) 18
- e)  $\frac{2}{\sqrt{3}}$