

King Fahd University of Petroleum and Minerals
Department of Mathematical Sciences

Dr. A. Lyaghfour

MATH 301/Term 062/Hw#3(9.7)/

14. Let $\mathbf{F}(x, y, z) = yz \ln(x)\mathbf{i} + (2x - 3yz)\mathbf{j} + xy^2z^3\mathbf{k}$ be a vector function. It is clear that F is defined and differentiable at any point (x, y, z) such that $x > 0$. Moreover we have

$$\begin{aligned} \text{curl}\mathbf{F} &= \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ yz \ln(x) & 2x - 3yz & xy^2z^3 \end{vmatrix} \\ &= (2xyz^3 - (-3y))\mathbf{i} - (y^2z^3 - y \ln(x))\mathbf{j} + (2 - z \ln(x))\mathbf{k} \\ &= (2xyz^3 + 3y)\mathbf{i} + (y \ln(x) - y^2z^3)\mathbf{j} + (2 - z \ln(x))\mathbf{k}. \end{aligned}$$

$$\begin{aligned} \text{div}\mathbf{F} &= \nabla \cdot \mathbf{F} = \frac{\partial(yz \ln(x))}{\partial x} + \frac{\partial(2x - 3yz)}{\partial y} + \frac{\partial(xy^2z^3)}{\partial z} \\ &= \frac{yz}{x} - 3z + 3xy^2z^2. \end{aligned}$$

□

24. Let $\mathbf{a} = \alpha\mathbf{i} + \beta\mathbf{j} + \gamma\mathbf{k}$ be a constant vector and $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$. We would like to prove the identity

$$\nabla \cdot [(\mathbf{r} \cdot \mathbf{r})\mathbf{a}] = 2(\mathbf{r} \cdot \mathbf{a}).$$

Indeed we have

$$\begin{aligned} \nabla \cdot [(\mathbf{r} \cdot \mathbf{r})\mathbf{a}] &= \nabla \cdot [\alpha(x^2 + y^2 + z^2)\mathbf{i} + \beta(x^2 + y^2 + z^2)\mathbf{j} + \gamma(x^2 + y^2 + z^2)\mathbf{k}] \\ &= \frac{\partial(\alpha(x^2 + y^2 + z^2))}{\partial x} + \frac{\partial(\beta(x^2 + y^2 + z^2))}{\partial y} + \frac{\partial(\gamma(x^2 + y^2 + z^2))}{\partial z} \\ &= 2x\alpha + 2y\beta + 2z\gamma = 2(\mathbf{r} \cdot \mathbf{a}). \end{aligned}$$

□

28. Let $\mathbf{F} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ be a vector function and let f be a real valued function. We would like to prove the identity

$$\nabla \times (f\mathbf{F}) = f(\nabla \times \mathbf{F}) + (\nabla f) \times \mathbf{F}.$$

Indeed we have

$$\begin{aligned} \nabla \times (f\mathbf{F}) &= \nabla \times (fP\mathbf{i} + fQ\mathbf{j} + fR\mathbf{k}) \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ fP & fQ & fR \end{vmatrix} \\ &= \left(\frac{\partial(fR)}{\partial y} - \frac{\partial(fQ)}{\partial z} \right) \mathbf{i} + \left(\frac{\partial(fP)}{\partial z} - \frac{\partial(fR)}{\partial x} \right) \mathbf{j} + \left(\frac{\partial(fQ)}{\partial x} - \frac{\partial(fP)}{\partial y} \right) \mathbf{k} \\ &= f \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \mathbf{i} + f \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \mathbf{j} + f \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \mathbf{k} \\ &\quad + \left(R \frac{\partial f}{\partial y} - Q \frac{\partial f}{\partial z} \right) \mathbf{i} + \left(P \frac{\partial f}{\partial z} - R \frac{\partial f}{\partial x} \right) \mathbf{j} + \left(Q \frac{\partial f}{\partial x} - P \frac{\partial f}{\partial y} \right) \mathbf{k} \\ &= f(\nabla \times \mathbf{F}) + (\nabla f) \times \mathbf{F}. \end{aligned}$$

□