

On a class of Free Boundary Problems of type

$$\operatorname{div}(a(X)\nabla u) = -\operatorname{div}(\chi(u)H(X))$$

S. Challal and A. Lyaghfour
King Fahd University of Petroleum and Minerals
P.O. Box 728, Dhahran 31261, Saudi Arabia

Abstract

We consider a class of two dimensional free boundary problems of type $\operatorname{div}(a(X)\nabla u) = -\operatorname{div}(\chi(u)H(X))$, where H is a Lipschitz vector function satisfying $\operatorname{div}(H(X)) \geq 0$. We prove that the free boundary $\partial[u > 0] \cap \Omega$ is represented locally by a family of continuous functions.

AMS Mathematics Subject Classification: 35R35

Introduction

In [4], we studied the following problem :

$$(P_0) \left\{ \begin{array}{l} \text{Find } (u, \chi) \in H^1(\Omega) \times L^\infty(\Omega) \text{ such that :} \\ (i) \quad u \geq 0, \quad 0 \leq \chi \leq 1, \quad u(\chi - 1) = 0 \text{ a.e. in } \Omega \\ (ii) \quad \int_{\Omega} (a(X)\nabla u + \chi h(X)) \cdot \nabla \xi dX \leq 0 \\ \quad \quad \quad \forall \xi \in H^1(\Omega), \quad \xi = 0 \text{ on } \Gamma_1 \cup \Gamma_3, \quad \xi \geq 0 \text{ on } \Gamma_2, \end{array} \right.$$

where Ω is the open set $\{X = (x, y) \in \mathbb{R}^2 / y \in (a_0, b_0), \gamma_1(y) < x < \gamma_2(y)\}$ with $\gamma_1, \gamma_2 \in C^0(a_0, b_0)$, $\Gamma_1 = \{(\gamma_1(y), y) / y \in (a_0, b_0)\}$, $\Gamma_2 = \{(\gamma_2(y), y) / y \in (a_0, b_0)\}$ and $\Gamma_3 = \partial\Omega \setminus (\Gamma_1 \cup \Gamma_2)$. The two-by-two matrix $a = (a_{ij})_{i,j=1,2}$ satisfies the assumptions (1.1), (1.2), (4.1) and (4.2). The function $h : \Omega \rightarrow \mathbb{R}$ satisfies

$$\begin{aligned} 0 < \underline{h} \leq h(X) \leq \bar{h} \quad \text{for a.e. } X \in \Omega \\ h_x(X) \in L^p_{loc}, \quad p > 2, \quad h_x(X) \geq 0 \quad \text{for a.e. } X \in \Omega. \end{aligned}$$

Under these assumptions we proved that the free boundary $\partial[u > 0] \cap \Omega$ is a continuous curve $x = \Phi(y)$.

In this paper, we would like to consider the more general class of free boundary problems of type $\operatorname{div}(a(X)\nabla u) = -\operatorname{div}(\chi(u)H(X))$, where H is a Lipschitz continuous vector function with

$(\operatorname{div}H)(X) \geq 0$. Our objective is to prove that the free boundary can be parameterized by a family of continuous functions.

In the study of the problem (P_0) , the monotonicity of χ with respect to x , i.e. $\chi_x \leq 0$ in $\mathcal{D}'(\Omega)$, was essential to define the free boundary as a function $x = \Phi(y)$. In the problem we are considering, we shall prove a more general monotonicity result for χ . For this purpose we introduce, for each $h \in \pi_y(\Omega)$ and $\omega \in \pi_x(\Omega \cap [y = h])$, the differential equation $(E(\omega, h))$: $X'(t) = H(X(t))$ with the initial condition $X(0) = (\omega, h)$. We show that the mappings $T_h : (t, \omega) \mapsto X(t, \omega)$ are $C^{0,1}$ homeomorphisms from domains D_h into $T_h(D_h)$ and the family $(T_h(D_h))_h$ is a covering of Ω . Using the change of variables T_h , we prove that χ is non-increasing along the orbits of $(E(\omega, h))$. This allows us to define a local parameterization of the free boundary by a family of functions $(\phi_h)_h$.

In the first section, we state the problem. In the second section, we show the monotonicity of χ . In section 3, we define the free boundary and establish some properties. In section 4, we construct a barrier function that will be used to establish a key lemma for the proof of the continuity of the functions ϕ_h , which is done in section 5.

We end the paper with some remarks. First when H is $C^{1,1}$, T_h is a C^1 diffeomorphism and the use of this change of variables leads to a problem of type (P_0) i.e.

$$\operatorname{div}(\mathbb{A}(t, \omega)\nabla(u \circ T_h)) = -(\chi \circ T_h \cdot \mathbf{h})_t$$

with \mathbb{A} and \mathbf{h} satisfying the assumptions of [4]. However when H is only $C^{0,1}$, the matrix \mathbb{A} is not necessarily $C^{0,\alpha}$, and we are not in the framework of [4].

Finally, when $H(X) = a(X)e$, i.e. for the dam problem, we propose a proof for lemma 4.4 and thus for Theorem 5.1 which does not require the assumptions (4.1)-(4.2).

1 Statement of the problem

Let Ω be a Lipschitz bounded domain of \mathbb{R}^2 . Set $\partial\Omega = \bar{\Gamma}_1 \cup \bar{\Gamma}_2 \cup \bar{\Gamma}_3$, with Γ_1, Γ_2 and Γ_3 relatively open connected subsets of $\partial\Omega$. We are concerned by the study of the following problem :

$$(P) \left\{ \begin{array}{l} \text{Find } (u, \chi) \in H^1(\Omega) \times L^\infty(\Omega) \text{ such that :} \\ (i) \quad u \geq 0, \quad 0 \leq \chi \leq 1, \quad u(\chi - 1) = 0 \quad \text{a.e. in } \Omega \\ (ii) \quad u = \varphi \quad \text{on } \Gamma_2 \cup \Gamma_3 \\ (iii) \quad \int_{\Omega} (a(X)\nabla u + \chi H(X)) \cdot \nabla \xi dX \leq 0 \\ \quad \quad \quad \forall \xi \in H^1(\Omega), \quad \xi \geq 0 \text{ on } \Gamma_2, \quad \xi = 0 \text{ on } \Gamma_3 \end{array} \right.$$

where $a = (a_{ij})_{i,j=1,2}$ is a two-by-two matrix satisfying

$$a_{ij} \in L^\infty(\Omega), \quad |a|_\infty \leq M \tag{1.1}$$

$$a(X)\xi \cdot \xi \geq \lambda |\xi|^2 \quad \forall \xi \in \mathbb{R}^2, \quad \text{for a.e. } X \in \Omega, \tag{1.2}$$

with λ and M positive constants. φ a nonnegative Lipschitz function such that $\varphi = 0$ on Γ_2 and $\varphi > 0$ on Γ_3 . The function $H = (H_1, H_2) : \Omega \rightarrow \mathbb{R}^2$ satisfies for positive constants \underline{h} and \bar{h} :

$$|H_1(X)| \leq \bar{h}, \quad 0 < \underline{h} \leq H_2(X) \leq \bar{h} \quad \text{for a.e. } X \in \Omega \tag{1.3}$$

$$\operatorname{div}(H(X)) \in L^\infty(\Omega), \quad \operatorname{div}(H(X)) \geq 0 \quad \text{for a.e. } X \in \Omega. \tag{1.4}$$

The existence of a solution of (P) is classical. We start by giving the following properties

Proposition 1.1. *Let (u, χ) be a solution of (P). We have*

i)
$$\operatorname{div}(a(X)\nabla u) = -\operatorname{div}(\chi H(X)) \quad \text{in } \mathcal{D}'(\Omega). \quad (1.5)$$

ii)
$$\operatorname{div}(\chi H(X)) - \chi([u > 0])\operatorname{div}(H(X)) \leq 0 \quad \text{in } \mathcal{D}'(\Omega). \quad (1.6)$$

iii) $u \in C_{loc}^{0,\alpha}(\Omega \cup \Gamma_2 \cup \Gamma_3)$ for all $\alpha \in (0, 1)$.

iv) $[u > 0]$ is an open set.

v) If $a \in C_{loc}^{0,\alpha}(\Omega)$, then $u \in C_{loc}^{1,\alpha}([u > 0])$.

Proof. i) This is an immediate consequence of taking $\pm\xi$, with $\xi \in \mathcal{D}(\Omega)$, as test functions for (P).

ii) Let $\xi \in \mathcal{D}(\Omega)$, $\xi \geq 0$ and let $F_\epsilon(s) = \min\left(\frac{s^+}{\epsilon}, 1\right)$, $\epsilon > 0$. Taking $\pm F_\epsilon(u)\xi$ as test functions for (P), we obtain

$$\int_{\Omega} (a(X)\nabla u + \chi H(X)) \cdot \nabla(F_\epsilon(u)\xi) dX = 0$$

which can be written by taking into account (P)(i), (1.2) and the fact that F_ϵ is nondecreasing

$$\int_{\Omega} [F_\epsilon(u)a(X)\nabla u \cdot \nabla \xi - (F_\epsilon(u)\xi)\operatorname{div}H(X)] dX \leq 0.$$

Letting $\epsilon \rightarrow 0$, we obtain

$$\int_{\Omega} [a(X)\nabla u \cdot \nabla \xi - \chi([u > 0])\operatorname{div}H(X)\xi] dX \leq 0.$$

Combining the last inequality and (1.5), we get (1.6).

iii) This is a consequence of (P)(ii), (1.5) and the regularity theory of elliptic problems (see [5], Theorem 8.34 for example).

iv) This a consequence of iii).

v) Using (P)(i) and (1.5) we obtain $\operatorname{div}(a(X)\nabla u) = -\operatorname{div}(H(X))$ in $\mathcal{D}'([u > 0])$. Hence the result becomes a consequence of the regularity theory of elliptic problems (see [5], Corollary 8.36). \square

2 A monotonicity property of χ

In all what follows, we shall assume that

$$H \in C^{0,1}(\overline{\Omega}). \quad (2.1)$$

We consider the following differential system

$$(E(\omega, h)) \begin{cases} X'(t, \omega, h) &= H(X(t, \omega, h)) \\ X(0, \omega, h) &= (\omega, h) \end{cases}$$

where $h \in \pi_y(\Omega)$ and $\omega \in \pi_x(\Omega \cap [y = h])$. π_x and π_y are respectively the orthogonal projections on the x and y axes.

By the classical theory of ordinary differential equations, there exists a unique maximal solution $X(., \omega, h)$ of $E(\omega, h)$ defined on $(\alpha_-(\omega, h), \alpha_+(\omega, h))$ and continuous on the open set

$$\{(t, \omega, h) / \alpha_-(\omega, h) < t < \alpha_+(\omega, h), h \in \pi_y(\Omega), \omega \in \pi_x(\Omega \cap [y = h])\}.$$

Since H is bounded and continuous on $\bar{\Omega}$, $X(., \omega, h)$ is defined on $[\alpha_-(\omega, h), \alpha_+(\omega, h)]$ (see Lemma 2.1 p 16 [6]). Moreover by Corollary 7.7 p. 103 of [1], we know that $X(\alpha_-(\omega, h), \omega, h) \in \partial\Omega \cap [y < h]$, $X(\alpha_+(\omega, h), \omega, h) \in \partial\Omega \cap [y > h]$ (see Figure 1).

For simplicity we will denote in the sequel $X(t, \omega, h)$, $\alpha_-(\omega, h)$ and $\alpha_+(\omega, h)$ respectively by $X(t, \omega)$, $\alpha_-(\omega)$ and $\alpha_+(\omega)$. We shall also denote by $\gamma(\omega)$ the orbit of $X(., \omega)$.

Remark 2.1. *Note that α_+ and α_- are bounded. Indeed, we have by (1.3)*

$$\begin{aligned} \underline{h}\alpha_+(\omega) &\leq \int_0^{\alpha_+(\omega)} H_2(X(s, \omega)) ds = X_2(\alpha_+(\omega), \omega) - h \\ X_2(\alpha_-(\omega), \omega) - h &= - \int_{\alpha_-(\omega)}^0 H_2(X(s, \omega)) ds \leq \underline{h}\alpha_-(\omega). \end{aligned}$$

Hence

$$\frac{1}{\underline{h}} \left(\inf_{y \in \pi_y(\Omega)} y - h \right) \leq \alpha_-(\omega) < 0 < \alpha_+(\omega) \leq \frac{1}{\underline{h}} \left(\sup_{y \in \pi_y(\Omega)} y - h \right).$$

Definition 2.1. *For each $h \in \pi_y(\Omega)$ we define the set*

$$D_h = \{(t, \omega) / \omega \in \pi_x(\Omega \cap [y = h]), t \in (\alpha_-(\omega), \alpha_+(\omega))\}$$

and consider the mapping

$$\begin{aligned} T_h : D_h &\longrightarrow T_h(D_h) \\ (t, \omega) &\longmapsto T_h(t, \omega) = (T_h^1, T_h^2)(t, \omega) = X(t, \omega). \end{aligned}$$

Clearly each $(x, y) \in \Omega$ can be written as $(x, y) = X(0, \omega) = T_h(0, \omega)$ with $\omega = x$ and $h = y$. So

$$\Omega = \bigsqcup_{h \in \pi_y(\Omega)} T_h(D_h). \quad (2.2)$$

Moreover




Figure 1

Proposition 2.1.

T_h is continuous and one to one.

Proof. By the previous remarks on the regularity of X , we have $T_h \in C^0(D_h)$. Now let $(t_1, \omega_1), (t_2, \omega_2) \in D_h$ such that $T_h(t_1, \omega_1) = T_h(t_2, \omega_2)$ i.e. $X(t_1, \omega_1) = X(t_2, \omega_2)$. Then we consider the following ordinary differential equation

$$\begin{cases} Z'(t) = H(Z(t)) \\ Z(0) = X(t_1, \omega_1). \end{cases}$$

Clearly we have

$$\begin{aligned} Z(t) &= X(t + t_1, \omega_1) = X(t + t_2, \omega_2) \\ \forall t &\in [\alpha_-(\omega_1) - t_1, \alpha_+(\omega_1) - t_1] = [\alpha_-(\omega_2) - t_2, \alpha_+(\omega_2) - t_2]. \end{aligned}$$

In particular $t_2 - t_1 \in (\alpha_-(\omega_2), \alpha_+(\omega_2))$. Indeed we have

$$\alpha_-(\omega_1) - t_1 = \alpha_-(\omega_2) - t_2 \quad \text{and} \quad \alpha_+(\omega_1) - t_1 = \alpha_+(\omega_2) - t_2.$$

Then since $\alpha_-(\omega_1) < 0$ and $\alpha_+(\omega_1) > 0$, we get

$$\alpha_-(\omega_2) < \alpha_-(\omega_2) - \alpha_-(\omega_1) = t_2 - t_1 = \alpha_+(\omega_2) - \alpha_+(\omega_1) < \alpha_+(\omega_2).$$

So we can write $Z(-t_1) = X(0, \omega_1) = X(t_2 - t_1, \omega_2)$, i.e.

$$(\omega_1, h) = (\omega_2, h) + \int_0^{t_2-t_1} H(X(s, \omega_2)) ds.$$

Therefore

$$\int_0^{t_2-t_1} H_2(X(s, \omega_2)) ds = 0$$

which leads by (1.3) to $t_2 = t_1$. We then deduce that $\omega_1 = \omega_2$. \square

Proposition 2.2.

$$T_h \quad \text{and} \quad T_h^{-1} \quad \text{are } C^{0,1}.$$

Proof. The proof is done in several steps. For the Lipschitz continuity of T_h , we refer to [?], Theorem 8.3 p 110.

Step 1. Extension.

Since $H \in C^{0,1}(\bar{\Omega})$, there exists by Kirszbraun's theorem (see [2], Theorem 2.10.43 p.210) an extension $\tilde{H} \in C^{0,1}(\mathbb{R}^2)$ of H with the same Lipschitz constant L . Then

$$\begin{aligned} \bar{H} &= (\min(\bar{h}, \max(\tilde{H}_1, -\bar{h})), \min(\bar{h}, \max(\tilde{H}_2, \underline{h}))) \in C^{0,1}(\mathbb{R}^2) \\ \text{with } |\bar{H}_1| &\leq \bar{h} \quad \text{and} \quad \underline{h} \leq \bar{H}_2 \leq \bar{h}. \end{aligned}$$

Step 2. Regularization.

Let $H_\epsilon = \rho_\epsilon * \bar{H}$, where ρ_ϵ is the usual mollifier function. Then, it is well known that $H_\epsilon \in C^\infty(\mathbb{R}^2)$ and satisfies

$$\begin{cases} |H_\epsilon^1(X)| \leq \bar{h}, & \underline{h} \leq H_\epsilon^2(X) \leq \bar{h} & \forall X \in \mathbb{R}^2 \\ H_\epsilon \longrightarrow \bar{H} & \text{uniformly on each compact set of } \mathbb{R}^2 & \text{as } \epsilon \rightarrow 0 \\ \|\nabla H_\epsilon\|_{L^\infty(\mathbb{R}^2)} \leq \|\nabla \bar{H}\|_{L^\infty(\mathbb{R}^2)} \leq L. \end{cases}$$

Now, for $(\omega, h) \in \mathbb{R}^2$, let X_ϵ and \bar{X} be respectively the unique solutions of the differential equations

$$\begin{cases} X'_\epsilon(t, \omega) = H_\epsilon(X_\epsilon(t, \omega)) \\ X_\epsilon(0, \omega) = (\omega, h) \end{cases} \quad \text{and} \quad \begin{cases} \bar{X}'(t, \omega) = \bar{H}(\bar{X}(t, \omega)) \\ \bar{X}(0, \omega) = (\omega, h). \end{cases}$$

X_ϵ and \bar{X} are defined on the maximal interval $(-\infty, +\infty)$. Moreover X_ϵ is C^∞ with respect to $t \in \mathbb{R}$ and the initial value $(w, h) \in \mathbb{R}^2$.

Step 3. Local uniform convergence.

Let K be a compact set of \mathbb{R}^2 . There exists $T > 0$, $\omega_1, \omega_2 \in \mathbb{R}$ such that $K \subset \subset [-T, T] \times [\omega_1, \omega_2] = K'$. For each $(t, \omega) \in K$, we have

$$\begin{aligned} |X_\epsilon(t, \omega) - \bar{X}(t, \omega)| &= \left| \int_0^t (H_\epsilon(X_\epsilon(s, \omega)) - \bar{H}(\bar{X}(s, \omega))) ds \right| \\ &\leq \left| \int_0^t (H_\epsilon(X_\epsilon(s, \omega)) - H_\epsilon(\bar{X}(s, \omega))) ds \right| + \left| \int_0^t (H_\epsilon(\bar{X}(s, \omega)) - \bar{H}(\bar{X}(s, \omega))) ds \right| \\ &\leq \left| \int_0^t L |X_\epsilon(s, \omega) - \bar{X}(s, \omega)| ds \right| + |t| |H_\epsilon - \bar{H}|_{\infty, \bar{X}(K')} \end{aligned}$$

By Gronwall's Lemma (see [?], p 90), we obtain

$$|X_\epsilon(t, \omega) - \bar{X}(t, \omega)| \leq |t| |H_\epsilon - \bar{H}|_{\infty, \bar{X}(K')} \exp(L|t|).$$

So we have with $C = T \exp(LT)$

$$|X_\epsilon - \bar{X}|_{\infty, K} \leq C |H_\epsilon - \bar{H}|_{\infty, \bar{X}(K')} \rightarrow 0 \quad \text{when } \epsilon \rightarrow 0.$$

Step 4. $X_\epsilon : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a C^1 diffeomorphism.

• $X_\epsilon(\mathbb{R}^2) = \mathbb{R}^2$: Indeed let $(x_0, y_0) \in \mathbb{R}^2$ and let $Z = (Z_1, Z_2)$ be the unique maximal solution of the following differential equation

$$\begin{cases} Z'(t) = H_\epsilon(Z(t)) \\ Z(0) = (x_0, y_0). \end{cases}$$

It is not difficult to see that Z is defined on $(-\infty, +\infty)$ and that $\lim_{t \rightarrow \pm\infty} Z_2(t) = \pm\infty$. Moreover since $Z_2'(t) = H_\epsilon^2(Z(t)) \geq \underline{h} > 0$, we deduce that Z_2 is bijective from \mathbb{R} into \mathbb{R} . Therefore there exists $t_0 \in \mathbb{R}$ such that $Z_2(t_0) = h$.

Let $\omega_0 = Z_1(t_0)$. Then $Z(t_0) = (\omega_0, h)$ and it is easy to verify that $X_\epsilon(t, \omega_0) = Z(t + t_0)$. In particular $X_\epsilon(-t_0, \omega_0) = Z(0) = (x_0, y_0)$.

• Since X_ϵ is onto, it suffices then to verify that $\det(\mathcal{J}X_\epsilon)$ does not vanish. Here we denote by $\mathcal{J}F$ the Jacobian matrix of the mapping F and by $\det(\mathcal{J}F)$ the determinant of $\mathcal{J}F$. One can easily check that

$$\begin{aligned} Y_h^\epsilon(t, \omega) &= \det(\mathcal{J}X_\epsilon) = H_\epsilon^1(X_\epsilon(t, \omega)) \frac{\partial X_{2\epsilon}}{\partial \omega} - H_\epsilon^2(X_\epsilon(t, \omega)) \frac{\partial X_{1\epsilon}}{\partial \omega}, \\ \frac{\partial Y_h^\epsilon}{\partial t}(t, \omega) &= Y_h^{\epsilon\epsilon}(t, \omega) \cdot (\text{div}(H_\epsilon))(X_\epsilon(t, \omega)). \end{aligned}$$

Therefore

$$Y_h^\epsilon(t, \omega) = Y_h^\epsilon(0, \omega) \cdot \exp\left(\int_0^t \{\text{div}(H_\epsilon)\}(X_\epsilon(s, \omega)) ds\right). \quad (2.3)$$

Since $Y_h^\epsilon(0, \omega) = -H_\epsilon^2(X_\epsilon(0, \omega)) = -H_\epsilon^2(\omega, h) < 0$, we get $Y_h^\epsilon(t, \omega) < 0 \forall (t, \omega) \in \mathbb{R}^2$.

Step 5. We have :

$$\|\mathcal{J}X_\epsilon^{-1}(x, y)\|_\infty \leq \frac{1}{\underline{h}} \left(\exp\left(\frac{L|y-h|}{\underline{h}}\right) + \bar{h} \right) \quad \forall (x, y) \in \mathbb{R}^2.$$

Indeed, we have for $(t, \omega) = X_\epsilon^{-1}(x, y)$

$$\mathcal{J}X_\epsilon^{-1}(x, y) = \frac{1}{Y_h^\epsilon(t, \omega)} \begin{pmatrix} \frac{\partial X_{2\epsilon}}{\partial \omega}(t, \omega) & -\frac{\partial X_{1\epsilon}}{\partial \omega}(t, \omega) \\ -H_\epsilon^2(X_\epsilon(t, \omega)) & H_\epsilon^1(X_\epsilon(t, \omega)) \end{pmatrix}.$$

$$|H_\epsilon^i(X_\epsilon(t, \omega))| \leq \bar{h}, \quad H_\epsilon^2(X_\epsilon(t, \omega)) \geq \underline{h}, \quad \text{and} \quad \text{div}(H_\epsilon) \geq 0.$$

It follows that

$$\frac{1}{|Y_h^\epsilon(t, \omega)|} = \frac{1}{H_\epsilon^2(\omega, h)} \exp\left(-\int_0^t \{\text{div}(H_\epsilon)\}(X_\epsilon(s, \omega)) ds\right) \leq \frac{1}{\underline{h}}.$$

We claim that $\left| \frac{\partial X_\epsilon}{\partial \omega} \right| \leq \exp(L|t|)$. Indeed for $\omega_1, \omega_2 \in \mathbb{R}$, we have

$$\begin{aligned} |X_\epsilon(t, \omega_1) - X_\epsilon(t, \omega_2)| &= \left| (\omega_1 - \omega_2, 0) + \int_0^t (H_\epsilon(X_\epsilon(s, \omega_1)) - H_\epsilon(X_\epsilon(s, \omega_2))) ds \right| \\ &\leq |\omega_1 - \omega_2| + L \left| \int_0^t |X_\epsilon(s, \omega_1) - X_\epsilon(s, \omega_2)| ds \right|. \end{aligned}$$

By Gronwall's Lemma, we obtain

$$|X_\epsilon(t, \omega_1) - X_\epsilon(t, \omega_2)| \leq |\omega_1 - \omega_2| \exp(L|t|).$$

Now we conclude that

$$\begin{aligned} \|\mathcal{J}X_\epsilon^{-1}(x, y)\|_\infty &= \max \left\{ \frac{1}{|Y_h^\epsilon|} \left(\left| \frac{\partial X_{2\epsilon}}{\partial \omega} \right| + |H_\epsilon^2 \circ X_\epsilon| \right), \frac{1}{|Y_h^\epsilon|} \left(\left| \frac{\partial X_{1\epsilon}}{\partial \omega} \right| + |H_\epsilon^1 \circ X_\epsilon| \right) \right\} (t, \omega) \\ &\leq \frac{1}{\underline{h}} (\exp(L|t|) + \bar{h}) \leq \frac{1}{\underline{h}} \left(\exp\left(\frac{L|y-h|}{\underline{h}}\right) + \bar{h} \right) \end{aligned}$$

since

$$|y-h| = \left| \int_0^t H_\epsilon^2(X_\epsilon(s, \omega)) ds \right| \geq \underline{h}|t|.$$

Step 6. X_ϵ^{-1} is uniformly Lipschitz continuous on each compact set with a Lipschitz constant independent of ϵ .

Let K be a compact set of \mathbb{R}^2 and $(x_1, y_1), (x_2, y_2)$ be two points in K . We denote by $|(x, y)|_\infty = \max(|x|, |y|)$. Then we have

$$\begin{aligned}
|X_\epsilon^{-1}(x_1, y_1) - X_\epsilon^{-1}(x_2, y_2)|_\infty &= \left| \int_0^1 \frac{d}{d\tau} X_\epsilon^{-1}(\tau(x_1, y_1) + (1-\tau)(x_2, y_2)) d\tau \right|_\infty \\
&= \left| \int_0^1 \mathcal{J} X_\epsilon^{-1}(\tau(x_1, y_1) + (1-\tau)(x_2, y_2)) \cdot (x_1 - x_2, y_1 - y_2) d\tau \right|_\infty \\
&\leq \int_0^1 |\mathcal{J} X_\epsilon^{-1}(\tau(x_1, y_1) + (1-\tau)(x_2, y_2))|_\infty \cdot |(x_1, y_1) - (x_2, y_2)|_\infty d\tau \\
&\leq \left(\frac{1}{\underline{h}} \int_0^1 (\exp(\frac{L}{\underline{h}} |(1-\tau)y_2 + \tau y_1 - h|) + \bar{h}) d\tau \right) |(x_1, y_1) - (x_2, y_2)|_\infty \\
&\leq c(K) |(x_1, y_1) - (x_2, y_2)|_\infty,
\end{aligned}$$

with $c(K) = \frac{1}{\underline{h}} (\exp(\frac{L}{\underline{h}} [m + h]) + \bar{h})$ and $m = \max\{|y| / y \in \pi_y(K)\}$.

Step 7. Conclusion.

There exists a subsequence $(X_{\epsilon_n})_{n \geq 0}$ such that $(X_{\epsilon_n}^{-1})_n$ converges uniformly to an element $X^* \in C_{loc}^{0,1}(\mathbb{R}^2)$ on each compact set of \mathbb{R}^2 .

We claim that $X^* = \bar{X}^{-1}$. Indeed we have

$$X_\epsilon \circ X_\epsilon^{-1}(x, y) = (x, y) \quad \text{and} \quad X_\epsilon^{-1} \circ X_\epsilon(t, \omega) = (t, \omega) \quad \forall (x, y), (t, \omega) \in \mathbb{R}^2.$$

Passing to the limit, we obtain

$$\bar{X} \circ X^*(x, y) = (x, y) \quad \text{and} \quad X^* \circ \bar{X}(t, \omega) = (t, \omega) \quad \forall (x, y), (t, \omega) \in \mathbb{R}^2.$$

Since $\bar{X}|_{D_h} = X = T_h$, we have $T_h^{-1} = \bar{X}^{-1}|_{T_h(D_h)} \in C^{0,1}(T_h(D_h))$. □

Now we have the following Proposition

Proposition 2.3. *Let $X(\cdot, \omega)$ be the maximal solution of $E(\omega, h)$. We have*

i)

$$\mathcal{J}T_h = \begin{pmatrix} H_1(X(t, \omega)) & \frac{\partial X_1}{\partial \omega}(t, \omega) \\ H_2(X(t, \omega)) & \frac{\partial X_2}{\partial \omega}(t, \omega) \end{pmatrix} \in L^\infty(D_h)$$

$$Y_h(t, \omega) = \det \mathcal{J}T_h = H_1(X(t, \omega)) \frac{\partial X_2}{\partial \omega}(t, \omega) - H_2(X(t, \omega)) \frac{\partial X_1}{\partial \omega}(t, \omega) \text{ in } L^\infty(D_h).$$

ii) $\frac{\partial Y_h}{\partial t}(t, \omega) = Y_h(t, \omega) (\operatorname{div} H)(X(t, \omega)) \quad \text{a.e. in } D_h.$

iii) $Y_h(t, \omega) = -H_2(\omega, h) \exp\left(\int_0^t (\operatorname{div} H)(X(s, \omega)) ds\right) \quad \text{a.e. in } D_h.$

iv) $\underline{h} \leq -Y_h(t, \omega) \leq C\bar{h}, \quad C > 0.$

Proof. i) Note that since $T_h \in C^{0,1}(D_h)$, we have $T_h \in W^{1,\infty}(D_h)$ and therefore we can talk about $\mathcal{J}T_h$. The formula is trivial.

ii) Given that H, T_h, T_h^{-1} are $C^{0,1}$, we can use the chain rule for $H_i \circ X$ (see [7]) to get

$$\frac{\partial(H_i \circ X)}{\partial t} = H_1 \frac{\partial H_i}{\partial x} + H_2 \frac{\partial H_i}{\partial y}. \quad (2.4)$$

Moreover $H_\epsilon, X_\epsilon \in C^\infty(\mathbb{R}^2)$ and $X_\epsilon(t, \omega) = (\omega, h) + \int_0^t H_\epsilon(X_\epsilon(s, \omega)) ds$. So we have

$$\frac{\partial X_\epsilon}{\partial \omega}(t, \omega) = (1, 0) + \int_0^t \left(\frac{\partial X_{1\epsilon}}{\partial \omega}(s, \omega) \frac{\partial H_\epsilon}{\partial x}(X_\epsilon(s, \omega)) + \frac{\partial X_{2\epsilon}}{\partial \omega}(s, \omega) \frac{\partial H_\epsilon}{\partial y}(X_\epsilon(s, \omega)) \right) ds.$$

Since H_ϵ and X_ϵ converge uniformly to H and X respectively in Ω and D_h , $\frac{\partial X_\epsilon}{\partial \omega}$ and ∇H_ϵ converge to $\frac{\partial X}{\partial \omega}$ and ∇H respectively in $L^p(D_h)$ and $L^p(\Omega)$ for each $p \geq 1$, we obtain for a.e. $(t, \omega) \in D_h$, by letting $\epsilon \rightarrow 0$

$$\frac{\partial X}{\partial \omega}(t, \omega) = (1, 0) + \int_0^t \left(\frac{\partial X_1}{\partial \omega}(s, \omega) \frac{\partial H}{\partial x}(X(s, \omega)) + \frac{\partial X_2}{\partial \omega}(s, \omega) \frac{\partial H}{\partial y}(X(s, \omega)) \right) ds. \quad (2.5)$$

It follows from (2.5) that

$$\frac{\partial^2 X}{\partial t \partial \omega}(t, \omega) = \frac{\partial X_1}{\partial \omega}(t, \omega) \cdot \frac{\partial H}{\partial x}(X(t, \omega)) + \frac{\partial X_2}{\partial \omega}(t, \omega) \cdot \frac{\partial H}{\partial y}(X(t, \omega)) \text{ in } L^\infty(D_h). \quad (2.6)$$

Now, since $H_i \circ X \in W^{1,\infty}(D_h)$ and $\frac{\partial^2 X_i}{\partial t \partial \omega} \in L^\infty(D_h)$, we obtain

$$\frac{\partial}{\partial t} \left(H_i \circ X \cdot \frac{\partial X_i}{\partial \omega} \right) = \frac{\partial}{\partial t} (H_i \circ X) \cdot \frac{\partial X_i}{\partial \omega} + (H_i \circ X) \cdot \frac{\partial^2 X_i}{\partial t \partial \omega}. \quad (2.7)$$

Using (2.4)-(2.7), we obtain

$$\begin{aligned} \frac{\partial Y_h}{\partial t}(t, \omega) &= \left(H_1(X(t, \omega)) \frac{\partial H_1}{\partial x}(X(t, \omega)) + H_2(X(t, \omega)) \frac{\partial H_1}{\partial y}(X(t, \omega)) \right) \frac{\partial X_2}{\partial \omega}(t, \omega) \\ &\quad - \left(H_1(X(t, \omega)) \frac{\partial H_2}{\partial x}(X(t, \omega)) + H_2(X(t, \omega)) \frac{\partial H_2}{\partial y}(X(t, \omega)) \right) \frac{\partial X_1}{\partial \omega}(t, \omega) \\ &\quad + H_1(X(t, \omega)) \frac{\partial^2 X_2}{\partial t \partial \omega}(t, \omega) - H_2(X(t, \omega)) \frac{\partial^2 X_1}{\partial t \partial \omega}(t, \omega) \\ &= \left(H_1(X(t, \omega)) \frac{\partial X_2}{\partial \omega}(t, \omega) - H_2(X(t, \omega)) \frac{\partial X_1}{\partial \omega}(t, \omega) \right) \cdot \left(\frac{\partial H_1}{\partial x} + \frac{\partial H_2}{\partial y} \right) (X(t, \omega)) \\ &= Y_h(t, \omega) (\operatorname{div} H)(X(t, \omega)). \end{aligned}$$

iii) By using the product formula and chain rule, we obtain

$$\begin{aligned} \frac{\partial}{\partial t} \left((Y_h(t, \omega) \cdot \exp(-\int_0^t (\operatorname{div} H)(X(s, \omega)) ds)) \right) &= \frac{\partial Y_h}{\partial t} \cdot \exp(-\int_0^t (\operatorname{div} H)(X(s, \omega)) ds) \\ &+ Y_h(t, \omega) \cdot (-\operatorname{div} H)(X(t, \omega)) \exp(-\int_0^t (\operatorname{div} H)(X(s, \omega)) ds) = 0. \end{aligned}$$

Then $Y_h(t, \omega) \cdot \exp(-\int_0^t (\operatorname{div} H)(X(s, \omega)) ds) = Cst = Y_h(0, \omega)$ which exists because $Y_h \in C^0(\alpha_-(\omega), \alpha_+(\omega))$. Since we have $\frac{\partial X}{\partial \omega}(0, \omega) = (1, 0)$, then $Y_h(0, \omega) = -H_2(X(0, \omega)) = -H_2(\omega, h)$.

iv) Since $0 \leq \operatorname{div} H \leq L$, it follows that $\left| \int_0^t (\operatorname{div} H)(X(s, \omega)) ds \right| \leq 2L|t| \leq 2L \max(\alpha_+(\omega), -\alpha_-(\omega)) \leq L'$. We deduce, since $\underline{h} \leq H_2(\omega, h) \leq \bar{h}$, that $\underline{h} \leq -Y_h(t, \omega) \leq \bar{h} \exp(L')$. \square

Now we can prove the main result of this section.

Theorem 2.1. *Let (u, χ) be a solution of (P). We have for each $h \in \pi_y(\Omega)$*

$$\frac{\partial}{\partial t} (\chi \circ T_h) \leq 0 \quad \text{in } \mathcal{D}'(D_h).$$

Proof. Let $\varphi \in \mathcal{D}(D_h)$, $\varphi \geq 0$. By (1.6), we have

$$\int_{T_h(D_h)} (-\chi H(X) \cdot \nabla(\varphi \circ T_h^{-1}) - \chi([u > 0]) \operatorname{div} H(X) \cdot \varphi \circ T_h^{-1}) dX \leq 0.$$

Since $T_h, T_h^{-1} \in C^{0,1}$, we can use T_h as a change of variables (see [7]) to obtain

$$\int_{D_h} \left(-\chi \circ T_h \frac{\partial \varphi}{\partial t} - \chi([u \circ T_h > 0]) (\operatorname{div} H) \circ T_h \cdot \varphi \right) (-Y_h(t, \omega)) dt d\omega \leq 0.$$

Given that $\frac{\partial Y_h}{\partial t} = Y_h \cdot (\operatorname{div} H) \circ T_h$, we obtain

$$\begin{aligned} \int_{D_h} \chi \circ T_h \frac{\partial(-Y_h \cdot \varphi)}{\partial t} dt d\omega &= \int_{D_h} \chi \circ T_h \frac{\partial \varphi}{\partial t} (-Y_h) + \chi \circ T_h \cdot (\operatorname{div} H) \circ T_h \cdot \varphi \cdot (-Y_h) dt d\omega \\ &\geq \int_{D_h} (\chi \circ T_h - \chi([u \circ T_h > 0])) \cdot (\operatorname{div} H) \circ T_h \cdot \varphi \cdot (-Y_h) dt d\omega \geq 0. \end{aligned}$$

By approximation the last inequality remains valid for all nonnegative functions φ with compact support and such that $\varphi_t \in L^1(D_h)$. Since $Y_h \in L^\infty(D_h)$ and does not vanish, one can choose $\varphi = -\frac{\psi}{Y_h}$, with $\psi \in \mathcal{D}(D_h)$ and $\psi \geq 0$. Thus we get the result. \square

3 Definition of the Free Boundary and some Technical Results

In this section, we use the monotonicity result of the previous section and the continuity of u to define the free boundary. We also give some other results. First, we have the following key proposition.

Proposition 3.1. *Let (u, χ) be a solution of (P) and $X_0 = (x_0, y_0) = T_h(t_0, \omega_0) \in T_h(D_h)$.*

i) If $u(X_0) = uoT_h(t_0, \omega_0) > 0$, then there exists $\epsilon > 0$ such that

$$uoT_h(t, \omega) > 0 \quad \forall (t, \omega) \in C_\epsilon = \{(t, \omega) \in D_h / |\omega - \omega_0| < \epsilon, t < t_0 + \epsilon\}.$$

ii) If $u(X_0) = uoT_h(t_0, \omega_0) = 0$, then $uoT_h(t, \omega_0) = 0 \quad \forall t \geq t_0$.

Proof. It suffices to verify *i*). By continuity, there exists $\epsilon > 0$ such that

$$uoT_h(t, \omega) > 0 \quad \forall (t, \omega) \in (t_0 - \epsilon, t_0 + \epsilon) \times (\omega_0 - \epsilon, \omega_0 + \epsilon) = Q_\epsilon.$$

Then $\chi oT_h(t, \omega) = 1$ for a.e. $(t, \omega) \in Q_\epsilon$. By Theorem 2.1 and since $\chi oT_h \leq 1$, we get $\chi oT_h = 1$ a.e. in C_ϵ , i.e. $\chi = 1$ a.e. in $T_h(C_\epsilon)$.

From (1.4) and (1.5), we have $div(a(X)\nabla u) = -div(H(X)) \leq 0$ in $\mathcal{D}'(T_h(C_\epsilon))$. Then by the strong maximum principle we deduce, since $u \geq 0$ in Ω and $u > 0$ in $T_h(Q_\epsilon) \subset T_h(C_\epsilon)$, that $u > 0$ in $T_h(C_\epsilon)$ (see Figure 2). \square

Thanks to Proposition 3.1, we can define for each $h \in \pi_y(\Omega)$, the following function ϕ_h on $\pi_x(\Omega \cap [y = h])$ by

$$\phi_h(\omega) = \begin{cases} \sup\{t / (t, \omega) \in D_h, \quad uoT_h(t, \omega) > 0\} \\ \text{if this set is not empty} \\ \alpha_-(\omega) & \text{otherwise.} \end{cases} \quad (3.1)$$

Remark 3.1. *Since $u = \varphi > 0$ on Γ_3 and $u \in C^0(\Omega \cup \Gamma_3)$, we have $u > 0$ below Γ_3 in the following sense :*

$$u(X(t, \omega)) > 0 \quad \forall t \in [\alpha_-(\omega), \alpha_+(\omega)] \quad \text{such that} \quad X(\alpha_+(\omega), \omega) \in \Gamma_3.$$

Consequently, if $X(t_0, \omega_0) \in \Omega$ and $u(X(t_0, \omega_0)) = 0$, we have necessarily $X(\alpha_+(\omega_0), \omega_0) \in \bar{\Gamma}_1 \cup \bar{\Gamma}_2$.

Arguing as in [3], we have the following results

Proposition 3.2. *ϕ_h is lower semi-continuous on each $\omega \in \pi_x(\Omega \cap [y = h])$ such that $T_h(\phi_h(\omega), \omega) \in \Omega$. Moreover*

$$[uoT_h(t, \omega) > 0] \cap D_h = [t < \phi_h(\omega)].$$




Figure 2

The following important lemmas will be useful in Sections 4 and 5. Some of them are extensions of lemmas in [3].

Lemma 3.1. *Let $h \in \pi_y(\Omega)$, $\omega_1, \omega_2 \in \pi_x(\Omega \cap [y = h])$ with $\omega_1 < \omega_2$, and $\underline{y} \in \pi_y(\Omega)$. We denote by $t_{\underline{y}}(\omega)$ the unique t (if it exists) at which the orbit $\gamma(\omega)$ meets the line $[y = \underline{y}]$. Assume that for $i = 1, 2$, $\gamma(\omega_i) \cap [y = \underline{y}] \neq \emptyset$ and that $[X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_2), \omega_2)] \subset \subset \Omega$. Then we have*

- i) $\gamma(\omega) \cap [y = \underline{y}] \neq \emptyset \quad \forall \omega \in [\omega_1, \omega_2]$
- ii) $[X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_2), \omega_2)] = \{X(t_{\underline{y}}(\omega), \omega) / \omega \in [\omega_1, \omega_2]\}$.

Proof. i) First note that it is enough to prove the assertion for $\omega \in (\omega_1, \omega_2)$. Moreover if $\underline{y} = h$, then the assertion is trivial since in this case for all $\omega \in [\omega_1, \omega_2]$, $\gamma(\omega) \cap [y = \underline{y}] = \{(\omega, h)\}$. So we assume that $\underline{y} \neq h$ and discuss the two cases :

* $\underline{y} > h$: For each $\omega \in (\omega_1, \omega_2)$, the half orbit $\gamma^+(\omega) = \gamma(\omega) \cap [t \geq 0]$ is enclosed between

$\gamma^+(\omega_1)$ and $\gamma^+(\omega_2)$. So if $\gamma^+(\omega) \cap [y = \underline{y}] = \emptyset$, then $\gamma^+(\omega)$ will never reach $\partial\Omega$, which contradicts $X(\alpha_+(\omega), \omega) \in \partial\Omega \cap [y > h]$.

$*y < h$: For each $\omega \in (\omega_1, \omega_2)$, the half orbit $\gamma^-(\omega) = \gamma(\omega) \cap [t \leq 0]$ is enclosed between $\gamma^-(\omega_1)$ and $\gamma^-(\omega_2)$. So if $\gamma^-(\omega) \cap [y = \underline{y}] = \emptyset$, then $\gamma^-(\omega)$ will never reach $\partial\Omega$, which contradicts $X(\alpha_-(\omega), \omega) \in \partial\Omega \cap [y < h]$.

ii) First note that it is enough to show that

$$(X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_2), \omega_2)) = \{X(t_{\underline{y}}(\omega), \omega) / \omega \in (\omega_1, \omega_2)\},$$

where $(X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_2), \omega_2))$ denotes the line segment without the extreme points.

- Let $\omega \in (\omega_1, \omega_2)$. Since the orbit $\gamma(\omega)$ is strictly enclosed between the orbits $\gamma(\omega_1)$ and $\gamma(\omega_2)$, and meets the line $[y = \underline{y}]$ at the point $X(t_{\underline{y}}(\omega), \omega)$, we have $X_1(t_{\underline{y}}(\omega_1), \omega_1) < X_1(t_{\underline{y}}(\omega), \omega) < X_1(t_{\underline{y}}(\omega_2), \omega_2)$ and therefore $X(t_{\underline{y}}(\omega), \omega) \in (X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_2), \omega_2))$.

- Let $(x_*, \underline{y}) \in (X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_2), \omega_2))$. We consider $X(\cdot, x_*, \underline{y})$ the maximal solution of the differential equation $X'(t) = H(\bar{X}(t))$, $X(0) = (x_*, \underline{y})$.

Note that the orbit $\gamma(x_*, \underline{y})$ of $X(\cdot, x_*, \underline{y})$ has no intersection with $\gamma(\omega_i)$, $i = 1, 2$, because otherwise we will have $\gamma(x_*, \underline{y}) = \gamma(\omega_1)$ or $\gamma(x_*, \underline{y}) = \gamma(\omega_2)$, which is impossible since $x_* \in (X_1(t_{\underline{y}}(\omega_1), \omega_1), X_1(t_{\underline{y}}(\omega_2), \omega_2))$. Hence $\gamma(x_*, \underline{y})$ is strictly enclosed between $\gamma(\omega_1)$ and $\gamma(\omega_2)$. Therefore it meets the line $[y = h]$ at the point (ω_*, h) , with $\omega_* \in (\omega_1, \omega_2)$. It follows that $X(t, x_*, \underline{y}) = X(t + t_{\underline{y}}(\omega_*), \omega_*, h)$ and in particular $(x_*, \underline{y}) = X(0, x_*, \underline{y}) = X(t_{\underline{y}}(\omega_*), \omega_*, h)$. \square

Lemma 3.2. *Let (u, χ) be a solution of (P) . Let $h \in \pi_y(\Omega)$, $\omega_1, \omega_2 \in \pi_x(\Omega \cap [y = h])$ with $\omega_1 < \omega_2$. Let $\underline{y} \in \pi_y(\Omega)$ such that $[y = \underline{y}] \cap \gamma(\omega_i) \neq \emptyset$ $i = 1, 2$.*

Set $D_{\underline{y}} = T_h(\{(t, \omega) \in D_h, \omega \in (\omega_1, \omega_2), t > t_{\underline{y}}(\omega)\}) = T_h([\omega_1 < \omega < \omega_2]) \cap [y > \underline{y}]$, and assume that $\bar{D}_{\underline{y}} \cap \bar{\Gamma}_3 = \emptyset$ (see Figure 3). Then if $uoT_h(t_{\underline{y}}(\omega_i), \omega_i) = 0$ for $i = 1, 2$, we have

$$\int_{D_{\underline{y}}} (a(X)\nabla u + \chi H(X)) \cdot \nabla \zeta dX \leq 0$$

$$\forall \zeta \in H^1(D_{\underline{y}}), \quad \zeta \geq 0, \quad \zeta(x, \underline{y}) = 0 \quad \text{for a.e. } (x, \underline{y}) \in \bar{D}_{\underline{y}}.$$

Proof. First note that $D_{\underline{y}}$ is well defined since by Lemma 3.1 i), $t_{\underline{y}}(\omega)$ exists for each $\omega \in (\omega_1, \omega_2)$. Next we claim that

$$\int_{D_{\underline{y}}} (a(X)\nabla u + \chi([u > 0])H(X)) \cdot \nabla \zeta dX \leq \int_{\omega_1}^{\omega_2} -(Y_h \cdot \zeta \circ T_h)(\phi_h(\omega), \omega) d\omega \quad (3.2)$$

$$\forall \zeta \in H^1(D_{\underline{y}}) \cap C^0(\bar{D}_{\underline{y}}), \quad \zeta \geq 0, \quad \zeta(x, \underline{y}) = 0 \quad \text{for all } (x, \underline{y}) \in \bar{D}_{\underline{y}}.$$

Indeed, we deduce from $uoT_h(t_{\underline{y}}(\omega_i), \omega_i) = 0$, $i = 1, 2$ and Proposition 3.1 ii) that $uoT_h(t, \omega_i) = 0$, for all $t \geq t_{\underline{y}}(\omega_i)$, $i = 1, 2$. Therefore for $\epsilon > 0$, $\chi(D_{\underline{y}}) \cdot \min(\frac{u}{\epsilon}, \zeta)$ is a test function for (P) and we have




Figure 3

$$\begin{aligned}
& \int_{D_{\underline{y}} \cap [u \geq \epsilon \zeta]} a(X) \nabla u \cdot \nabla \zeta dX + \int_{D_{\underline{y}}} \chi([u > 0]) H(X) \cdot \nabla \zeta dX \\
& \leq \int_{D_{\underline{y}}} \chi([u > 0]) H(X) \cdot \nabla \left(\zeta - \frac{u}{\epsilon} \right)^+ dX = I_\epsilon.
\end{aligned}$$

Using the change of variables T_h and the second mean value theorem, we obtain

$$\begin{aligned}
I_\epsilon &= \int_{J = \{\omega \in (\omega_1, \omega_2) / \phi_h(\omega) > t_{\underline{y}}(\omega)\}} \int_{t_{\underline{y}}(\omega)}^{\phi_h(\omega)} \frac{\partial}{\partial t} \left(\left(\zeta - \frac{u}{\epsilon} \right)^+ \circ T_h \right) \cdot (-Y_h(t, \omega)) dt d\omega \\
&= \int_J (-Y_h(\phi_h(\omega), \omega)) \left\{ \int_{t^*(\omega)}^{\phi_h(\omega)} \frac{\partial}{\partial t} \left(\left(\zeta - \frac{u}{\epsilon} \right)^+ \circ T_h \right) (t, \omega) dt \right\} d\omega \\
&\leq \int_{\omega_1}^{\omega_2} -Y_h(\phi_h(\omega), \omega) \cdot \zeta \circ T_h(\phi_h(\omega), \omega) d\omega, \quad t^*(\omega) \in [t_{\underline{y}}(\omega), \phi_h(\omega)].
\end{aligned}$$

Then by letting ϵ go to 0, the inequality (3.2) holds.

Now to prove the lemma, it suffices to do it for $\zeta \in H^1(D_{\underline{y}}) \cap C^0(\overline{D_{\underline{y}}})$, $\zeta \geq 0$, $\zeta(x, \underline{y}) = 0$ for all $(x, \underline{y}) \in \overline{D_{\underline{y}}}$ and conclude by density. So let $\epsilon > 0$ and $h_\epsilon = \theta_\epsilon \circ T_h^{-1}$, with $\theta_\epsilon(\omega) = \min\left(\frac{(\omega - \omega_1)^+}{\epsilon}, 1\right) \cdot \min\left(\frac{(\omega_2 - \omega)^+}{\epsilon}, 1\right)$. Since $\chi(D_{\underline{y}}) \cdot \zeta \cdot h_\epsilon$ is a test function for (P), we have

$$\begin{aligned} & \int_{D_{\underline{y}}} (a(X)\nabla u + \chi H(X)) \cdot \nabla \zeta dX \leq \int_{D_{\underline{y}}} (a(X)\nabla u + \chi H(X)) \cdot \nabla((1 - h_\epsilon)\zeta) dX \\ & = \int_{D_{\underline{y}}} (a(X)\nabla u + \chi([u > 0])H(X)) \cdot \nabla((1 - h_\epsilon)\zeta) dX \\ & \quad + \int_{D_{\underline{y}}} (\chi - \chi([u > 0]))H(X) \cdot \nabla((1 - h_\epsilon)\zeta) dX = I_\epsilon^1 + I_\epsilon^2. \end{aligned}$$

Using (3.2) and the fact that $\theta_\epsilon \xrightarrow{\epsilon \rightarrow 0} 1$, we obtain the lemma since we have

$$\begin{aligned} I_\epsilon^1 & \leq \int_{\omega_1}^{\omega_2} -Y_h(\phi_h(\omega), \omega) \cdot \zeta \circ T_h(\phi_h(\omega), \omega) \cdot (1 - \theta_\epsilon(\omega)) d\omega, \\ I_\epsilon^2 & = \int_{T_h^{-1}(D_{\underline{y}})} (\chi \circ T_h - \chi([u \circ T_h > 0]))(-Y_h(t, \omega)) \cdot \frac{\partial}{\partial t}(\zeta \circ T_h) \cdot (1 - \theta_\epsilon(\omega)) dt d\omega. \end{aligned}$$

□

Lemma 3.3. *Let (u, χ) be a solution of (P) and $X_0 = (x_0, y_0) = T_h(t_0, \omega_0)$ be a point of $T_h(D_h)$. We denote by $B_r(t_0, \omega_0)$ a ball with center (t_0, ω_0) and radius r contained in D_h . If $u \circ T_h = 0$ in $B_r(t_0, \omega_0)$, then*

$$u \circ T_h = 0 \quad \text{in } C_r \quad \text{and} \quad \chi \circ T_h = 0 \quad \text{a.e. in } C_r$$

where $C_r = \{(t, \omega) \in D_h, |\omega - \omega_0| < r, t > t_0\} \cup B_r(t_0, \omega_0)$. In other words if $u = 0$ in $T_h(B_r(t_0, \omega_0))$, then $u = 0$ and $\chi = 0$ a.e. in $T_h(C_r)$ (see Figure 4).

Proof. By Proposition 3.1, we have $u \circ T_h = 0$ in C_r . Applying Lemma 3.2 with domains $D_{\underline{y}} = T_h([\omega_1 < \omega < \omega_2]) \cap [y > \underline{y}] \subset T_h(C_r)$, ($\underline{y} \in \pi_y(\Omega)$) satisfying $[y = \underline{y}] \cap \gamma(\omega) \neq \emptyset \forall \omega \in [\omega_1, \omega_2]$ and taking $\zeta = (y - \underline{y})\chi(D_{\underline{y}})$, we obtain

$$\int_{D_{\underline{y}}} \chi H_2(X) dX \leq 0.$$

From (1.3), we deduce that $\chi = 0$ a.e in $D_{\underline{y}}$. This holds for all domains $D_{\underline{y}}$ in $T_h(C_r)$. Hence $\chi = 0$ a.e in $T_h(C_r)$. □

Lemma 3.4. *Let (u, χ) be a solution of (P), $X_0 = (x_0, y_0) = T_h(t_0, \omega_0)$ be a point of Ω and B_r the open ball in D_h with center (t_0, ω_0) and radius r . Then we cannot have the following situations (see Figure 5)*




Figure 4

$$\begin{aligned}
 (i) \quad & \begin{cases} uoT_h(t, \omega_0) = 0 & \forall t \in (t_0 - r, t_0 + r) \\ uoT_h(t, \omega) > 0 & \forall (t, \omega) \in B_r \setminus S, \quad S = (t_0 - r, t_0 + r) \times \{\omega_0\}, \end{cases} \\
 (ii) \quad & \begin{cases} uoT_h(t, \omega) = 0 & \forall (t, \omega) \in B_r \cap [\omega \leq \omega_0] \\ uoT_h(t, \omega) > 0 & \forall (t, \omega) \in B_r \cap [\omega > \omega_0], \end{cases} \\
 (iii) \quad & \begin{cases} uoT_h(t, \omega) = 0 & \forall (t, \omega) \in B_r \cap [\omega \geq \omega_0] \\ uoT_h(t, \omega) > 0 & \forall (t, \omega) \in B_r \cap [\omega < \omega_0]. \end{cases}
 \end{aligned}$$

Proof. Assume ii) holds. The proof of i) and iii) is based on the same arguments. Let $\zeta \in \mathcal{D}(T_h(B_r))$, $\zeta \geq 0$. Using the fact that, by Lemma 3.2, $\chi oT_h = 0$ a.e on $B_r \cap [\omega < \omega_0]$ and $\pm\zeta$ are tests functions for (P) , we obtain after using the change of variables T_h

$$\int_{T_h(B_r)} a(X) \nabla u \cdot \nabla \zeta dX = \int_{B_r \cap [\omega > \omega_0]} \frac{\partial}{\partial t} (-Y_h(t, \omega)) \zeta oT_h dt d\omega \geq 0.$$




Figure 5 (i)

We deduce that $\operatorname{div}(a(X)\nabla u) \leq 0$ in $\mathcal{D}'(T_h(B_r))$. By the strong maximum principle, we have either $u > 0$ or $u = 0$ in $T_h(B_r)$, which contradicts the assumption. \square

Lemma 3.5. *Let (u, χ) be a solution of (P), $X_0 = (x_0, y_0) = T_h(t_0, \omega_0)$ be a point of Ω such that $u \circ T_h(t_0, \omega_0) = 0$. Then there exists $\rho > 0$ such that one of the following situations holds :*

- $$\begin{aligned}
 (i) \quad & \left\{ \begin{array}{l} u \circ T_h(t, \omega) > 0 \quad \forall (t, \omega) \in B_\rho(t_0, \omega_0) \cap [\omega < \omega_0], \\ \text{there exists a sequence } (t_n, \omega_n)_{n \geq 1} \subset B_\rho(t_0, \omega_0) \cap [\omega > \omega_0] \\ \text{such that } \forall n \geq 1 \quad u \circ T_h(t_n, \omega_n) = 0 \quad \text{and} \quad X(t_n, \omega_n)_{n \rightarrow \infty} X_0 \end{array} \right. \\
 (ii) \quad & \left\{ \begin{array}{l} u \circ T_h(t, \omega) > 0 \quad \forall (t, \omega) \in B_\rho(t_0, \omega_0) \cap [\omega > \omega_0], \\ \text{there exists a sequence } (t_n, \omega_n)_{n \geq 1} \subset B_\rho(t_0, \omega_0) \cap [\omega < \omega_0] \\ \text{such that } \forall n \geq 1 \quad u \circ T_h(t_n, \omega_n) = 0 \quad \text{and} \quad X(t_n, \omega_n)_{n \rightarrow \infty} X_0 \end{array} \right.
 \end{aligned}$$




Figure 5 (ii)

$$(iii) \left\{ \begin{array}{l} \text{There exists two sequences } (t_n^\pm, \omega_n^\pm)_{n \geq 1} \subset B_\rho(t_0, \omega_0), \\ \text{such that } \forall n \geq 1 \quad \omega_n^- < \omega_0 < \omega_n^+ \quad uoT_h(t_n^-, \omega_n^-) = uoT_h(t_n^+, \omega_n^+) = 0 \\ \text{and } X(t_n^-, \omega_n^-)_{n \rightarrow \infty} X_0, \quad X(t_n^+, \omega_n^+)_{n \rightarrow \infty} X_0. \end{array} \right.$$

Proof. Let $\eta > 0$ such that $B_\eta(t_0, \omega_0) \subset D_h$. By Proposition 3.1, we have $uoT_h(t, \omega_0) = 0 \forall t \geq t_0$. Then for any $\rho \in (0, \eta)$, by Lemma 3.4, one of the following situations holds necessarily

$$\begin{array}{ll} \alpha) & \exists (t_1^-, \omega_1^-) \in B_\rho(t_0, \omega_0) \cap [\omega < \omega_0] \quad \text{such that} \quad uoT_h(t_1^-, \omega_1^-) = 0 \\ \beta) & \exists (t_1^+, \omega_1^+) \in B_\rho(t_0, \omega_0) \cap [\omega > \omega_0] \quad \text{such that} \quad uoT_h(t_1^+, \omega_1^+) = 0. \end{array}$$

We discuss the following cases

- If α) and β) holds simultaneously for any $\rho \in (0, \eta)$, then we are in the situation *iii*).
- If for example α) does not hold for some $\rho \in (0, \eta)$. Then $uoT_h > 0$ in $B_\rho(t_0, \omega_0) \cap [\omega < \omega_0]$. Moreover by Lemma 3.4, β) holds for any $\rho' \in (0, \rho)$. In this case we are in the situation *i*).




Figure 5 (iii)

• If for example β does not hold, then we show as in the previous case that we obtain the situation *ii*). \square

4 A Comparison Result

In all what follows, we assume that

$$a \in C_{loc}^{0,\alpha}(\Omega) \quad (0 < \alpha < 1) \quad (4.1)$$

$$\exists c_0 \in \mathbb{R} \quad / \quad \forall Y \in \Omega \quad : \quad \operatorname{div}(a(X)(X - Y)) \leq c_0 \quad \text{in } \mathcal{D}'(\Omega). \quad (4.2)$$

Note that (4.2) is satisfied in particular if $a \in C^{0,1}$ or simply if $\operatorname{div}(a(X)e_1), \operatorname{div}(a(X)e_2) \in L^\infty(\Omega)$, where $e_1 = (1, 0)$ and $e_2 = (0, 1)$. Moreover, one can adapt the proof in [4] (see Remark 2.2 of this reference) to verify that $u \in C_{loc}^{0,1}(\Omega)$. The main result of this section is the comparison Lemma 4.4. First, we construct a barrier function and establish some of its properties.

Lemma 4.1. *Let $k > 0$, $(x_1, \underline{y}), (x_2, \underline{y}) \in \Omega$ with $x_1 < x_2$ and $x_2 - x_1 = 2k\epsilon$, where ϵ is small enough so that*

$$(x_1 - \epsilon, x_2 + \epsilon) \times (\underline{y}, \underline{y} + 2\epsilon) \subset\subset \Omega.$$

Let $Z = (x_1 - \epsilon, x_2 + \epsilon) \times (\underline{y}, \underline{y} + \epsilon)$ and denote by v the unique solution in $H^1(Z)$ of

$$\begin{cases} \operatorname{div}(a(X)\nabla v) = -\operatorname{div}(H(X)) & \text{in } Z \\ v = \epsilon(\underline{y} + \epsilon - \underline{y})^+ & \text{on } \partial Z. \end{cases} \quad (4.3)$$

Then, there exists a positive constant C independent of ϵ such that

$$\begin{aligned} i) & \quad 0 < v \leq C\epsilon^2 \quad \text{in } Z \\ ii) & \quad |\nabla v(X)| \leq C\epsilon \quad \forall X \in T = [x_1, x_2] \times \{\underline{y} + \epsilon\}. \end{aligned}$$

Proof. *i)* Since $\operatorname{div}(a(X)\nabla v) = -\operatorname{div}(H(X)) \leq 0$ in Z and due to the boundary condition, we deduce by the weak and strong maximum principles (see [5]) that $v > 0$ in Z .

To prove the second inequality, we introduce the function

$$\begin{aligned} \omega & : \widehat{Z} = (0, 2k + 2) \times (0, 1) \longrightarrow \mathbb{R}^+ \\ X' & = (x', y') \longmapsto \omega(X') = v(x_1 - \epsilon + \epsilon x', \underline{y} + \epsilon y'). \end{aligned}$$

It is not difficult to check that

$$\begin{cases} \operatorname{div}(\widehat{a}(X')\nabla\omega) = -\epsilon^2 \widehat{\operatorname{div}H} & \text{in } \widehat{Z} \\ \omega = \epsilon^2(1 - y')^+ & \text{on } \partial\widehat{Z} \end{cases} \quad (4.4)$$

where

$$\widehat{a}(X') = a(x_1 - \epsilon + \epsilon x', \underline{y} + \epsilon y'), \quad \widehat{\operatorname{div}H}(X') = (\operatorname{div}H)(x_1 - \epsilon + \epsilon x', \underline{y} + \epsilon y').$$

Moreover we have

$$\begin{aligned} \widehat{a}(X')\xi \cdot \xi & \geq \lambda|\xi|^2, \quad \forall \xi \in \mathbb{R}^2, \quad \forall X' \in \widehat{Z} \\ |\widehat{a}|_{\infty, \widehat{Z}} & \leq M, \quad 0 \leq \widehat{\operatorname{div}H}(X') \leq C = |\operatorname{div}H|_{\infty}, \quad \forall X' \in \widehat{Z}. \end{aligned}$$

Applying Theorem 8.16 p. 191 of [5], we get

$$\sup_{\widehat{Z}} \omega \leq \sup_{\partial\widehat{Z}} \omega + C_1 \frac{|\epsilon^2 \widehat{\operatorname{div}H}|_{L^{q/2}}}{\lambda}$$

where $q > 2$ and C_1 is a positive constant depending only on Y . So

$$\sup_Z v = \sup_{\widehat{Z}} \omega \leq \epsilon^2 + C_2 \epsilon^2 = C\epsilon^2.$$

ii) Let $S = (\frac{1}{2}, 2k + \frac{3}{2}) \times \{1\}$ and $\widehat{Z}' = (\frac{1}{2}, 2k + \frac{3}{2}) \times (\frac{1}{2}, 1)$. Since S is a $C^{1,\alpha}$ boundary portion of $\partial\widehat{Z}$, $\omega = 0$ on S , we deduce from (4.4) by applying Corollary 8.36 p. 212 [5] that $\omega \in C^{1,\alpha}(\widehat{Z} \cup S)$ with the following estimate

$$|\omega|_{1,\alpha,\widehat{Z}'} \leq C \left(|\omega|_{0,\widehat{Z}} + |\epsilon^2 \widehat{\operatorname{div}H}|_{0,\widehat{Z}} \right)$$

where $C = C(\lambda, M, K, d', S)$ is a constant independent of ϵ , $d' = d(\widehat{Z}', \partial\widehat{Z}\setminus S)$ and $K = \max_{i,j}(|a_{ij}|_{0,\alpha})$.

Taking into account the estimate in i), we obtain

$$|\nabla\omega|_{0,\widehat{Z}'} \leq |\omega|_{1,\alpha,\widehat{Z}'} \leq C\epsilon^2$$

which, in particular, leads to

$$|\nabla\omega(x', 1)| \leq C\epsilon^2 \quad \forall x' \in [1, 1 + 2k].$$

Therefore

$$|\nabla v(x, \underline{y} + \epsilon)| = \frac{1}{\epsilon} \left| \nabla\omega\left(\frac{x - x_1 + \epsilon}{\epsilon}, 1\right) \right| \leq C\epsilon \quad \forall x \in [x_1, x_2].$$

□

Lemma 4.2. *Let $h \in \pi_y(\Omega)$, $\omega_1, \omega_2 \in \pi_x(\Omega \cap [y = h])$ with $\omega_1 < \omega_2$.*

Let $y \in \pi_y(\Omega)$ such that $\gamma(\omega_i) \cap [y = y] \neq \emptyset$ $i = 1, 2$.

Set $\overline{D}_y = T_h([\omega_1 < \omega < \omega_2]) \cap [y > y]$. Assume that $D_y \cap [y < \underline{y} + \epsilon] \subset (x_1, x_2) \times (\underline{y}, \underline{y} + \epsilon) \subset Z$ with \overline{Z} defined in Lemma 4.1. Then after extending v by 0 to \overline{D}_y , we obtain

$$\int_{D_y} (a(X)\nabla v + \chi([v > 0])H(X))\nabla\zeta dX \geq 0 \quad \forall \zeta \in H^1(D_y), \zeta \geq 0, \zeta = 0 \text{ on } \partial D_y \cap \Omega.$$

Proof. Set $T' = [y = \underline{y} + \epsilon] \cap \overline{D}_y \subset T$ and let ν be the outward unit normal vector to T . We have by Lemma 4.1 *ii*) $a(X)\nabla v \cdot \nu + H(X) \cdot \nu = a(X)\nabla v \cdot e_y + H_2(X) \geq -C\epsilon + \underline{h} \geq 0$ on T' for ϵ small enough. Now, for $\zeta \in H^1(D_y)$, $\zeta \geq 0$, $\zeta = 0$ on $\partial D_y \cap \Omega$, we have

$$\int_{D_y} (a(X)\nabla v + \chi([v > 0])H(X))\nabla\zeta dX = \int_{T'} (a(X)\nabla v \cdot \nu + H(X) \cdot \nu)\zeta dX \geq 0.$$

□

The following lemma extends a lemma proved in [4] for $H(X) = h(X)e_2$.

Lemma 4.3. *Let (u, χ) be a solution of (P). Assume that the hypothesis of Lemma 4.2 hold, $\overline{D}_y \cap \Gamma_3 = \emptyset$ and (see Figure 6)*

$$\begin{aligned} uoT_h(t_y(\omega_1), \omega_1) &= uoT_h(t_y(\omega_2), \omega_2) = 0 \\ uoT_h(t_y(\omega), \omega) &\leq \epsilon^2 = v(t_y(\omega), \omega) \quad \forall \omega \in (\omega_1, \omega_2), \end{aligned}$$

then we have

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta} \int_{D_y \cap [v > 0] \cap [0 < u - v < \delta]} a(X)\nabla(u - v)^+ \cdot \nabla(u - v)^+ dX = 0.$$




Figure 6

Proof. For $\delta, \eta > 0$, let $F_\delta(s)$ be the function introduced in the proof of Proposition 1.1, $d_\eta(y) = F_\eta(y - \bar{y})$ and $\bar{y} = y + \epsilon$. By applying Lemma 3.2 and Lemma 4.2 for $\zeta = F_\delta(u - v) + d_\eta(1 - H_\delta(u))$ and for $\zeta = F_\delta(u - v)$ respectively, we get

$$\begin{aligned} & \int_{D_{\underline{y}}} (a(X)\nabla u + \chi H(X)) \cdot \nabla (F_\delta(u - v)) dX \\ & \leq - \int_{D_{\underline{y}}} (a(X)\nabla u + \chi H(X)) \cdot \nabla (d_\eta(1 - F_\delta(u))) dX. \end{aligned} \quad (4.5)$$

$$\int_{D_{\underline{y}}} (a(X)\nabla v + \chi([v > 0])H(X)) \cdot \nabla (F_\delta(u - v)) dX \geq 0. \quad (4.6)$$

Using (4.5) and (4.6), we get since $d_\eta = 0$ on $[v > 0]$

$$\int_{D_{\underline{y}} \cap [v > 0]} F'_\delta(u - v) a(X) \nabla(u - v) \cdot \nabla(u - v) dX$$

$$\begin{aligned}
&\leq - \int_{D_{\underline{y}} \cap [v=0]} (1 - d_\eta)(a(X)\nabla u + \chi H(X)) \cdot \nabla(F_\delta(u)) dX \\
&\quad - \int_{D_{\underline{y}} \cap [v=0]} (1 - F_\delta(u))(a(X)\nabla u + \chi H(X)) \cdot \nabla d_\eta dX = I_1^{\delta\eta} + I_2^{\delta\eta}.
\end{aligned}$$

Since

$$|I_1^{\delta\eta}| \leq \int_{D_{\underline{y}} \cap [\bar{y} < y < \bar{y} + \eta]} |(a(X)\nabla u + \chi H(X)) \cdot \nabla(F_\delta(u))| dX,$$

we obtain $\lim_{\eta \rightarrow 0} I_1^{\delta\eta} = 0$.

As for $I_2^{\delta\eta}$, we have

$$\begin{aligned}
I_2^{\delta\eta} &= - \int_{D_{\underline{y}} \cap [u=v=0]} \chi H(X) \cdot \nabla d_\eta dX \\
&\quad - \int_{D_{\underline{y}} \cap [u > 0=v]} (1 - F_\delta(u))(a(X)\nabla u + H(X)) \cdot \nabla d_\eta dX = I_3^{\delta\eta} + I_4^{\delta\eta} \leq I_4^{\delta\eta}.
\end{aligned}$$

since

$$I_3^{\delta\eta} = - \int_{D_{\underline{y}} \cap [u=v=0]} H_2(X) \cdot \chi \cdot \partial_y d_\eta dX = \frac{-1}{\eta} \int_{D_{\underline{y}} \cap [u=v=0] \cap [\bar{y} < y < \bar{y} + \eta]} H_2(X) \chi dX \leq 0.$$

Moreover since $u \in C_{loc}^{0,1}(\Omega)$, one has for some constant C

$$\begin{aligned}
|I_4^{\delta\eta}| &\leq \frac{C}{\eta} \int_{D_{\underline{y}} \cap [u > 0=v] \cap [\bar{y} < y < \bar{y} + \eta]} (1 - F_\delta(u)) dX \\
&= \frac{C}{\eta} \int_J \int_{t_{\bar{y}}(\omega)}^{\min(\phi_h(\omega), t_{\bar{y}+\eta}(\omega))} (1 - F_\delta(uoT_h))(t, \omega) \cdot (-Y_h(t, \omega)) dt d\omega \\
&\leq C \int_J \left(\frac{1}{\eta} \int_{t_{\bar{y}}(\omega)}^{t_{\bar{y}}(\omega) + \frac{\eta}{h}} (1 - F_\delta(uoT_h)) dt \right) d\omega,
\end{aligned}$$

where $J = \{\omega \in (\omega_1, \omega_2) / \phi_h(\omega) > t_{\bar{y}}(\omega)\}$.

Since the function $t \mapsto 1 - F_\delta(uoT_h(t, \omega))$ is continuous, we obtain

$$\limsup_{\eta \rightarrow 0} |I_4^{\delta\eta}| \leq C \int_J (1 - F_\delta(uoT_h(t_{\bar{y}}(\omega), \omega))) d\omega.$$

Hence

$$\int_{D_{\underline{y}} \cap [v > 0] \cap [0 < u - v < \delta]} \frac{1}{\delta} a(X) \nabla(u - v)^+ \cdot \nabla(u - v)^+ dX \leq C \int_J (1 - F_\delta(uoT_h(t_{\bar{y}}(\omega), \omega))) d\omega.$$

But given that $\omega \in J$, we have $uoT_h(t_{\bar{y}}(\omega), \omega) > 0$. Thus $\lim_{\delta \rightarrow 0} (1 - F_\delta(uoT_h(t_{\bar{y}}(\omega), \omega))) = 0$ and the result follows. \square

Lemma 4.4. *Let (u, χ) be a solution of (P). Assume that the hypothesis of Lemma 4.3 hold. Then we have*

$$u \equiv 0 \quad \text{in } D_{\underline{y}} \cap [y > \underline{y} + \epsilon].$$

Proof. Let

$$\begin{aligned} D^+ &= D_{\underline{y}} \cap [v > 0] = D_{\underline{y}} \cap [\underline{y} < y < \underline{y} + \epsilon] \\ \Delta &= T_h(\{(t, \omega) \in D_h / \omega \in (\omega_1, \omega_2), \alpha_-(\omega) < t < t_{\underline{y}+\epsilon}(\omega)\}) \\ w &= \begin{cases} (u - v)^+ & \text{in } D^+ \\ 0 & \text{in } \Delta \setminus \overline{D^+}. \end{cases} \end{aligned}$$

We have $w \in H^1(\Delta)$ since by assumption $u \leq v$ on $\Delta \cap [y = \underline{y}]$. Let $\zeta \in \mathcal{D}(\Delta)$. We have

$$\begin{aligned} \int_{\Delta} a(X) \nabla w \cdot \nabla \zeta dX &= \int_{D^+} a(X) \nabla (u - v)^+ \cdot \nabla \zeta dX \\ &= \lim_{\delta \rightarrow 0} \int_{D^+} F_{\delta}(u - v) a(X) \nabla (u - v)^+ \cdot \nabla \zeta dX = \lim_{\delta \rightarrow 0} I_{\delta}. \end{aligned}$$

Note that

$$\begin{aligned} I_{\delta} &= \int_{D^+} a(X) \nabla (u - v)^+ \cdot \nabla (F_{\delta}(u - v) \zeta) dX \\ &\quad - \frac{1}{\delta} \int_{D^+ \cap [0 < u - v < \delta]} \zeta a(X) \nabla (u - v) \cdot \nabla (u - v) dX = I_{\delta}^1 - I_{\delta}^2. \end{aligned}$$

By Lemma 4.3, $\lim_{\delta \rightarrow 0} I_{\delta}^2 = 0$, since we have

$$|I_{\delta}^2| \leq \sup_{\Delta} |\zeta| \cdot \frac{1}{\delta} \int_{D^+ \cap [0 < u - v < \delta]} a(X) \nabla (u - v) \cdot \nabla (u - v) dX.$$

Moreover, we have since $(F_{\delta}(u - v) \zeta) \in H_0^1(D^+)$,

$$\begin{aligned} I_{\delta}^1 &= \int_{D^+} a(X) \nabla u \cdot \nabla (F_{\delta}(u - v) \zeta) dX - \int_{D^+} a(X) \nabla v \cdot \nabla (F_{\delta}(u - v) \zeta) dX \\ &= - \int_{D^+} \chi H(X) \cdot \nabla (F_{\delta}(u - v) \zeta) dX + \int_{D^+} H(X) \cdot \nabla (F_{\delta}(u - v) \zeta) dX \\ &= 0 \quad \text{since } \chi = 1 \quad \text{a.e. in } [u > 0]. \end{aligned}$$

It follows that

$$\int_{\Delta} a(X) \nabla w \cdot \nabla \zeta dX = 0 \quad \forall \zeta \in \mathcal{D}(\Delta).$$

Since $w = 0$ in $\Delta \setminus \overline{D^+}$, we obtain by the strong maximum principle : $w = 0$ in Δ . Consequently, $u \leq v$ in D^+ and then $u \circ T_h(t_{\underline{y}+\epsilon}(\omega), \omega) = 0 \quad \forall \omega \in [\omega_1, \omega_2]$. Therefore

$$u \circ T_h(t, \omega) = 0 \quad \forall t \geq t_{\underline{y}+\epsilon}(\omega) \quad \forall \omega \in [\omega_1, \omega_2].$$

□

Combining Lemma 3.5 and Lemma 4.4, we obtain the following useful lemma

Lemma 4.5. *Let $X_0 = T_h(t_0, \omega_0) = (x_0, y_0) \in \Omega$, $\omega_{01}, \omega_{02} \in \pi_y(\Omega \cap [y = h])$ such that $u(X_0) = 0$, $\omega_{01} < \omega_0 < \omega_{02}$ and $\gamma(\omega_{0i}) \cap [y = y_0] \neq \emptyset$, $i = 1, 2$. Let $\epsilon > 0$ and $D_{y_0} = T_h([\omega_{01} < \omega < \omega_{02}]) \cap [y > y_0]$. We assume that for some $k > 0$, $D_{y_0} \cap [y < y_0 + \epsilon] \subset (x_0 - 2k\epsilon, x_0 + 2k\epsilon) \times (y_0, y_0 + 2\epsilon) \subset \subset \Omega$, and for all $\omega \in (\omega_{01}, \omega_{02})$ $uoT_h(t_{y_0}(\omega), \omega) \leq \epsilon^2$. Then the following situations cannot hold :*

- (i) $\left\{ \begin{array}{l} \text{There exists a sequence } (t_n, \omega_n)_{n \geq 1} \subset B_{\rho_0}(t_0, \omega_0) \cap [\omega < \omega_0] \text{ satisfying} \\ uoT_h(t_n, \omega_n) = 0 \quad \forall n \geq 1, \quad X(t_n, \omega_n) \xrightarrow{n \rightarrow \infty} X_0, \\ \forall n \geq 1, \quad X(\alpha_+(\omega_n), \omega_n) \text{ does not belong to the connected component of } \Gamma_1 \cup \Gamma_2 \\ \text{which contains } X(\alpha_+(\omega_0), \omega_0) \end{array} \right.$
- (ii) $\left\{ \begin{array}{l} \text{There exists a sequence } (t_n, \omega_n)_{n \geq 1} \subset B_{\rho_0}(t_0, \omega_0) \cap [\omega > \omega_0] \text{ satisfying} \\ uoT_h(t_n, \omega_n) = 0 \quad \forall n \geq 1, \quad X(t_n, \omega_n) \xrightarrow{n \rightarrow \infty} X_0 \\ \forall n \geq 1, \quad X(\alpha_+(\omega_n), \omega_n) \text{ does not belong to the connected component of } \Gamma_1 \cup \Gamma_2 \\ \text{which contains } X(\alpha_+(\omega_0), \omega_0). \end{array} \right.$

Proof. We will consider only the first situation. The second one can be treated similarly.

Let $(x^*, y^*) \in M$ such that $y^* > y_0 + \epsilon$, where M is the domain enclosed between $\gamma(\omega_{01})$, $\gamma(\omega_0)$, $[y = y_0]$ and $\partial\Omega$. Consider the maximal solution $X(\cdot, x^*, y^*)$ of $X'(t) = H(X(t))$, $X(0) = (x^*, y^*)$. The orbit $\gamma(x^*, y^*)$ of $X(\cdot, x^*, y^*)$ leaves M from the top at a point of $\partial\Omega$ and from the bottom at a point (x_*, y_0) of $[y = y_0]$.

From Lemma 3.1 ii), we know that $(x_*, y_0) = X(t_{y_0}(\omega_*), \omega_*, h)$ for some $\omega_* \in (\omega_{01}, \omega_0)$. It follows that the two orbits $\gamma(x^*, y^*)$ and $\gamma(\omega_*, h)$ coincide. Therefore we have $X(t, x^*, y^*) = X(t + t^*, \omega_*, h)$, where $t^* = t_{y^*}(\omega_*)$ is defined by $(x^*, y^*) = X(t^*, \omega_*, h)$.

We have $X_1(t_{y^*}(\omega_{01}), \omega_{01}, h) < x^* < X_1(t_{y^*}(\omega_0), \omega_0, h)$ and $X_{1n}(t_{y^*}(\omega_0), \omega_n, h)$ converges to $X_1(t_{y^*}(\omega_0), \omega_0, h)$ when $n \rightarrow \infty$. So there exists $n_1 > 1$ such that $x^* < X_{1n_1}(t_{y^*}(\omega_0), \omega_{n_1}, h)$.

We deduce that $(x^*, y^*) \in M_{n_1}$: the domain enclosed between $\gamma(\omega_{01})$, $\gamma(\omega_{n_1})$, $[y = y_0]$ and $\partial\Omega$. It follows, by Lemma 4.4, that $u \equiv 0$ in $M_{n_1} \cap [y \geq y_0 + \epsilon]$. In particular, we obtain $u(x^*, y^*) = 0$. This holds for any point of M . Then $u \equiv 0$ in $M \cap [y \geq y_0 + \epsilon]$. But (see Remark 3.1), this contradicts $\bar{M} \cap \Gamma_3 \neq \emptyset$ and $u > 0$ on Γ_3 . □

Remark 4.1. *Lemma 4.5 becomes trivial if α_+ is continuous. However we know only that α_+ is lower semi-continuous (see Lemma 10.5 p. 125, [1]). Of course one can have more regularity for α_+ if one assumes more regularity on H and the boundary of Ω . Actually one can verify that α_+ is C^1 if $H \in C^1(\bar{\Omega})$, $\partial\Omega$ is C^1 and $H(X) \cdot \nu$ does not vanish on $\partial\Omega$ (see Proposition 2.1, [3]).*

5 Continuity of the Free Boundary

The main result of this section is the continuity of the functions ϕ_h representing the free boundary. Note that by Remark 3.1, if $X(\phi_h(\omega), \omega) \in \Omega$, then $X(\alpha_+(\omega), \omega) \in \partial\Omega \setminus \Gamma_3$. Here we will consider the case where $X(\alpha_+(\omega), \omega) \in \partial\Omega \setminus \bar{\Gamma}_3$.

Theorem 5.1. *For each $h \in \pi_y(\Omega)$, the function ϕ_h is continuous at each $\omega \in \pi_x(\Omega \cap [y = h])$ such that $X(\phi_h(\omega), \omega) \in \Omega$ and $X(\alpha_+(\omega), \omega) \in \partial\Omega \setminus \bar{\Gamma}_3$.*

Proof. Let $\omega_0 \in \pi_x(\Omega \cap [y = h])$ such that $X(\phi_h(\omega_0), \omega_0) = T_h(\phi_h(\omega_0), \omega_0) = T_h(t_0, \omega_0) = (x_0, y_0) = X_0 \in \Omega$ and $X(\alpha_+(\omega_0), \omega_0) \in \partial\Omega \setminus \bar{\Gamma}_3$.

Let $0 < \epsilon < \min\left(\frac{h}{3}(\alpha_+(\omega_0) - t_0), \frac{h}{2}(t_0 - \alpha_-(\omega_0))\right)$.

Since $u(X_0) = 0$ and u continuous, there exists $\rho^* \in (0, \epsilon)$ such that

$$u(X) \leq \epsilon^2 \quad \forall X \in B_{\rho^*}(X_0) \subset T_h(D_h). \quad (5.1)$$

Since (t_0, ω_0) belongs to the open set $T_h^{-1}(B_{\rho^*}(X_0))$, there exists $\eta_1 \in (0, \rho^*)$ such that

$$B_{\eta_1}(t_0, \omega_0) \subset\subset T_h^{-1}(B_{q\rho^*}(X_0)) \quad \text{with } q = \underline{h}/4\bar{h}. \quad (5.2)$$

By Theorem 3.4 p 24 [6], $\exists \eta_2 \in (0, \eta_1)$ such that

$$\begin{aligned} X(t, \omega) & \text{ exists for all } (t, \omega) \in [\alpha_-(\omega_0), \alpha_+(\omega_0)] \times (\omega_0 - \eta_2, \omega_0 + \eta_2) \\ \text{and } (t, \omega) & \longmapsto X(t, \omega) \text{ is continuous.} \end{aligned} \quad (5.3)$$

So there exists $\eta_3 \in (0, \eta_2)$ such that

$$|X(t, \omega) - X(t_0, \omega_0)| < \epsilon \quad \forall (t, \omega) \in B_{\eta_3}(t_0, \omega_0). \quad (5.4)$$

Set $\rho = \eta_3 < \epsilon$. By Lemma 3.4, one of the following situations is true :

- i) $\exists (t_1, \omega_1) \in B_\rho(t_0, \omega_0)$ such that $\omega_1 < \omega_0$ and $u \circ T_h(t_1, \omega_1) = 0$
- ii) $\exists (t_2, \omega_2) \in B_\rho(t_0, \omega_0)$ such that $\omega_2 > \omega_0$ and $u \circ T_h(t_2, \omega_2) = 0$.

We will consider only the case where i) holds (see Figure 7). The other case can obviously be treated in a similar way. Note that $X(t_1, \omega_1)$ is at the left hand side of the orbit $\gamma(\omega_0)$ since $X(0, \omega_1) = (\omega_1, h)$, $X(0, \omega_0) = (\omega_0, h)$ and $\omega_1 < \omega_0$.

Set $\underline{y} = \max(X_2(t_0, \omega_0), X_2(t_1, \omega_1))$. Then

$$u \circ T_h(t_{\underline{y}}(\omega_i), \omega_i) = 0 \quad i = 0, 1. \quad (5.5)$$




Figure 7

Consider the set

$$\mathcal{O} = \{(x, y) = X(t, \omega) \in T_h(D_h) / |\omega - \omega_0| < \rho\} \cap [\underline{y} < y < \underline{y} + \epsilon].$$

Then we have

Lemma 5.1. *For all ω in $(\omega_0 - \rho, \omega_0 + \rho)$, we have*

$$\gamma(\omega) \cap [y = \underline{y}] \neq \emptyset \quad \text{and} \quad X_2(\alpha_-(\omega_0), \omega) < \underline{y} < \underline{y} + \epsilon < X_2(\alpha_+(\omega_0), \omega).$$

Moreover the open set \mathcal{O} can be written

$$\mathcal{O} = T_h\left(\{(t, \omega) \in D_h / |\omega - \omega_0| < \rho, t_{\underline{y}}(\omega) < t < t_{\underline{y}+\epsilon}(\omega)\}\right).$$

Proof. *i)* First we show that

$$X_2(\alpha_-(\omega_0), \omega) < \underline{y} < \underline{y} + \epsilon < X_2(\alpha_+(\omega_0), \omega) \quad \forall \omega \in (\omega_0 - \rho, \omega_0 + \rho). \quad (5.6)$$

Indeed we have for $\omega \in (\omega_0 - \rho, \omega_0 + \rho)$

$$\begin{aligned} X_2(\alpha_+(\omega_0), \omega) - X_2(t_0, \omega) &= \int_{t_0}^{\alpha_+(\omega_0)} H_2(X(s, \omega)) ds \geq \underline{h}(\alpha_+(\omega_0) - t_0) \\ X_2(\alpha_-(\omega_0), \omega) - X_2(t_0, \omega) &= - \int_{\alpha_-(\omega_0)}^{t_0} H_2(X(s, \omega)) ds \leq -\underline{h}(t_0 - \alpha_-(\omega_0)). \end{aligned}$$

Using (5.4), we get

$$\begin{aligned} X_2(\alpha_+(\omega_0), \omega) &\geq X_2(t_0, \omega_0) - \epsilon + \underline{h}(\alpha_+(\omega_0) - t_0) \\ X_2(\alpha_-(\omega_0), \omega) &\leq X_2(t_0, \omega_0) + \epsilon - \underline{h}(t_0 - \alpha_-(\omega_0)). \end{aligned}$$

Since $|X_2(t_0, \omega_0) - \underline{y}| \leq |X_2(t_0, \omega_0) - X_2(t_1, \omega_1)| < \rho < \epsilon$ (by (5.4)), we get

$$\begin{aligned} X_2(\alpha_+(\omega_0), \omega) &\geq \underline{y} - 2\epsilon + \underline{h}(\alpha_+(\omega_0) - t_0) \\ X_2(\alpha_-(\omega_0), \omega) &\leq \underline{y} + 2\epsilon - \underline{h}(t_0 - \alpha_-(\omega_0)). \end{aligned}$$

To conclude it is enough to verify that

$$-2\epsilon + \underline{h}(\alpha_+(\omega_0) - t_0) > \epsilon \quad \text{and} \quad 2\epsilon - \underline{h}(t_0 - \alpha_-(\omega_0)) < 0$$

which is assured by the choice of ϵ .

As a consequence of (5.6), we obtain by the intermediate value theorem that $\gamma(\omega) \cap [y = \underline{y}] \neq \emptyset$ for all $\omega \in (\omega_0 - \rho, \omega_0 + \rho)$.

ii) Clearly $\mathcal{O}' = T_h\left(\{(t, \omega) \in D_h, |\omega - \omega_0| < \rho, t_{\underline{y}}(\omega) < t < t_{\underline{y}+\epsilon}(\omega)\}\right) \subset \mathcal{O}$. To prove that $\mathcal{O} \subset \mathcal{O}'$, it is enough to show that

$$\forall (\omega, y) \in (\omega_0 - \rho, \omega_0 + \rho) \times [\underline{y}, \underline{y} + \epsilon] \quad \exists t_y(\omega) \in (\alpha_-(\omega_0), \alpha_+(\omega_0)) : \quad X_2(t_y(\omega), \omega) = y$$

which is a consequence of (5.6) and the continuity of the function $t \mapsto X_2(t, \omega)$. \square

Lemma 5.2. *We have*

$$|t_{\underline{y}}(\omega) - t_0| < \frac{2\epsilon}{\underline{h}} \quad \forall \omega \in (\omega_0 - \rho, \omega_0 + \rho). \quad (5.7)$$

Proof. Let $\omega \in (\omega_0 - \rho, \omega_0 + \rho)$. We have

$$\underline{y} - X_2(t_0, \omega) = X_2(t_{\underline{y}}(\omega), \omega) - X_2(t_0, \omega) = \int_{t_0}^{t_{\underline{y}}(\omega)} H_2(X(s, \omega)) ds.$$

If $\underline{y} = X_2(t_0, \omega_0)$, then by (5.4), $|\underline{y} - X_2(t_0, \omega)| < \epsilon$.

If $\underline{y} = X_2(t_1, \omega_1)$, then we have

$$|\underline{y} - X_2(t_0, \omega)| \leq |X_2(t_1, \omega_1) - X_2(t_0, \omega_0)| + |X_2(t_0, \omega_0) - X_2(t_0, \omega)|.$$

Using (5.4) and the fact that $(t_1, \omega_1) \in B_\rho(t_0, \omega_0)$, we deduce that $|\underline{y} - X_2(t_0, \omega)| < 2\epsilon$.

We conclude by distinguishing the cases $t_{\underline{y}}(\omega) > t_0$, $t_{\underline{y}}(\omega) < t_0$ and use (1.3) to conclude. \square

We claim that

Lemma 5.3.

$$\mathcal{O} \subset (x_0 - k\epsilon, x_0 + k\epsilon) \times (\underline{y}, \underline{y} + \epsilon), \quad k = c_0 \left(1 + \frac{2}{\underline{h}}\right) + \frac{\bar{h}}{\underline{h}}.$$

Proof. Indeed, let $X(t, \omega) \in \mathcal{O}$. By definition of \mathcal{O} , we have $\underline{y} < X_2(t, \omega) < \underline{y} + \epsilon$. So we only need to verify that $|X_1(t, \omega) - x_0| < k\epsilon$.

Note that, since $T_h \in C^{0,1}(D_h)$, we have

$$|X_1(t_{\underline{y}}(\omega), \omega) - X_1(t_0, \omega_0)| \leq c_0(|t_{\underline{y}}(\omega) - t_0| + |\omega - \omega_0|).$$

Using (5.7), we get for all ω in $(\omega_0 - \rho, \omega_0 + \rho)$

$$|X_1(t_{\underline{y}}(\omega), \omega) - X_1(t_0, \omega_0)| < c_0 \left(\frac{2\epsilon}{\underline{h}} + \rho\right) < c_0 \left(1 + \frac{2}{\underline{h}}\right) \epsilon. \quad (5.8)$$

We also have for $\omega \in (\omega_0 - \rho, \omega_0 + \rho)$ and $t_{\underline{y}}(\omega) < t = t_y(\omega) < t_{\underline{y}+\epsilon}(\omega)$

$$\epsilon \geq y - \underline{y} = X_2(t, \omega) - X_2(t_{\underline{y}}(\omega), \omega) = \int_{t_{\underline{y}}(\omega)}^t H_2(X(s, \omega)) ds \geq \underline{h}(t - t_{\underline{y}}(\omega))$$

$$|X_1(t, \omega) - X_1(t_{\underline{y}}(\omega), \omega)| = \left| \int_{t_{\underline{y}}(\omega)}^t H_1(X(s, \omega)) ds \right| \leq \bar{h}(t - t_{\underline{y}}(\omega)) \leq \frac{\bar{h}}{\underline{h}} \epsilon. \quad (5.9)$$

Combining (5.8) and (5.9), we obtain

$$\begin{aligned} |X_1(t, \omega) - X_1(t_0, \omega_0)| &\leq |X_1(t, \omega) - X_1(t_{\underline{y}}(\omega), \omega)| + |X_1(t_{\underline{y}}(\omega), \omega) - X_1(t_0, \omega_0)| \\ &< \left(c_0 \left(1 + \frac{2}{\underline{h}}\right) + \frac{\bar{h}}{\underline{h}}\right) \epsilon = k\epsilon \quad \forall X(t, \omega) \in \mathcal{O}. \end{aligned}$$

□

From now on, we assume that ϵ is small enough to ensure that

$$(x_0 - (k+1)\epsilon, x_0 + (k+1)\epsilon) \times (\underline{y}, \underline{y} + 2\epsilon) \subset\subset \Omega.$$

We set

$$\begin{aligned} x_1 &= x_0 - k\epsilon, & x_2 &= x_0 + k\epsilon \\ Z &= (x_1 - \epsilon, x_2 + \epsilon) \times (\underline{y}, \underline{y} + \epsilon) \\ D_{\underline{y}} &= T_h \left(\{ (t, \omega) \in D_h, \omega \in (\omega_1, \omega_0), t > t_{\underline{y}}(\omega) \} \right). \end{aligned}$$

We have

Lemma 5.4. *The line segment $S = [X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_0), \omega_0)] \subset B_{\rho^*}(X_0)$.*

Proof. Since $B_{\rho^*}(X_0)$ is convex, it suffices to prove that $X(t_{\underline{y}}(\omega_1), \omega_1), X(t_{\underline{y}}(\omega_0), \omega_0) \in B_{\rho^*}(X_0)$. First, we have $(t_1, \omega_1) \in B_{\rho}(t_0, \omega_0)$, and by (5.2), we are led to $X(t_1, \omega_1) \in B_{q\rho^*}(X_0)$. Using the definition of \underline{y} , we get

$$|\underline{y} - X_2(t_0, \omega_0)| < q\rho^* < \rho^*/4.$$

In the same way we have

$$|\underline{y} - X_2(t_1, \omega_1)| \leq |X_2(t_0, \omega_0) - X_2(t_1, \omega_1)| < q\rho^* < \rho^*/4.$$

Now, for $i = 0, 1$, we have

$$\begin{aligned} X_1(t_{\underline{y}}(\omega_i), \omega_i) - X_1(t_i, \omega_i) &= \int_{t_i}^{t_{\underline{y}}(\omega_i)} H_1(X(s, \omega_i)) ds \\ \underline{y} - X_2(t_i, \omega_i) &= X_2(t_{\underline{y}}(\omega_i), \omega_i) - X_2(t_i, \omega_i) = \int_{t_i}^{t_{\underline{y}}(\omega_i)} H_2(X(s, \omega_i)) ds \geq 0, \end{aligned}$$

from which we deduce that

$$|X_1(t_{\underline{y}}(\omega_i), \omega_i) - X_1(t_i, \omega_i)| \leq \frac{\bar{h}}{\underline{h}} |\underline{y} - X_2(t_i, \omega_i)| < \rho^*/4.$$

Hence

$$\begin{aligned} |X_1(t_{\underline{y}}(\omega_0), \omega_0) - X_1(t_0, \omega_0)| &< \rho^*/4 \\ |X_1(t_{\underline{y}}(\omega_1), \omega_1) - X_1(t_0, \omega_0)| &\leq |X_1(t_{\underline{y}}(\omega_1), \omega_1) - X_1(t_1, \omega_1)| + |X_1(t_1, \omega_1) - X_1(t_0, \omega_0)| \\ &< \rho^*/4 + q\rho^* < \rho^*/2. \end{aligned}$$

We conclude that $|X(t_{\underline{y}}(\omega_i), \omega_i) - X(t_0, \omega_0)| \leq \sqrt{(\rho^*/4)^2 + (\rho^*/2)^2} < \rho^*$. □

End of the Proof of Theorem 5.1. As a consequence of Lemma 3.1 ii) and Lemma 5.4, we have

$$u_0 T_h(t_{\underline{y}}(\omega), \omega) \leq \epsilon^2 \quad \forall \omega \in (\omega_1, \omega_0). \quad (5.10)$$

Moreover by Lemma 5.3, we have $D_{\underline{y}} \cap [\underline{y} < y < \underline{y} + \epsilon] \subset (x_1, x_2) \times (\underline{y}, \underline{y} + \epsilon)$.
We discuss the following cases :

1st case: $\overline{D_{\underline{y}}} \cap \Gamma_3 = \emptyset$

Applying Lemma 4.4, we deduce that $u \equiv 0$ in $D_{\underline{y}} \cap [y \geq \underline{y} + \epsilon]$.

Set $X'_0 = X(t'_0, \omega'_0) = X(t_{\underline{y}+\epsilon}, \omega_0)$. Arguing as before, one can find $(t_2, \omega_2) \in B_{\rho'}(t'_0, \omega'_0) \cap [\omega > \omega_0]$ such that $u \circ T_h(t_2, \omega_2) = 0$. We define $\underline{y}' = \max(X_2(t'_0, \omega'_0), X_2(t_2, \omega_2))$ and $D_{\underline{y}'} = T_h(\omega_0 < \omega < \omega_2) \cap [y > \underline{y}']$.

• If $D_{\underline{y}'} \cap \Gamma_3 = \emptyset$, then $u \equiv 0$ in $T_h(\omega_0 < \omega < \omega_2) \cap [y > \underline{y}' + \epsilon]$. So for all $\omega \in (\omega_1, \omega_2)$, we have

$$\begin{aligned} \phi_h(\omega) &\leq t_{\underline{y}'+\epsilon}(\omega) \leq t_{\underline{y}'}(\omega) + \frac{\epsilon}{h} < t'_0 + \frac{3\epsilon}{h} \\ &< t_{\underline{y}}(\omega_0) + \frac{\epsilon}{h} + \frac{3\epsilon}{h} < t_0 + 2\frac{\epsilon}{h} + \frac{4\epsilon}{h} = \phi_h(\omega_0) + \frac{6\epsilon}{h} \end{aligned}$$

which is the upper semi-continuity (u.s.c) of ϕ_h at ω_0 .

• If $D_{\underline{y}'} \cap \Gamma_3 \neq \emptyset$, then $X(\alpha_+(\omega_2), \omega_2)$ does not belong to the same connected component of $\Gamma_1 \cup \Gamma_2$ containing $X(\alpha_+(\omega_0), \omega_0)$. Moreover we are now in the situation iii) of Lemma 3.5. So there exists $(t_n^+, \omega_n^+) \in B_{\rho'}(t'_0, \omega'_0) \cap [\omega > \omega_0]$ satisfying $u \circ T_h(t_n^+, \omega_n^+) = 0$ and $X(t_n^+, \omega_n^+)_{n \rightarrow \infty} X'_0$. By Lemma 4.5, there exists $n_0 \geq 1$ such that $X(\alpha_+(\omega_{n_0}), \omega_{n_0})$ belongs to the same connected component of $\Gamma_1 \cup \Gamma_2$ which is containing $X(\alpha_+(\omega_0), \omega_0)$. Necessarily, the set $\{X(\alpha_+(\omega), \omega), \omega \in [\omega_0, \omega_{n_0}^+]\}$ is contained in this connected component. Then, by considering $D_{\underline{y}'} \cap [\omega_1 < \omega < \omega_{n_0}^+]$, we can argue as in the previous case, since $\overline{D_{\underline{y}'}} \cap [\omega_1 < \omega < \omega_{n_0}^+] \cap \Gamma_3 = \emptyset$, to show that ϕ_h is u.s.c at ω_0 .

2nd case: $\overline{D_{\underline{y}}} \cap \Gamma_3 \neq \emptyset$

From Lemma 3.5, we can have a sequence $(t_n^-, \omega_n^-)_{n \geq 1}$ in $B_{\rho}(t_0, \omega_0) \cap [\omega < \omega_0]$ or a sequence $(t_n^+, \omega_n^+)_{n \geq 1}$ in $B_{\rho}(t_0, \omega_0) \cap [\omega > \omega_0]$ or both of them, converging to X_0 and such that $u \circ T_h$ vanishes on each point of the sequences. By Lemma 4.5, we can find $\omega_{n_1}^- < \omega_0$ or $\omega_{n_2}^+ > \omega_0$ such that $X(\alpha_+(\omega_{n_1}^-), \omega_{n_1}^-)$ or $X(\alpha_+(\omega_{n_2}^+), \omega_{n_2}^+)$ or both of them belong to the same connected component of $\Gamma_1 \cup \Gamma_2$ which is containing $X(\alpha_+(\omega_0), \omega_0)$. We conclude for the last case by considering $D_{\underline{y}'} \cap [\omega_{n_1}^- < \omega < \omega_{n_2}^+]$. For the other cases, we are back to the 1st one. \square

6 Some Remarks

In this section we first propose a different proof for Theorem 5.1 when H is more regular. Then we show that conditions (4.1)-(4.2) are not sharp. Finally we show that in condition (3.1), one can replace the direction $e = (0, 1)$ by any other direction.

Remark 6.1. *When $H \in C^{1,1}(\overline{\Omega})$, it is possible to give another proof for Theorem 5.1 much simpler than the above one. It consists on using the change of variables T_h , which is now a $C^{1,1}$ diffeomorphism, to reduce the problem to a problem of type (P_0) .*

Prof of Theorem 5.1 when $H \in C^{1,1}(\bar{\Omega})$. Indeed let $h \in \pi_y(\Omega)$, $\xi \in H^1(D_h)$, $\xi = 0$ on $(\partial D_h \cap T_h^{-1}(\Gamma_3)) \cup (\partial D_h \cap \Omega)$ and $\xi \geq 0$ on $\partial D_h \cap T_h^{-1}(\Gamma_2)$. Then $\xi \circ T_h^{-1} \chi(T_h(D_h))$ is a test function for (P) and we have

$$\int_{T_h(D_h)} (a(X)\nabla u + \chi H(X)) \cdot \nabla(\xi \circ T_h^{-1}) dX \leq 0$$

which can be written using the change of variables T_h

$$\int_{D_h} (\mathbb{A}(t, \omega)\nabla(u \circ T_h) + \chi \circ T_h \cdot \mathbf{h}(t, \omega)e_t) \cdot \nabla \xi dt d\omega \leq 0$$

where the matrix \mathbb{A} and the function \mathbf{h} are given by

$$\begin{aligned} \mathbf{h}(t, \omega) &= |Y_h(t, \omega)|, & e_t &= (1, 0) \\ \mathbb{A}(t, \omega) &= |Y_h(t, \omega)|^t P(t, \omega) \cdot a(X(t, \omega)) \cdot P(t, \omega) \\ \text{with } P &= ({}^t \mathcal{J}T_h)^{-1} = \frac{1}{Y_h(t, \omega)} \begin{pmatrix} \frac{\partial X_2}{\partial \omega}(t, \omega) & -H_2(X(t, \omega)) \\ -\frac{\partial X_1}{\partial \omega}(t, \omega) & H_1(X(t, \omega)) \end{pmatrix}. \end{aligned}$$

Note that from Proposition 2.3, the function \mathbf{h} satisfies

$$\begin{cases} 0 < \underline{h} \leq \mathbf{h}(t, \omega) \leq C\bar{h} & \text{for a.e. } (t, \omega) \in D_h \\ 0 \leq \mathbf{h}_t(t, \omega) \leq C\bar{h} & \text{for a.e. } (t, \omega) \in D_h. \end{cases}$$

From the proof of Proposition 2.3, $\frac{\partial X}{\partial \omega} = U(t, \omega)$ satisfies the following differential equation

$$\begin{cases} U'(t, \omega) = DH(X(t, \omega)) \cdot U(t, \omega) \\ U(0, \omega) = (1, 0). \end{cases}$$

Arguing as in the proof of Proposition 2.3, we deduce, since $DH \in C^{0,1}(\bar{\Omega})$, that $\frac{\partial X}{\partial \omega} \in C^{0,1}(D_h)$. Moreover $\frac{1}{Y_h(t, \omega)} = -\frac{1}{H_2(\omega, h)} \exp\left(-\int_0^t (\text{div}H)(X(s, \omega)) ds\right)$ clearly belongs to $C^{0,1}(D_h)$. Hence the matrix \mathbb{A} satisfies

$$\mathbb{A} \in C^{0,1}(D_h) \quad \text{and} \quad |\mathbb{A}(t, \omega)| \leq C$$

where C is a positive constant. To conclude, it remains to verify the following ellipticity condition

$$\mathbb{A}(t, \omega)\xi \cdot \xi \geq \mu|\xi|^2 \quad \forall \xi \in \mathbb{R}^2, \quad \text{for a.e. } (t, \omega) \in D_h, \quad \text{for some positive constant } \mu.$$

So, let $\xi \in \mathbb{R}^2$. We have

$$\mathbb{A}(t, \omega)\xi \cdot \xi = |Y_h| \cdot \langle a \circ T_h \cdot P\xi, P\xi \rangle \geq \lambda|Y_h| |P\xi|^2 = \lambda|Y_h| \langle {}^t P P \xi, \xi \rangle.$$

Denote by Q the matrix ${}^t P P$. Since Q is symmetric, its eigenvalues κ_1 and κ_2 are real numbers. Moreover, we have

$$\kappa_1 \cdot \kappa_2 = \det Q = (\det P)^2 = \frac{1}{Y_h^2} \tag{6.1}$$

$$\kappa_1 + \kappa_2 = \text{tr}Q = \frac{1}{Y_h^2} \left(H_1^2 + H_2^2 + \left(\frac{\partial X_1}{\partial \omega} \right)^2 + \left(\frac{\partial X_2}{\partial \omega} \right)^2 \right). \quad (6.2)$$

Then $\kappa_1 > 0$ and $\kappa_2 > 0$. Assume for example that $\kappa_1 \leq \kappa_2$ and set $m = \inf_{(t,\omega) \in D_h} \kappa_1(t,\omega)$. Suppose $m = 0$. There exists a sequence $(t_n, \omega_n) \in D_h$ such that $m = \lim_{n \rightarrow \infty} \kappa_1(t_n, \omega_n) = 0$.

Since H and $\frac{\partial X}{\partial \omega}$ are bounded, we deduce from (6.2) that the sequence $\kappa_2(t_n, \omega_n)$ is bounded in \mathbb{R}^+ . So there exists a subsequence $(n_k)_{k \geq 1}$ such that $\lim_{k \rightarrow \infty} \kappa_2(t_{n_k}, \omega_{n_k}) = \kappa^*$ with $0 \leq \kappa^* < \infty$.

Now, letting $k \rightarrow \infty$ in (6.1), we get $\lim_{k \rightarrow \infty} \frac{1}{Y_h^2(t_{n_k}, \omega_{n_k})} = 0$ which is a contradiction with Proposition 2.3 iv). So $m > 0$.

Now since Q is symmetric, there exists an orthogonal matrix O (i.e $O^t O = {}^t O O = I_2$) such that $Q = O D O^{-1}$, D is a diagonal matrix with diagonal coefficients equal to the eigenvalues of Q . Then we have

$$\langle Q\xi, \xi \rangle = \langle D O^{-1} \xi, {}^t O \xi \rangle = \langle D {}^t O \xi, {}^t O \xi \rangle \geq m |{}^t O \xi|^2 = m |\xi|^2.$$

Hence

$$\langle \mathbb{A}\xi, \xi \rangle \geq \lambda m |Y_h| |\xi|^2 \geq \lambda m \underline{h} |\xi|^2 \quad \forall \xi \in \mathbb{R}^2.$$

We conclude (see [4]) that the free boundary $\partial[uoT_h > 0] \cap D_h$ is a continuous curve $[t = \phi_h(\omega)]$. \square

Remark 6.2. *The conditions under which Theorem 5.1 is proved are not sharp. Indeed we present below a proof when $H(X) = a(X)e$, that is to say when (P) is the weak formulation of the dam problem with Dirichlet boundary conditions, with $a(X)$ satisfying (1.1)-(1.2), $a(X)e \in C^{0,1}(\bar{\Omega})$, but not the assumptions (4.1)-(4.2). Note that only the proof of Lemma 4.4. requires the last assumptions. Actually the proof given in section 4 is based on the comparison of u with respect to the barrier function defined by (4.3). It uses the local Lipschitz continuity of u which requires the assumptions (4.1)-(4.2). For this special case, we propose another proof using an explicit barrier function. Moreover the assumption “ $uoT_h(t_{\underline{y}}(\omega), \omega) \leq \epsilon^2 \forall \omega \in (\omega_1, \omega_2)$ ” in Lemma 4.3, will be modified by changing ϵ^2 to ϵ .*

Proof of lemma 4.4 when $H(X) = a(X)e$. Let $v(y) = (\epsilon + \underline{y} - y)^+$ and $\xi(x, y) = \chi(D_{\underline{y}})(u - v)^+$. Since $v \geq 0 = u$ on $(\partial D_{\underline{y}} \setminus ([y = \underline{y}])) \cap \Omega$, we have $\xi = 0$ on $(\partial D_{\underline{y}} \setminus ([y = \underline{y}])) \cap \Omega$. Moreover $v(\underline{y}) = \epsilon \geq u(x, \underline{y})$ and then $\xi(x, \underline{y}) = 0$. It follows that $\xi = 0$ on $(\partial D_{\underline{y}} \cap \Omega) \cup (\partial D_{\underline{y}} \cap \Gamma_2)$, and $\pm \xi$ are test functions for (P). So we have

$$\int_{D_{\underline{y}}} (a(X)\nabla u + \chi a(X)e) \cdot \nabla (u - v)^+ dX \leq 0. \quad (6.3)$$

We also have

$$\int_{D_{\underline{y}}} (a(X)\nabla v + \chi([v > 0])a(X)e) \cdot \nabla (u - v)^+ dX = 0. \quad (6.4)$$

Subtracting (6.4) from (6.3), we obtain

$$\int_{D_{\underline{y}} \cap [v > 0]} a(X)\nabla (u - v) \cdot \nabla (u - v)^+ dX$$

$$+ \int_{D_{\underline{y}} \cap [v=0]} a(X)(\nabla u + \chi e) \cdot \nabla u dX \leq 0. \quad (6.5)$$

By Lemma 3.2, we have for $D_{\underline{y}+\epsilon} = [y > \underline{y} + \epsilon] \cap D_{\underline{y}} = D_{\underline{y}} \cap [v = 0]$ and $\zeta = y - (\underline{y} + \epsilon)$

$$\int_{D_{\underline{y}} \cap [v=0]} a(X)(\nabla u + \chi e) \cdot e dX \leq 0. \quad (6.6)$$

Adding (6.5) and (6.6), we get by taking into account (P)i)

$$\begin{aligned} & \int_{D_{\underline{y}} \cap [v>0]} a(X) \nabla(u-v) \cdot \nabla(u-v)^+ dX \\ & + \int_{D_{\underline{y}} \cap [u>v=0]} a(X)(\nabla u + e) \cdot (\nabla u + e) dX \\ & + \int_{D_{\underline{y}} \cap [u=v=0]} \chi a(X) e \cdot e dX \leq 0. \end{aligned}$$

or by (1.2)

$$\int_{D_{\underline{y}} \cap [v>0]} |\nabla(u-v)^+|^2 dX + \int_{D_{\underline{y}} \cap [u>v=0]} |\nabla u + e|^2 dX + \int_{D_{\underline{y}} \cap [u=v=0]} \chi dX \leq 0.$$

Since the three integrals in the left hand side of the above inequality are all nonnegative, we obtain $\nabla(u-v)^+ = 0$ a.e. in $D_{\underline{y}} \cap [v > 0]$ and then, since $(u-v)^+ = 0$ on $\partial D_{\underline{y}} \cap [y = \underline{y}]$, we get $u \leq v$ in $D_{\underline{y}} \cap [v > 0]$. This leads to $u(x, \underline{y} + \epsilon) = 0 \forall x \in \pi_x(D_{\underline{y}} \cap [y = \underline{y} + \epsilon])$. Hence $u = 0$ in $D_{\underline{y}} \cap [y \geq \underline{y} + \epsilon]$. \square

Remark 6.3. *The assumption (1.3) can be replaced by the more general one*

$$|H_1(X)| \leq \bar{h}, \quad 0 < \underline{h} \leq H(X) \cdot \nu \leq \bar{h} \quad \text{a.e. } X \in \Omega \quad (6.7)$$

where $\nu \neq 0$ is a constant vector.

Proof of Remark 6.3 Indeed, set $\nu = (\nu_1, \nu_2)$, $n = (-\nu_2, \nu_1)$. We can assume that $|\nu| = \nu_1^2 + \nu_2^2 = 1$. Clearly (n, ν) is an orthonormal basis of \mathbb{R}^2 .

For a point $M \in \Omega$, we denote by X (resp. Y) its coordinates in the canonical (resp. new) basis (e_1, e_2) (resp. (n, ν)). We have

$$X = RY \quad \text{with} \quad R = R^{-1} = \begin{pmatrix} -\nu_2 & \nu_1 \\ \nu_1 & \nu_2 \end{pmatrix}.$$

Consider the change of variables $\theta : Y \mapsto X = RY$ from $\theta^{-1}(\Omega) = \tilde{\Omega}$ into Ω . Let $\xi \in H^1(\tilde{\Omega})$, $\xi = 0$ on $\tilde{\Gamma}_3$, $\xi \geq 0$ on $\tilde{\Gamma}_2$, where $\tilde{\Gamma}_i = \theta^{-1}(\Gamma_i)$ for $i = 1, 2, 3$. Using $\xi \circ \theta^{-1}$ as a test function for (P), we obtain

$$\begin{aligned}
& \int_{\Omega} (a(X)\nabla u + \chi H(X)) \cdot \nabla(\xi o\theta^{-1}) dX \\
&= \int_{\theta^{-1}(\Omega)} (R.a o\theta.R\nabla_Y(u o\theta) + \chi o\theta R.H o\theta) \cdot \nabla_Y \xi dY \\
&= \int_{\tilde{\Omega}} (\tilde{a}(Y)\nabla \tilde{u} + \tilde{\chi}\tilde{H}(Y)) \nabla \xi dY.
\end{aligned}$$

where $\tilde{a}(Y) = R.a o\theta(Y).R$, $\tilde{u} = u o\theta$, $\tilde{\chi} = \chi o\theta$, and $\tilde{H}(Y) = R.H o\theta(Y)$. Note that $H o\theta = H_1 o\theta e_1 + H_2 o\theta e_2 = \tilde{H}_1(Y)n + \tilde{H}_2(Y)\nu = R.\tilde{H}(Y)$. Then

$$\tilde{H}_1(Y) = -\nu_2 H_1 o\theta(Y) + \nu_1 H_2 o\theta(Y) \quad \text{and} \quad \tilde{H}_2(Y) = \nu_1 H_1 o\theta(Y) + \nu_2 H_2 o\theta(Y) = H o\theta(Y).\nu.$$

We deduce that

$$|\tilde{H}_1(Y)| \leq 2\bar{h}, \quad 0 < \underline{h} \leq \tilde{H}_2(Y) \leq \bar{h} \quad \text{a.e. } Y \in \theta^{-1}(\Omega).$$

Finally, one can check easily that $(\operatorname{div}_Y \tilde{H})(Y) = (\operatorname{div}_X H)(X)$ from which we deduce that

$$\operatorname{div}_Y \tilde{H} \in L^\infty(\theta^{-1}(\Omega)) \quad \text{and} \quad (\operatorname{div}_Y \tilde{H})(Y) \geq 0 \quad \text{a.e. } Y \in \theta^{-1}(\Omega).$$

Similarly one can check that $\tilde{a}(Y)$ satisfies the assumptions (1.1)-(1.2) and (4.1)-(4.2). \square

Acknowledgments We would like to thank KFUPM for its continuing support and facilities provided.

References

- [1] Amann, H. : Ordinary Differential Equations. An Introduction to Nonlinear Analysis. Walter de Gruyter. Berlin. New York 1990.
- [2] Federer, H. : Geometric Measure Theory. Springer-Verlag 1969.
- [3] Challal, S., Lyaghfour, A. : A Filtration Problem through a Heterogeneous Porous Medium. Interfaces and Free Boundaries 6, 55-79 (2004).
- [4] Challal, S., Lyaghfour, A. : Continuity of the Free Boundary in Problems of type $\operatorname{div}(a(x)\nabla u) = -(\chi(u)h(x))_{x_1}$. Nonlinear Analysis : Theory, Methods & Applications, Vol. 62, No. 2, 283-300 (2005).
- [5] Gilbarg, D., Trudinger, N.S. : Elliptic Partial Differential Equations of Second Order. Springer-Verlag 1983.
- [6] Hale, J. : Ordinary Differential Equations. Second Edition. Robert E. Krieger Publishing Co., Inc., Huntington, N.Y., 1980.
- [7] Ziemer, W.P. : Weakly Differential Functions. Graduate Texts in Mathematics 120. Springer-Verlag 1989.