

# 1 (1.3) Subdifferentials of nonsmooth functions

- $X$  is a Banach space,  $\varphi : X \rightarrow \overline{\mathbb{R}}$ .
- $\varphi$  is called proper if  $\varphi(x) > -\infty \forall x \in X$ . The domain of  $\varphi$  is defined as

$$\text{dom } \varphi := \{x \in X : \varphi(x) < \infty\} \neq \emptyset.$$

- The epigraph and hypergraph of  $\varphi$  are defined by

$$\begin{aligned} \text{epi } \varphi & : = \{(x, a) \in X \times \mathbb{R} : a \geq \varphi(x)\}, \\ \text{hypo } \varphi & : = \{(x, a) \in X \times \mathbb{R} : a \leq \varphi(x)\}. \end{aligned}$$

- Observe that  $\text{gr } \varphi = \text{epi } \varphi \cap \text{hypo } \varphi$ .
- $\text{epi } \varphi$  is locally closed at  $(\bar{x}, \varphi(\bar{x}))$  if and only if  $\varphi$  is lower semi-continuous around  $\bar{x}$ .
- $\text{hypo } \varphi$  is locally closed at  $(\bar{x}, \varphi(\bar{x}))$  if and only if  $\varphi$  is upper semi-continuous around  $\bar{x}$ .
- $\text{gr } \varphi$  is locally closed at  $(\bar{x}, \varphi(\bar{x}))$  if and only if  $\varphi$  is continuous around  $\bar{x}$ .
- The notation  $x \xrightarrow{\varphi} \bar{x}$  means  $(x, \varphi(x)) \rightarrow (\bar{x}, \varphi(\bar{x}))$ .

## 1.1 (1.3.1) Basic definitions and results

**Proposition 1** (1.76 basic normals to epigraphs)

Suppose  $\varphi : X \rightarrow \overline{\mathbb{R}}$ ,  $(\bar{x}, \bar{\alpha}) \in \text{epi } \varphi$ . Then  $\lambda \geq 0$  for all  $(x^*, -\lambda) \in N((\bar{x}, \bar{\alpha}); \text{epi } \varphi)$ .

- There are uniquely defined subsets  $D$  and  $D^\infty$  of  $X^*$  such that

$$N((\bar{x}, \varphi(\bar{x})); \text{epi } \varphi) = \{\lambda(x^*, -1) : x^* \in D, \lambda > 0\} \cup \{(x^*, 0) : x^* \in D^\infty\}.$$

**Definition 2** (1.77 basic and singular subdifferentials)

Suppose  $\varphi : X \rightarrow \overline{\mathbb{R}}$ ,  $\bar{x} \in X$ ,  $|\varphi(\bar{x})| < \infty$ .

- (i) The set

$$\partial\varphi(\bar{x}) = \{x^* \in X^* : (x^*, -1) \in N((\bar{x}, \varphi(\bar{x})); \text{epi } \varphi)\}$$

is the basic (limiting) subdifferential of  $\varphi$  at  $\bar{x}$ . Its elements are the basic subgradients of  $\varphi$  at  $\bar{x}$ . (We put  $\partial\varphi(\bar{x}) = \emptyset$  if  $|\varphi(\bar{x})| = \infty$ ).

- (ii) The set

$$\partial^\infty\varphi(\bar{x}) = \{x^* \in X^* : (x^*, 0) \in N((\bar{x}, \varphi(\bar{x})); \text{epi } \varphi)\}$$

is the singular subdifferential of  $\varphi$  at  $\bar{x}$ . Its elements are singular subdifferentials of  $\varphi$  at  $\bar{x}$ . (We put  $\partial^\infty\varphi(\bar{x}) = \emptyset$  if  $|\varphi(\bar{x})| = \infty$ ).

**Definition 3** (1.78 upper subgradients)

Suppose  $\varphi : X \rightarrow \overline{\mathbb{R}}$ ,  $\bar{x} \in X$ ,  $|\varphi(\bar{x})| < \infty$ .

(i) The set

$$\partial^+ \varphi(\bar{x}) = \{x^* \in X^* : (-x^*, 1) \in N((\bar{x}, \varphi(\bar{x})); \text{hypo } \varphi)\}$$

is the upper (basic, limitting) subdifferential of  $\varphi$  at  $\bar{x}$ . (We put  $\partial\varphi(\bar{x}) = \phi$  if  $|\varphi(\bar{x})| = \infty$ ).

(ii) The set

$$\partial^{\infty,+} \varphi(\bar{x}) = \{x^* \in X^* : (-x^*, 0) \in N((\bar{x}, \varphi(\bar{x})); \text{hypo } \varphi)\}$$

is the singular upper subdifferential of  $\varphi$  at  $\bar{x}$ . (We put  $\partial^\infty \varphi(\bar{x}) = \phi$  if  $|\varphi(\bar{x})| = \infty$ ).

- Subgradients and upper subgradients may be substantially different.

**Example** For  $\varphi(x) = -|x|$ ,  $\partial\varphi(0) = \{-1, 1\}$  while  $\partial^+ \varphi(0) = [-1, 1]$ .

- $\partial^+ \varphi(\bar{x}) = -\partial(-\varphi)(\bar{x})$ ,  $\partial^{\infty,+} \varphi(\bar{x}) = -\partial^{\infty}(-\varphi)(\bar{x})$ .
- The symmetric and singular symmetric subdifferentials are defined by

$$\begin{aligned} \partial^0 \varphi(\bar{x}) & : = \partial\varphi(\bar{x}) \cup \partial^+ \varphi(\bar{x}), \\ \partial^{\infty,0} \varphi(\bar{x}) & = \partial^{\infty,+} \varphi(\bar{x}) \cup \partial^\infty \varphi(\bar{x}) \end{aligned}$$

respectively.

- We have

$$\begin{aligned} \partial^0(-\varphi)(\bar{x}) & = -\partial^0 \varphi(\bar{x}), \\ \partial^{\infty,0}(-\varphi)(\bar{x}) & = -\partial^{\infty,0} \varphi(\bar{x}). \end{aligned}$$

- Since properties of upper subdifferentials can be deduced from the corresponding ones for subgradients, we deal from now on only with the subdifferential  $\partial$ .
- For  $\Omega \subset X$ , the indicator function  $\delta(\cdot; \Omega) : X \rightarrow \overline{\mathbb{R}}$  is defined by

$$\delta(x; \Omega) = \begin{cases} 0, & x \in \Omega \\ \infty, & x \notin \Omega. \end{cases}$$

**Proposition 4** (1.79 subdifferentials of indicator functions)

We have

$$\partial\delta(\bar{x}; \Omega) = \partial^\infty \delta(\bar{x}; \Omega) = N(\bar{x}; \Omega).$$

- For a function  $\varphi : X \rightarrow \overline{\mathbb{R}}$ , the epigraphical function  $E_\varphi : X \rightarrow \mathbb{R}$  is defined by

$$E_\varphi(x) = \{\alpha \in \mathbb{R} : \alpha \geq \varphi(x)\}.$$

- $\text{gr } E_\varphi = \text{epi } \varphi$ .
- $D_N^* E_\varphi = D_M^* E_\varphi$  since  $E_\varphi$  takes values in  $\mathbb{R}$ .
- $\partial\varphi(\bar{x}) = D^* E_\varphi(\bar{x}, \varphi(\bar{x}))(1)$  and  $\partial^\infty\varphi(\bar{x}) = D^* E_\varphi(\bar{x}, \varphi(\bar{x}))(0)$ .

**Theorem 5** (1.80 subdifferentials from coderivatives of continuous functions)  
 Suppose  $\varphi : X \rightarrow \overline{\mathbb{R}}$  is continuous around  $\bar{x} \in X$ . Then

$$\begin{aligned}\partial\varphi(\bar{x}) &= D^*\varphi(\bar{x})(1), \\ \partial^\infty\varphi(\bar{x}) &\subset D^*\varphi(\bar{x})(0).\end{aligned}$$

- The inclusion in Theorem 1.80 for  $\partial^\infty$  could be strict.

### Example

**Corollary 6** (1.81 subdifferentials of Lipschitzian functions)

Suppose  $\varphi : X \rightarrow \overline{\mathbb{R}}$  is Lipschitz continuous around  $\bar{x} \in X$  with modulus  $\ell \geq 0$ . Then

$$\begin{aligned}\partial^\infty\varphi(\bar{x}) &= \{0\}, \\ \|x^*\| &\leq \ell \quad \forall x^* \in \partial\varphi(\bar{x}).\end{aligned}$$

- It follows that, if  $\varphi$  is Lipschitz continuous around  $\bar{x}$  then  $\partial^{\infty,0}\varphi(\bar{x}) = \{0\}$  and  $\|x^*\| \leq \ell \quad \forall x^* \in \partial^0\varphi(\bar{x})$ .

**Corollary 7** (1.82 subdifferentials of strictly differentiable functions)

Suppose  $\varphi : X \rightarrow \overline{\mathbb{R}}$  is strictly differentiable at  $\bar{x} \in X$ . Then

$$\partial\varphi(\bar{x}) = \partial^+\varphi(\bar{x}) = \partial^0\varphi(\bar{x}) = \{\varphi'(\bar{x})\}.$$

- If  $\varphi$  is not strictly differentiable at  $\bar{x}$ , it could happen that the equalities in Corollary 1.82 do not hold. For example, the function

$$\varphi(x) = \begin{cases} x^2 \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

is differentiable but not strictly differentiable at 0. Here,  $\varphi'(0) = 0$  while  $\partial\varphi(\bar{x}) = \partial^+\varphi(\bar{x}) = [-1, 1]$ .