

**Q.1:** Let parametric equations of a curve are:  $x(t) = 2 + 3t^2$  and  $y(t) = 3 + 2t^3$ .

(a) Find  $\frac{dx}{dt}$  (1 pts)  $\frac{dx}{dt} = 6t$

(b) Find  $\frac{dy}{dt}$  (1 pts)  $\frac{dy}{dt} = 6t^2$

(c) Find  $\frac{dy}{dx}$  (2 pts)  $\frac{dy}{dx} = t$


(d) Find  $\frac{d^2y}{dx^2}$  (3 pts)  $\frac{d^2y}{dx^2} = \frac{1}{6t}$

(e) Find length of the curve for  $0 \leq t \leq 1$ . (8 pts)

$$\begin{aligned} \text{Arc Length } s &= \int_0^1 \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^1 \sqrt{36t^2 + 36t^4} dt \\ &= 3 \int_0^1 2t\sqrt{1+t^2} dt = 3 \left. \frac{2}{3} (1+t^2)^{\frac{3}{2}} \right|_0^1 = 2(2\sqrt{2} - 1) \end{aligned}$$

**Q.2:** Sketch graph of the polar curve  $r = 2 + 2\sin(\theta)$ . Show the points on the curve for  $\theta = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi$ . (4 pts)

$\theta$	$r$
0	2
$\frac{\pi}{2}$	4
$\pi$	2
$\frac{3\pi}{2}$	0
$2\pi$	2



$$x = r \cos(\theta) = (2 + 2\sin(\theta)) \cos(\theta) = 2\cos(\theta) + 2\sin(\theta)\cos(\theta) = 2\cos(\theta) + \sin(2\theta)$$

$$y = r \sin(\theta) = (2 + 2\sin(\theta)) \sin(\theta) = 2\sin(\theta) + 2\sin^2(\theta)$$

(a) Find  $\frac{dx}{d\theta}$  (2 pts)  $\frac{dx}{d\theta} = -2\sin(\theta) + 2\cos(2\theta)$

(b) Find  $\frac{dy}{d\theta}$  (2 pts)  $\frac{dy}{d\theta} = 2\cos(\theta) + 4\sin(\theta)\cos(\theta) = 2\cos(\theta) + 2\sin(2\theta)$

(c) Find  $\frac{dy}{dx}$  (2 pts)  $\frac{dy}{dx} = \frac{2\cos(\theta) + 2\sin(2\theta)}{-2\sin(\theta) + 2\cos(2\theta)}$


(d) Find equation of tangent line at  $\theta = \frac{\pi}{4}$  (5 pts)

$$\text{Slope } m = \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{4}} = \left. \frac{2\cos(\theta) + 2\sin(2\theta)}{-2\sin(\theta) + 2\cos(2\theta)} \right|_{\theta=\frac{\pi}{4}} = \frac{2\cos(\frac{\pi}{4}) + 2\sin(\frac{\pi}{2})}{-2\sin(\frac{\pi}{4}) + 2\cos(\frac{\pi}{2})} = -\sqrt{2} - 1$$

Also at  $\theta = \frac{\pi}{4}$ ,  $x = 2\cos(\frac{\pi}{4}) + \sin(\frac{\pi}{2}) = \sqrt{2} + 1$  and  $y = 2\sin(\frac{\pi}{4}) + 2\sin^2(\frac{\pi}{4}) = \sqrt{2} + 1$

Equation of tangent line is  $(y - \sqrt{2} - 1) = (-\sqrt{2} - 1)(x - \sqrt{2} - 1)$ .

**Q.3:** Find area of the region that lies inside  $r = 2$  and outside  $r = 2\sin(2\theta)$ . (10 pts)



Area of one loop is  $A = \frac{1}{2} \int_0^{\frac{\pi}{2}} 4 \sin^2(t) dt = \frac{1}{2} \pi$ . Total area of rose is  $4A = 2\pi$  and area outside of rose and inside of circle is  $4\pi - 2\pi = 2\pi$ .

**Q.4:** If  $r = \langle x, y, z \rangle$ ,  $a = \langle 2, 3, 1 \rangle$ , and  $b = \langle -2, 1, 5 \rangle$ . Show that the vector equation  $(r - a) \cdot (r - b)$  represents a sphere. Find center and radius of the sphere (10 pts)

$$(r - a) = \langle x - 2, y - 3, z - 1 \rangle \text{ and } (r - b) = \langle x + 2, y - 1, z - 5 \rangle$$

$$(r - a) \cdot (r - b) = (x - 2)(x + 2) + (y - 3)(y - 1) + (z - 1)(z - 5) = x^2 - 6z - 4y + y^2 + z^2 + 4$$

$$(r - a) \cdot (r - b) = 0 \Rightarrow x^2 - 6z - 4y + y^2 + z^2 + 4 = 0 \Rightarrow x^2 + y^2 - 4y + 4 + z^2 - 6z + 9 = 9 \Rightarrow x^2 + (y - 2)^2 + (z - 3)^2 = 3^2$$

The equation represents sphere with center at  $(0, 2, 3)$  and radius equal to 3.

**Q.5:** Find scalar and vector projection of the vector  $\mathbf{u} = 3\mathbf{i} - \mathbf{j} + 2\mathbf{k}$  in the direction of  $\mathbf{v} = 2\mathbf{i} + 3\mathbf{j} - 4\mathbf{k}$ . (5+5 pts)

$$\text{Comp}_{\mathbf{v}} \mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|} = \frac{6 - 3 - 8}{\sqrt{29}} = \frac{-5}{\sqrt{29}} \text{ and } \text{Proj}_{\mathbf{v}} \mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|} \left( \frac{\mathbf{v}}{\|\mathbf{v}\|} \right) = \frac{-5}{\sqrt{29}} \left( \frac{2\mathbf{i} + 3\mathbf{j} - 4\mathbf{k}}{\sqrt{29}} \right) = \frac{-10\mathbf{i} - 15\mathbf{j} + 20\mathbf{k}}{29}$$

**Q.6:** Determine whether the three vectors  $\mathbf{u} = \mathbf{i} + 4\mathbf{j} - 7\mathbf{k}$ ,  $\mathbf{v} = 2\mathbf{i} - \mathbf{j} + 4\mathbf{k}$ , and  $\mathbf{w} = 0\mathbf{i} - 9\mathbf{j} + 18\mathbf{k}$  are coplanar or not. Show all your work. (10 pts)

$$\mathbf{v} \times \mathbf{w} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 4 \\ 0 & -9 & 18 \end{vmatrix} = 18\mathbf{i} - 36\mathbf{j} - 18\mathbf{k} \text{ and } \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = 18 + 4(-36) - 7(-18) = 0.$$

The three vectors are coplanar.

**Q.7:** Find equation of a line containing the point  $(1, 2, -3)$  and is perpendicular to the plane  $3x - 2y + z = 5$ . Write your answer in symmetric and parametric form. (10 pts)

Since line is perpendicular to the plane therefore it is parallel to the normal of the plane which is  $\vec{n} = \langle 3, -2, 1 \rangle$ . Thus equation of the line is:

$$\text{Symmetric form } \frac{(x - 1)}{3} = \frac{(y - 2)}{-2} = \frac{(z + 3)}{1} = t$$

$$\text{Parametric form } x = 3t + 1, y = -2t + 2, z = t - 3.$$

**Q.8:** Find equation of a plane passing through the points  $(1, 2, 3)$ ,  $(-3, 4, 2)$ , and  $(2, 4, 5)$ . (10 pts)

Let  $P(1, 2, 3)$ ,  $Q(-3, 4, 2)$ , and  $R(2, 4, 5)$ . Then two vectors in the plane are

$P\vec{Q} = \langle -4, 2, -1 \rangle$  and  $P\vec{R} = \langle 1, 2, 2 \rangle$ . Cross product of these two vectors is a vector normal to the plane.

$$P\vec{Q} \times P\vec{R} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & 2 & -1 \\ 1 & 2 & 2 \end{vmatrix} = 6\mathbf{i} + 7\mathbf{j} - 10\mathbf{k}. \text{ Thus equation of the plane passing through } P(1, 2, 3) \text{ and } \vec{n} = 6\mathbf{i} + 7\mathbf{j} - 10\mathbf{k} \text{ is } 6x + 7y - 10z = 1(6) + 2(7) + 3(-10) = -10.$$

**Q.9:** Find angle between the planes  $3x - 2y + z = 5$  and  $2x + y - 4z = 5$ . (5 pts)

Normal to these planes are  $\vec{n}_1 = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$  and  $\vec{n}_2 = 2\mathbf{i} + \mathbf{j} - 4\mathbf{k}$ . Angle between two planes is the angle between their normals. Thus  $\cos(\theta) = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|} = 0 \Rightarrow \theta = \frac{\pi}{2}$ .

**Q.10:** Prove that  $(\mathbf{u} - \mathbf{v}) \times (\mathbf{u} + \mathbf{v}) = 2(\mathbf{u} \times \mathbf{v})$  (5 pts)

$$(\mathbf{u} - \mathbf{v}) \times (\mathbf{u} + \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} - \mathbf{v} \cdot \mathbf{v} = \mathbf{0} + \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{v} - \mathbf{0} = 2(\mathbf{u} \cdot \mathbf{v}).$$