

King Fahd University of Petroleum and Minerals
 Department of Mathematical Sciences
 Solution Exam 1 for Math 201 (081)

Dr. Muhammad Yousuf

Q.1: (12 pts) Find equations of all tangent lines to the parametric curve given by $x = t^5 - 4t^3$, $y = t^2$, at $(0, 4)$.

Sol: $x = 0$ if $t^3(t^2 - 4) = 0 \Rightarrow t = 0, \pm 2$ and $y = 4$ if $t^2 = 4 \Rightarrow t = \pm 2$.
 The common values of t at which $x = 0$ and $y = 4$ are ± 2 .

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{2t}{5t^4 - 12t^2}$$


At $t = 2$, $m = \left. \frac{dy}{dx} \right|_{t=2} = \frac{2(2)}{5(2)^4 - 12(2)^2} = \frac{1}{8}$ and equation of tangent line is $y = \frac{1}{8}x + 4$.

At $t = -2$, $m = \left. \frac{dy}{dx} \right|_{t=-2} = \frac{2(-2)}{5(-2)^4 - 12(-2)^2} = -\frac{1}{8}$ and equation of tangent line is $y = -\frac{1}{8}x + 4$.

Q.2: (14 pts) Consider the polar curve $C : r = f(\theta) = \cos(2\theta)$, $-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$

- (a) Sketch the curve.
- (b) Setup the integral for the area enclosed by the curve.
- (c) Setup the integral for the arc length of curve.

Sol: (a) The graph of $r = f(\theta) = \cos(2\theta)$




(b) $A = \frac{1}{2} \int_{-\pi/4}^{\pi/4} \cos^2(2\theta) d\theta$.

(c) $S = \int_{-\pi/4}^{\pi/4} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta = \int_{-\pi/4}^{\pi/4} \sqrt{\cos^2 \theta + 4 \sin^2(2\theta)} d\theta$.

Q.3: Find the area inside the curve $r = 1 + \sin(\theta)$ and outside the curve $r = \sin(\theta)$ when $\frac{\pi}{3} \leq \theta \leq \frac{\pi}{2}$.

Sol:



$$\begin{aligned}
A &= \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} ((1 + \sin(\theta))^2 - \sin^2(\theta)) d\theta \\
&= \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} (1 + \sin^2(\theta) + 2\sin(\theta) - \sin^2(\theta)) d\theta \\
&= \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} (1 + 2\sin(\theta)) d\theta = \frac{1}{12}\pi + \frac{1}{2}.
\end{aligned}$$

Q.4: (9 pts) Find an equation of the sphere if one of its diameters has end points at $A(1, 4, -2)$ and $B(-7, 1, 2)$. What is the intersection of this sphere with the xz -plane?

Sol: Center is $\left(\frac{1-7}{2}, \frac{4+1}{2}, \frac{-2+2}{2}\right) = \left(-3, \frac{5}{2}, 0\right)$ and radius is $r = \frac{1}{2}\sqrt{(-8)^2 + (-3)^2 + (4)^2} = \frac{1}{2}\sqrt{89}$.

$$\text{Equation of the sphere is } (x+3)^2 + \left(y - \frac{5}{2}\right)^2 + (z-0)^2 = \frac{89}{4}.$$

$$\text{Intersection with } xz\text{-plane: Set } y = 0, \text{ then } (x+3)^2 + \left(0 - \frac{5}{2}\right)^2 + (z-0)^2 = \frac{89}{4}$$

$$(x+3)^2 + z^2 = \frac{89}{4} - \frac{25}{4} = \frac{64}{4} = 16, \text{ a circle in the } xz\text{-plane with radius 4 and center at } (-3, 0, 0).$$

Q.5: (9 pts) Find the distance from the point $P(1,1,1)$ to the line passing through the points $Q(2,-1,3)$ and $R(5,0,1)$.

Sol: $Q\vec{P} = \langle -1, 2, -2 \rangle$ and $Q\vec{R} = \langle 3, 1, -2 \rangle$.

$$Q\vec{R} \times Q\vec{P} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & -2 \\ -1 & 2 & -2 \end{vmatrix} = 2\hat{i} + 8\hat{j} + 7\hat{k}.$$

$$D = \frac{|Q\vec{R} \times Q\vec{P}|}{|Q\vec{R}|} = \frac{|2\hat{i} + 8\hat{j} + 7\hat{k}|}{|3\hat{i} + \hat{j} - 2\hat{k}|} = \frac{\sqrt{4 + 64 + 49}}{\sqrt{9 + 1 + 4}} = \frac{\sqrt{117}}{\sqrt{14}} = \frac{3}{14}\sqrt{182}.$$

Q.6: (8 pts) Find the Cartesian equation of the curve whose polar equation is given by $r = \sec(\theta) - \csc(\theta)$.

Sol: $r = \frac{1}{\cos \theta} - \frac{1}{\sin \theta} = \frac{\sin \theta - \cos \theta}{\sin \theta \cos \theta}$
 $r \sin \theta \cdot \cos \theta = \sin \theta - \cos \theta$
 $r \sin \theta \cdot r \cos \theta = r \sin \theta - r \cos \theta$
 $xy = y - x$
 $y = \frac{x}{1-x}$.

OR

$$r = r = \frac{1}{\cos \theta} - \frac{1}{\sin \theta} = \frac{1}{\frac{x}{r}} - \frac{1}{\frac{y}{r}} = \frac{r}{x} - \frac{r}{y} = r \left(\frac{1}{x} - \frac{1}{y} \right)$$

$$1 = \frac{1}{x} - \frac{1}{y} \Rightarrow xy = y - x \Rightarrow y = \frac{x}{1-x}.$$

OR

$$x = r \cos \theta = (\sec \theta - \csc \theta) \cos \theta = 1 - \tan \theta$$

$$x = 1 - \frac{y}{x} \Rightarrow xy = y - x \Rightarrow y = \frac{x}{1-x}.$$

Q.7: (6 pts) The graph of the curve represented by $x = 4 \cos \theta$, and $y = 5 \sin \theta$, is:

Sol: $x^2 = 16 \cos^2 \theta$, $y^2 = 25 \sin^2 \theta$

$$\frac{x^2}{16} + \frac{y^2}{25} = \cos^2 \theta + \sin^2 \theta = 1$$

This is an ellipse with center at the origin $(0, 0)$ and intercepts $(\pm 4, 0)$ and $(0, \pm 5)$.

Q.8: (6 pts) What does the polar equation $r = \tan \theta \sec \theta$ represents ?

Sol: $r = \frac{\sin \theta}{\cos \theta} \frac{1}{\cos \theta} = \frac{\sin \theta}{\cos^2 \theta} \Rightarrow r \cos^2 \theta = \sin \theta \Rightarrow r^2 \cos^2 \theta = r \sin \theta$
 $y^2 = x$, a parabola.

Q.9 : (6 pts) The vector that has the same direction as $\langle -3, 4, 1 \rangle$ but has length 5 is:

Sol: $|\langle -3, 4, 1 \rangle| = \sqrt{9 + 16 + 1} = \sqrt{26}$

A unit vector in the direction of $\langle -3, 4, 1 \rangle$ is $\langle -\frac{3}{\sqrt{26}}, \frac{4}{\sqrt{26}}, \frac{1}{\sqrt{26}} \rangle$.

A vector of length 5 in the direction of $\langle -3, 4, 1 \rangle$ is $\langle -\frac{15}{\sqrt{26}}, \frac{20}{\sqrt{26}}, \frac{5}{\sqrt{26}} \rangle$.

Q.10: (6 pts) The vector projection of $u = \hat{i} + 2\hat{j} + 3\hat{k}$ onto $v = 5\hat{i} - \hat{j} + 2\hat{k}$ is given by:


Sol: $Proj_{\vec{v}} \vec{u} = \left(\frac{\vec{u} \cdot \vec{v}}{|\vec{v}|^2} \right) \vec{v} = \frac{5 - 2 + 6}{25 + 1 + 4} \langle 5, -1, 2 \rangle = \frac{3}{10} \langle 5, -1, 2 \rangle = \langle \frac{3}{2}, -\frac{3}{10}, \frac{3}{5} \rangle$.

Q.11: (6 pts) The Cartesian equation for the parametric equations $x = 9 \sec t$ and $y = 8 \tan t$ is:

Sol: $\frac{x^2}{9^2} = \sec^2 t$ and $\frac{y^2}{8^2} = \tan^2 t$
 $\frac{x^2}{81} - \frac{y^2}{64} = \sec^2 t - \tan^2 t = 1$, a hyperbola.

Q.12: (6 pts) Let P be the parallelogram in with vertices $A = (1, -1, 2)$, $B = (2, 0, 1)$, $C = (3, 2, -1)$, $D = (2, 1, 0)$. The area of P is:

Sol: The vectors $\vec{AB} = \langle 1, 1, -1 \rangle$ and $\vec{DC} = \langle 1, 1, -1 \rangle$ shows that sides AB and DC are parallel with AB and AD as two adjacent sides.



Now $\vec{AD} = \langle 1, 2, -2 \rangle$ and $\vec{AB} \times \vec{AD} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & -1 \\ 1 & 2 & -1 \end{vmatrix} = \hat{i} + \hat{k}$

Area of P is $\mathbf{A} = \left| \vec{AB} \times \vec{AD} \right| = \sqrt{2}$.