

Q.1: Find the limit if it exist $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{\sqrt{x^2 + y^2 + 9} - 3}$.

Sol:
$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{\sqrt{x^2 + y^2 + 9} - 3} = \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{\sqrt{x^2 + y^2 + 9} - 3} \cdot \frac{\sqrt{x^2 + y^2 + 9} + 3}{\sqrt{x^2 + y^2 + 9} + 3}$$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{(x^2 + y^2) (\sqrt{x^2 + y^2 + 9} + 3)}{x^2 + y^2 + 9 - 9} = \lim_{(x,y) \rightarrow (0,0)} \sqrt{x^2 + y^2 + 9} + 3 = 6.$$

Q.2: Use implicit differentiation to find the partial derivatives $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ for $\cos(xyz) = 4x + 3y + z$.

Sol: Implicit differentiation to find $\frac{\partial z}{\partial x}$,

$$-\sin(xyz) \left(yz + xy \frac{\partial z}{\partial x} \right) = 4 + 0 + \frac{\partial z}{\partial x} \Rightarrow (-xy \sin(xyz) - 1) \frac{\partial z}{\partial x} = 4 + yz \sin(xyz)$$

$$\frac{\partial z}{\partial x} = \frac{4 + yz \sin(xyz)}{-xy \sin(xyz) - 1} = -\frac{4 + yz \sin(xyz)}{xy \sin(xyz) + 1}$$

We can also use chain rule to find these derivatives. Let $F(x, y, z) = \cos(xyz) - 4x - 3y - z = 0$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{-\sin(xyz) xz - 3}{-\sin(xyz) xy - 1} = -\frac{xz \sin(xyz) + 3}{xy \sin(xyz) + 1}.$$

Q.3: Find equation of tangent plane to the surface $z = \sqrt{7 - x^2 - 2y^2}$ at the point $(1, -1, 2)$. Using linearization $L(x, y)$ of $f(x, y) = \sqrt{7 - x^2 - 2y^2}$ at $(1, -1)$, approximate $f(0.99, -1.01)$.

Sol: $f_x = \frac{-2x}{2\sqrt{7 - x^2 - 2y^2}}$ and $f_y = \frac{-4y}{2\sqrt{7 - x^2 - 2y^2}}$

$$f_x(1, -1) = -\frac{1}{2} \text{ and } f_y(1, -1) = 1$$

$$\text{Equation of tangent plane is } z - 2 = -\frac{1}{2}(x - 1) + (y + 1) \Rightarrow z = -\frac{1}{2}x + y + \frac{7}{2}.$$

$$\text{Linearization of } f(x, y) \text{ at } (1, -1) \text{ is } L(x, y) = -\frac{1}{2}x + y + \frac{7}{2}.$$

$$f(0.99, -1.01) \approx -\frac{1}{2}(0.99) + (-1.01) + \frac{7}{2} = 1.995.$$