


**Q.1:** Estimate volume of the solid that lies below the surface  $z = 2x^2 + y^2$  and above the rectangle  $R = [0, 2] \times [0, 4]$ . Use a Riemann sum with  $m = 2$  and  $n = 4$  and choose upper right corner points.

**Sol:** Here  $f(x, y) = x^2 + 2y^2$ .



The lower left points are  $(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (2, 4)$  and  $\Delta A = \Delta x \Delta y = 1$ .


$V \approx \Delta A [f(1, 1) + f(1, 2) + f(1, 3) + f(1, 4) + f(2, 1) + f(2, 2) + f(2, 3) + f(2, 4)] = 3 + 6 + 11 + 18 + 9 + 12 + 17 + 24 = 100$ .

**Q.2:** Find exact volume of the solid that lies below the surface  $z = 2x^2 + y^2$  and above the rectangle  $R = [0, 2] \times [0, 4]$ .

**Sol:**  $V = \int_0^4 \int_0^2 (2x^2 + y^2) dx dy = \int_0^4 \left( \frac{2x^3}{3} + xy^2 \right) \Big|_0^2 dy = \int_0^4 \left( \frac{16}{3} + 2y^2 \right) dy = \frac{16}{3}y + \frac{2y^3}{3} \Big|_0^4 = \frac{64}{3} + \frac{128}{3} = \frac{192}{3} = 64$ .

**Q.3:** Evaluate the double integral  $\int_0^2 \int_{2y}^4 6e^{x^2} dx dy$ . (Hint: Change order of integration)

**Sol:** The region is  $0 \leq y \leq 2, 2y \leq x \leq 4$ .



$$\int_0^4 \int_0^{\frac{x}{2}} 6e^{x^2} dy dx = \int_0^4 3xe^{x^2} dx = \frac{3}{2} e^{x^2} \Big|_0^4 = \frac{3}{2} (e^{16} - 1)$$

**Q.4:** Use double integral to find the area enclosed by one loop of the graph of  $r = \cos(3\theta)$ .

**Sol:**  $\int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \int_0^{\cos(3\theta)} r dr d\theta = \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} r^2 \Big|_0^{\cos(3\theta)} d\theta = \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \cos^2(3\theta) d\theta = \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \frac{1 + \cos(6\theta)}{2} d\theta = \frac{1}{4} \left( \theta + \frac{\sin(6\theta)}{6} \right) \Big|_{-\frac{\pi}{6}}^{\frac{\pi}{6}} = \frac{\pi}{12}$ .