

2.5 Solutions by Substitutions

Objectives:

- To learn technique of transforming an ODE equation to a separable equation
- To know homogeneous functions and homogeneous ODE
- To recognize and solve homogeneous first order ODEs
- To recognize and solve Bernoulli equation.

A differential equation of the form

$$\frac{dy}{dx} = f(ax + by + c); \quad b \neq 0 \quad \{\text{what happens for } b = 0?\}$$

can always be reduced to a separable equation by the

substitution $u = ax + by + c$

where a , b , and c are constants.

Question 24/78: Solve the ODE $\frac{dy}{dx} = \frac{1-x-y}{x+y}$ by an appropriate substitution.

Question 29/78: Solve the IVP $\frac{dy}{dx} = \cos(x+y)$, $y(0) = \pi/4$ by an appropriate substitution.

Homogeneous Function: If $f(tx, ty) = t^n f(x, y)$ for some real number n , then $f(x, y)$ is called a **homogeneous function** of degree n .

For example, $f(x, y) = x^3 + y^3$ is a homogeneous function of degree 3, because $f(tx, ty) = t^3 f(x, y)$. The function $f(x, y) = x^3 + y^3 + 1$ is not homogeneous.

Homogeneous ODE: A first order DE in differential form

$$M(x, y)dx + N(x, y)dy = 0$$

is said to be **homogeneous** if both coefficient functions M and N are homogeneous functions of the **same** degree.

If M and N are homogeneous functions of degree n , then we can always write

$M(x, y) = x^n M(1, u)$ and $N(x, y) = x^n N(1, u)$ where $u = y/x$ and

$M(x, y) = y^n M(v, 1)$ and $N(x, y) = y^n N(v, 1)$ where $v = x/y$.

The homogeneous ODE is solved by substituting $y = ux$ or $x = vy$.

In particular, a homogeneous 1st order ODE is the one that can be written in the form

$$\frac{dy}{dx} = F\left(\frac{y}{x}\right) \quad (*)$$

Examples:

a) $\frac{dy}{dx} = \frac{2x^2 y}{3x^3 + y^3}$

b) $x \frac{dy}{dx} = y(\ln y - \ln x)$

Why?

Method of solution

Main idea: Given $\frac{dy}{dx} = F\left(\frac{y}{x}\right)$

The substitution $u = \frac{y}{x}$ (or $y = ux$) and hence $\frac{dy}{dx} = u + x \frac{du}{dx}$ transforms the ODE into a separable equation.

Steps for finding the solution:

Step 1: Check if **homogeneous**. Try to write in the form $\frac{dy}{dx} = F\left(\frac{y}{x}\right)$ (*) {If homogeneous, proceed as follows}

Step 2: Substitute $u = \frac{y}{x}$ (or $y = ux$) and $\frac{dy}{dx} = u + x \frac{du}{dx}$ in (*).

This transforms it into separable equation.

Step 3: Solve the **separable equation**.

Step 4: Use the substitution $u = \frac{y}{x}$ to get the **general solution** in the variables **y and x**.

Step 6: In case of **IVP**, find the **particular solution**.

Question 9/78: Solve the ODE $-ydx + (x + \sqrt{xy})dy = 0$ by appropriate substitution.

Question 14/78: Solve the IVP $ydx + x(\ln x - \ln y - 1)dy = 0$, $y(1) = e$ by appropriate substitution.

Bernoulli Equation: A first order ODE of the form

$$\frac{dy}{dx} + P(x) \cdot y = Q(x) \cdot y^n \quad (*)$$

is called ***Bernoulli equation***.

Note that for $n = 0$ and $n = 1$, equation (*) is linear.

Method of solution

The substitution $u = y^{1-n}$ to transforms the Bernoulli equation into a linear equation as follows:

- $\frac{du}{dx} = (1-n)y^{-n} \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{y^n}{(1-n)} \frac{du}{dx}$
- $\frac{du}{dx} + (1-n)P(x)u = (1-n)Q(x)$

Question 20/78: Solve ODE $3(1+t^2) \frac{dy}{dt} = 2ty(y^3 - 1)$ by an

appropriate substitution.

Question 22/78: Solve IVP $y^{1/2} \frac{dy}{dx} + y^{3/2} = 1$, $y(0) = 4$ by an

appropriate substitution.