



# The Existence of Nonlinear Inequalities

Q. H. ANSARI\*

Department of Mathematics, A.M.U.

Aligarh 202 002, India

NGAI-CHING WONG AND JEN-CHIH YAO

Department of Applied Mathematics, National Sun Yat-sen University

Kaohsiung, Taiwan, R.O.C.

(Received and accepted May 1998)

**Abstract**—The object of this paper is to establish an existence result for nonlinear inequalities for not necessarily pseudomonotone maps. As a consequence of our result, we give some existence results for variational and variational-like inequalities. © 1999 Elsevier Science Ltd. All rights reserved.

**Keywords**—Nonlinear inequalities, Variational inequality problems, Variational-like inequality problems, Fixed points.

## 1. INTRODUCTION AND PRELIMINARIES

Let  $X$  be a real locally convex space with its dual  $X^*$  and  $K$  be a nonempty convex subset of  $X$ . Let  $\varphi : K \times K \rightarrow \mathbb{R}$  be a bifunction and  $f \in X^*$ . Gwinner [1] considered the problem of finding  $u_0$  in  $K$  such that

$$\varphi(u_0, v) \geq \langle f, v - u_0 \rangle, \quad \text{for all } v \text{ in } K, \quad (1)$$

where  $\langle \cdot, \cdot \rangle$  denotes the pairing between  $X^*$  and  $X$ . Equation (1) is called the *nonlinear inequality*.

Such types of nonlinear inequalities model some equilibrium problems drawn from operations research, as well as some unilateral boundary value problems stemming from mathematical physics. The existence theory and abstract stability analysis of (1) have been investigated by Gwinner [1] in the setting of reflexive Banach spaces.

When  $\varphi(u, v) = \langle T(u), v - u \rangle$ , where  $A : K \rightarrow X^*$  is a map, (1) reduces to the *classical variational inequality problem* introduced by Lions and Stampacchia [2], that is, to find a  $u_0$  in  $K$  such that

$$\langle T(u_0), v - u_0 \rangle \geq \langle f, v - u_0 \rangle, \quad \text{for all } v \text{ in } K. \quad (2)$$

Even more specifically, when  $\varphi(u, v) = \langle T(u), v - u \rangle$  and  $f \equiv 0$ , (1) becomes the *original variational inequality problem* (see, e.g., [3,4]), that is, to find a  $u_0$  in  $K$  such that

$$\langle T(u_0), v - u_0 \rangle \geq 0, \quad \text{for all } v \text{ in } K. \quad (3)$$

Many authors have studied this type of problem in the context of reflexive Banach spaces (see, for example, [3,5–7]). Watson [8] established the existence of solutions to problem (3) in

---

This research is partially supported by National Science Council of the Republic of China.

\*Presently visiting the Department of Applied Mathematics, National Sun Yat-sen University, Kaohsiung, Taiwan, R.O.C.

the setting of not necessarily reflexive Banach spaces and pseudomonotone and hemicontinuous maps. His assumptions are weaker than those needed in [9].

When  $\varphi(u, v) = \langle T(u), \eta(v, u) \rangle$ , and  $f \equiv 0$ , where  $\eta : K \times K \rightarrow X$  is a map, (1) is equivalent to find a  $u_0$  in  $K$  such that

$$\langle T(u_0), \eta(v, u_0) \rangle \geq 0, \quad \text{for all } v \text{ in } K. \quad (4)$$

Inequality (4) is known as a *variational-like inequality* which has many applications in operations research, optimization, and mathematical programming. For further details, we refer to [10–13].

In the case  $f \equiv 0$  and  $\varphi(u, u) = 0$ , for all  $u$  in  $K$ , (1) reduces to the *equilibrium problem* considered in [14–16], that is, to find a  $u_0$  in  $K$  such that

$$\varphi(u_0, v) \geq 0, \quad \text{for all } v \text{ in } K. \quad (5)$$

Apparently, our nonlinear inequality (1) contains all above variational inequalities and equilibrium problems as special cases.

The main object of this paper is to establish an existence result for the nonlinear inequality (1) for general maps without any pseudomonotone assumption. We shall employ Fan-Browder [17,18] type fixed-point theorems due to [19]. As a consequence of our result, we shall derive some existence results for the variational inequality (3) and the variational-like inequality (4) without any kind of monotonicity assumption.

The following fixed-point theorem will be used in this paper. We denote by  $2^Y$  the family of all nonempty subsets of a set  $Y$ .

**THEOREM 1.** (See [19].) *Let  $K$  be a nonempty convex subset of a Hausdorff topological vector space  $X$ . Let  $Q : K \rightarrow 2^K$  be a multifunction such that we have the following.*

- (a) *For each  $x$  in  $K$ ,  $Q(x)$  is a nonempty convex subset of  $K$ .*
- (b) *For each  $y$  in  $K$ ,  $Q^{-1}(y)$  contains relatively open subset  $O_y$  of  $K$  ( $O_y$  may be empty for some  $y$  in  $K$ ) such that  $\bigcup_{y \in K} O_y = K$ .*
- (c)  *$K$  contains a nonempty subset  $D_0$  which is contained in a compact convex subset  $D_1$  of  $K$  such that the set  $D = \bigcap_{x \in D_0} O_x^c$  is either empty or compact, where  $O_x^c$  denotes the complements of  $O_x$  in  $K$ .*

Then there exists a point  $x_0$  in  $K$  such that  $x_0 \in Q(x_0)$ .

## 2. MAIN RESULTS

We now prove the main result of this paper.

**THEOREM 2.** *Let  $K$  be a nonempty convex subset of a Hausdorff topological vector space  $X$  (over the real field). Let  $f$  be a nonzero continuous linear functional on  $X$ . Let  $\varphi : K \times K \rightarrow \mathbb{R}$  be a bifunction vanishing on the diagonal, i.e.,  $\varphi(u, u) = 0$ , for all  $u$  in  $K$ , and satisfying the following conditions.*

1.  *$\varphi$  is convex in the second variable.*
2.  *$\liminf_{u \rightarrow u^*} \varphi(u, v) \leq \varphi(u^*, v)$ , for all  $v$  in  $K$  whenever  $u \rightarrow u^*$  in  $K$ .*
3. *There is a compact convex subset  $D_1$  of  $K$  such that for each  $u$  in  $K \setminus D_1$ , there is a  $v$  in  $D_1$  with*

$$\varphi(u, v) < \langle f, v - u \rangle.$$

Then the nonlinear inequality (1) has a solution in  $K$ .

**PROOF.** We define  $A(v) = \{u \in K : \varphi(u, v) \geq \langle f, v - u \rangle\}$  for each  $v$  in  $K$ . Then the solution set of (1) is  $S = \bigcap_{v \in K} A(v)$ . We note that for each  $v$  in  $K$ ,  $A(v)$  is closed. Indeed, let  $\{u_\lambda\}_{\lambda \in \Lambda}$  be

a net in  $A(v)$  such that  $u_\lambda \rightarrow u$  in  $K$ . Then for all  $f$  in  $X^*$ , we have  $\langle f, u_\lambda - v \rangle \rightarrow \langle f, u - v \rangle$ . Since  $u_\lambda \in A(v)$  and  $\liminf_{u_\lambda \rightarrow u} \varphi(u_\lambda, v) \leq \varphi(u, v)$ , for all  $v$  in  $K$ , we have

$$\varphi(u, v) \geq \liminf_{u_\lambda \rightarrow u} \varphi(u_\lambda, v) \geq \liminf_{u_\lambda \rightarrow u} \langle f, u_\lambda - v \rangle = \langle f, u - v \rangle.$$

Hence,  $u \in A(v)$ . So for all  $v$  in  $K$ ,  $A(v)$  is closed.

Now we shall prove that the solution set  $S$  is nonempty. Assume contrary that  $S = \emptyset$ . Then for each  $u$  in  $K$ , the set

$$B(u) = \{v \in K : u \notin A(v)\} = \{v \in K : \varphi(u, v) < \langle f, v - u \rangle\} \neq \emptyset.$$

Since  $\varphi$  is convex in the second variable, we have for each  $u$  in  $K$ ,  $B(u)$  is convex. Thus,  $B : K \rightarrow 2^K$  defines a multifunction such that for each  $u$  in  $K$ ,  $B(u)$  is nonempty and convex. Now for each  $v$  in  $K$ , the set

$$\begin{aligned} B^{-1}(v) &= \{u \in K : v \in B(u)\} \\ &= \{u \in K : \varphi(u, v) < \langle f, v - u \rangle\} \\ &= \{u \in K : \varphi(u, v) \geq \langle f, v - u \rangle\}^c \\ &= [A(v)]^c \\ &= O_v \end{aligned}$$

is open in  $K$ . We claim that  $\bigcup_{v \in K} O_v = \bigcup_{v \in K} B^{-1}(v) = K$ . To see this, let  $u \in K$ . As  $B(u) \neq \emptyset$ , we can choose a  $v$  from  $B(u)$ . Hence,  $u \in B^{-1}(v) = O_v$ .

From the last condition of the theorem, for each  $u$  in  $K \setminus D_1$ , there is a  $v$  in  $D_1$  with  $\varphi(u, v) < \langle f, v - u \rangle$ , that is,  $u \notin A(v)$ . This implies that  $D = \bigcap_{v \in D_1} O_v^c = \bigcap_{v \in D_1} A(v) \subset D_1$ . Since for each  $v$  in  $K$ ,  $A(v)$  is closed,  $D$  is a closed subset of the compact set  $D_1$ , and hence,  $D$  is compact. Thus, the multifunction  $B : K \rightarrow 2^K$  satisfies all the conditions of Theorem 1, so there exists a point  $u_0$  in  $K$  such that  $u_0 \in B(u_0)$ , that is,  $0 = \varphi(u_0, u_0) < \langle f, u_0 - u_0 \rangle = 0$ , which is a contradiction. Hence, the solution set  $S$  is nonempty. Therefore, the nonlinear inequality (1) has a solution in  $K$ . ■

In case  $K$  is compact, the last condition in Theorem 2 is automatically satisfied (in particular,  $X$  is a Banach space), since we can set  $D_1 = K$ . Meanwhile, if a Hausdorff locally convex space  $X$  is barreled, then every weak\* closed and bounded subset  $K$  of  $X^*$  is weak\* compact (see, for example, [20, p. 141]).

**COROLLARY 1.** *Let  $X$  be a Hausdorff locally convex space with dual  $X^*$ . Let  $K$  be a weak\* compact convex subset of  $X^*$  and  $f \in X$ . Let  $\varphi : K \times K \rightarrow \mathbb{R}$  be a bifunction vanishing on the diagonal, convex in the second variable, and satisfying the condition that  $\liminf_{u \rightarrow u^*} \varphi(u, v) \leq \varphi(u^*, v)$ , for all  $v \in K$ , whenever  $u \rightarrow u^*$  in  $K$ . Then the nonlinear inequality (1) has a solution in  $K$ .*

**COROLLARY 2.** *Let  $K$  be a weak\* compact convex subset of the dual space  $X^*$  of a Hausdorff locally convex space  $X$ . Let  $\eta : K \times K \rightarrow X^*$  be a bifunction vanishing on the diagonal. Let  $T$  be a map from  $K$  into  $X$  such that  $h(v) := \langle \eta(v, u), T(u) \rangle$  is convex in  $v$ , for each fixed  $u$  in  $K$ , and  $\liminf_{u \rightarrow u^*} \langle \eta(v, u), T(u) \rangle \leq \langle \eta(v, u^*), T(u^*) \rangle$  for each  $v$  in  $K$ , whenever  $u \rightarrow u^*$  in the weak\* topology of  $K$ . Then the variational-like inequality (4) has a solution in  $K$ .*

**COROLLARY 3.** *Let  $K$  be a closed convex subset of the dual space  $X^*$  of a Hausdorff locally convex barreled space  $X$ . Let  $T$  be a map from  $K$  into  $X$  such that  $\liminf_{u \rightarrow u^*} \langle v - u, T(u) \rangle \leq \langle v - u^*, T(u^*) \rangle$ , for each  $v$  in  $K$ , whenever  $u \rightarrow u^*$  in the weak\* topology of  $K$ . Further assume that there is a compact subset  $D$  of  $K$  such that for each  $u$  in  $K \setminus D$ , there is a  $v$  in  $D$  such that  $\langle v - u, T(u) \rangle < 0$ . Then the variational inequality problem (3) has a solution in  $K$ .*

PROOF. We note that the convex hull of a totally bounded subset of any locally convex space is totally bounded as well. On the other hand, the dual space  $X^*$  of the Hausdorff locally convex barreled space  $X$  is quasi-complete in its weak\* topology, that is, closed and bounded subsets of  $X^*$  are complete (in fact, weak\* compact). As a result, the convex hull  $D_1$  of the weak\* compact subset  $D$  of  $X^*$  is still weak\* compact. Consequently, Theorem 2 applies. ■

COROLLARY 4. *Let  $X$  be a Hausdorff locally convex space with dual  $X^*$ . Let  $T$  be a map from  $X^*$  into  $X$  such that we have the following.*

1.  $\liminf_{u \rightarrow u^*} \langle v - u, T(u) \rangle \leq \langle v - u^*, T(u^*) \rangle$ , for each  $v$  in  $X^*$ , whenever  $u \rightarrow u^*$  in the weak\* topology of  $X^*$ .
2. There exists a weak\* compact convex subset  $D_1$  of  $X^*$  such that for each  $u$  not in  $D_1$  there is a  $v$  in  $D_1$  with  $\langle v - u, T(u) \rangle < 0$ .

Then  $T$  has a zero  $u_0$  in  $X^*$ , i.e.  $T(u_0) = 0$ . In case  $X$  is barreled, the convexity assumption on  $D_1$  can be dropped.

## REFERENCES

1. J. Gwinner, Stability of monotone variational inequalities with various applications, In *Variational Inequalities and Network Equilibrium Problems*, (Edited by F. Giannessi and A. Maugeri), pp. 123–142, Plenum Press, (1995).
2. J.L. Lions and G. Stampacchia, Variational inequalities, *Comm. Pure Appl. Math.* **20**, 493–519 (1967).
3. F.E. Browder, Nonlinear elliptic boundary value problems, *Trans. Amer. Math. Soc.* **117**, 530–550 (1965).
4. P. Hartman and G. Stampacchia, On some nonlinear elliptic differential functional equations, *Acta Math.* **115**, 153–188 (1966).
5. D. Pascali and S. Sburian, *Nonlinear Mappings of Monotone Type*, Romania, (1978).
6. D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, Academic Press, New York, (1980).
7. J.S. Guo and J.C. Yao, Variational inequalities with nonmonotone operators, *J. Optim. Theory Appl.* **80**, 63–74 (1994).
8. P.J. Watson, Variational inequalities in nonreflexive Banach spaces, *Appl. Math. Lett.* **10** (2), 45–48 (1997).
9. S. Chang, B.S. Lee and Y. Chen, Variational inequalities for monotone operators in nonreflexive Banach spaces, *Appl. Math. Lett.* **8** (6), 29–34 (1995).
10. J. Parida, M. Sahoo and A. Kumar, A variational-like inequality problem, *Bull. Austral. Math. Soc.* **39**, 225–231 (1989).
11. N.D. Dien, Some remarks on variational-like and quasivariational-like inequalities, *Bull. Austral. Math. Soc.* **46**, 335–342 (1992).
12. X.Q. Yang and G.Y. Chen, A class of nonconvex functions and pre-variational inequalities, *J. Math. Anal. Appl.* **169**, 359–373 (1992).
13. A.H. Siddiqi, A. Khaiq and Q.H. Ansari, On variational-like inequalities, *Ann. Sci. Math. Quebec* **18**, 39–48 (1994).
14. M. Bianchi and S. Schaible, Generalized monotone bifunctions and equilibrium problems, *J. Optim. Theory Appl.* **90**, 31–43 (1996).
15. E. Blum and W. Oettli, From optimization and variational inequalities to equilibrium problems, *The Math. Student* **63**, 123–145 (1993).
16. E. Blum and W. Oettli, Variational principle for equilibrium problems, In *Parametric Optimization and Related Topics III*, (Edited by J. Guddat *et al.*), pp. 79–88, Peter Lang, Frankfurt am Main, (1993).
17. F.E. Browder, The fixed point theory of multivalued mappings in topological vector spaces, *Math. Ann.* **177**, 283–301 (1968).
18. K. Fan, A generalization of Tychonoff's fixed point theorem, *Math. Ann.* **142**, 305–310 (1961).
19. E. Tarafdar, A fixed point theorem equivalent to Fan-Knaster-Kuratowski-Mazurkiewicz's theorem, *J. Math. Anal. Appl.* **128**, 475–479 (1987).
20. H. H. Schaefer, *Topological Vector Spaces*, Fifth edition, Springer-Verlag, New York, (1986).