

King Fahd University of Petroleum and Minerals
Prep-Year Math Program

Code: Master

Prep-Year Math II
FINAL EXAM
Semester I, 2003-04
Saturday, January 10, 2004
Net Time Allowed: 130 minutes

Code: Master

Student's Name:

ID #:

Section #:

Important Instructions:

1. All types of CALCULATORS, PAGERS, and OR MOBILES ARE NOT ALLOWED to be with you during the examination.
2. Use an HB $2\frac{1}{2}$ pencil.
3. Use a good eraser. Do not use the eraser attached to the pencil.
4. Write your name, ID number and Mathematics Section on the examination paper and in the upper left corner of the answer sheet.
5. When bubbling your ID number and Math Section number, be sure that bubbles match with the number that you write.
6. The test Code Number is already typed and bubbled in your answer sheet. Make sure that it is the same as that printed on your question paper.
7. When bubbling, make sure that the bubbled space is fully covered.
8. When erasing a bubble, make sure that you do not leave any trace of penciling.
9. Check that the exam paper has 25 questions.

1. The coordinates of the **focus** of the parabola $8x - y^2 + 4y - 12 = 0$ are

(a) (3, 2)

(b) (-3, 2)

(c) (3, -2)

(d) (3, 4)

(e) (2, 4)

2. If $y = -3 \sin x + 3\sqrt{3} \cos x$ is rewritten as $y = k \sin(x + \alpha)$, $0 \leq \alpha < 360^\circ$, then the values of k and α are given by

(a) $k = 6$, $\alpha = 120^\circ$

(b) $k = 18$, $\alpha = 210^\circ$

(c) $k = 6$, $\alpha = 240^\circ$

(d) $k = 9$, $\alpha = 300^\circ$

(e) $k = 6$, $\alpha = 330^\circ$

3. If (a, b) is the solution of the system

$$\begin{cases} 6x + 7y = -4 \\ 2x + 5y = 4 \end{cases}$$

then $a + b$ is equal to

(a) -1

(b) 3

(c) -2

(d) $\frac{3}{2}$

(e) $-\frac{1}{2}$

4. If $3^{2+x} = 6^x$, then $x =$

(a) $\log_2 9$

(b) $\log_3 2$

(c) $\log_2 3$

(d) $\log_9 6$

(e) $\log_3 \frac{3}{2}$

5. Given the vectors $\mathbf{v} = \langle 6, 7 \rangle$ and $\mathbf{w} = \langle 3, 4 \rangle$, the value of $\text{Proj}_{\mathbf{w}} \mathbf{v}$ is equal to

(a) -2

(b) $\frac{13}{5}$

(c) $-\frac{21}{5}$

(d) $\frac{41}{5}$

(e) -6

6. Let A , B , and C be any $n \times n$ matrices and let O be the $n \times n$ zero matrix. Which one of the following statements is **FALSE**?

(a) $AB - BA = O$

(b) $A(BC) = (AB)C$

(c) $OA = AO = O$

(d) $A(B + C) = AB + AC$

(e) $A^2 + B^2$ is an $n \times n$ matrix

7. If $A = \begin{bmatrix} 1 & 0 & 4 \\ 2 & 1 & 0 \\ 1 & 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 & 2 \\ m & 1 & -4 \\ \frac{1}{2} & 0 & k - \frac{1}{2} \end{bmatrix}$ are inverses of each other, then

- (a) $k = 0$ and $m = 2$
- (b) $k = -1$ and $m = 0$
- (c) $k = 1$ and $m = 2$
- (d) $k = 2$ and $m = 0$
- (e) $k = 3$ and $m = -1$
8. The equation of the **directrix** of the parabola that has vertex $(-4, 1)$, has its axis of symmetry parallel to the y axis, and passes through the point $(-2, 2)$ is given by

- (a) $y = 0$
- (b) $y = -1$
- (c) $x = 2$
- (d) $y = -2$
- (e) $x = 2$

9. Which one of the following statements is **FALSE** about the graph of

$$f(x) = \frac{e^{-x} + e^x}{2}?$$

- (a) the graph is increasing on $(-\infty, \infty)$
- (b) $(0, 1)$ is the lowest point on the graph
- (c) the graph is symmetric about the y -axis
- (d) the graph has only one y -intercept
- (e) the graph has no x -intercept
10. If A and B are 2×2 matrices with $|A| = 5$, $|B| = 2$, then $|3AB| + 2|B^{-1}|$ is equal to

- (a) 91
- (b) 63
- (c) 84
- (d) 103
- (e) 37

11. The expression

$$\log\left(y^{3/2}z^2\right) - 3\log(x\sqrt{y}) + 2\log\frac{x}{z}$$

simplifies to

(a) $-\log x$

(b) $\log\frac{y^2}{x}$

(c) $\frac{1}{2}\log(xy)$

(d) $-\log\left(\frac{y^3}{xz}\right)$

(e) $\log\left(\frac{y}{xz^3}\right)$

12. The number of solutions in the solution set of the equation

$$2\sin^2 x - \sin 2x = 0, \quad \text{where } 0 \leq x < \frac{5\pi}{2},$$

is equal to

(a) 6

(b) 5

(c) 4

(d) 3

(e) 2

13. The exact value of $\cos 157.5^\circ$ is equal to

(a) $-\frac{\sqrt{2 + \sqrt{2}}}{2}$

(b) $-\frac{\sqrt{2 - \sqrt{2}}}{2}$

(c) $\frac{\sqrt{2 + \sqrt{2}}}{2}$

(d) $\frac{\sqrt{2 - \sqrt{2}}}{2}$

(e) $2 + \sqrt{2}$

14. One of the foci of the ellipse $\frac{(x - 3)^2}{16} + \frac{(y + 2)^2}{25} = 1$ is at

(a) $(3, -5)$

(b) $(1, -2)$

(c) $(3, -9)$

(d) $(7, -2)$

(e) $(3, -1)$

15. Given the vectors $\mathbf{u} = 10\mathbf{i} - 8\mathbf{j}$ and $\mathbf{v} = 12\mathbf{i} - 6\mathbf{j}$, the direction angle of the vector $\frac{1}{2}\mathbf{u} - \frac{1}{6}\mathbf{v}$, in radians, is equal to

(a) $\frac{7\pi}{4}$

(b) $\frac{5\pi}{4}$

(c) $\frac{3\pi}{4}$

(d) $\frac{\pi}{4}$

(e) $\frac{5\pi}{3}$

16. The solution set of the equation

$$\log(-6 - 7x) = \log(3 + 2x) + \log(4 + 3x)$$

contains

(a) one negative integer only

(b) two negative integers

(c) one positive integer only

(d) one negative and one positive integers

(e) no real numbers

17. The equation of the asymptote with **positive slope** of the hyperbola $4x^2 - 9y^2 - 16x + 5y - 29 = 0$ is

(a) $2x - 3y + 5 = 0$

(b) $4x - 9y + 19 = 0$

(c) $2x - 6y + 10 = 0$

(d) $4x - 3y + 1 = 0$

(e) $3x - 4y + 6 = 0$

18. If $0 < x < \pi$, then $\csc x - \cot x$ simplifies to

(a) $\tan \frac{x}{2}$

(b) $\csc \frac{x}{2}$

(c) $\cos \frac{x}{2}$

(d) $\sin \frac{x}{2}$

(e) $\sec \frac{x}{2}$

19. Let x be a nonzero real number and $A = \begin{bmatrix} x & x^4 & x \\ 0 & x^3 & 1 \\ x^5 & 2x^8 & -3x^5 \end{bmatrix}$. Then the determinant of A is equal to

- (a) $-5x^9$
(b) $-3x^5$
(c) $-3x^9$
(d) $6(x^5 - x^3)$
(e) $4x^5$

20. The equation in standard form of the hyperbola with vertices $(2, 3)$ and $(-2, 3)$, and eccentricity $\frac{5}{2}$ is given by

- (a) $\frac{x^2}{4} - \frac{(y-3)^2}{21} = 1$
(b) $\frac{(x-2)^2}{4} - \frac{(y-6)^2}{21} = 1$
(c) $\frac{x^2}{16} - \frac{(y-3)^2}{25} = 1$
(d) $\frac{(x-4)^2}{4} - \frac{(y-3)^2}{21} = 1$
(e) $\frac{x^2}{4} - \frac{(y-3)^2}{36} = 1$

21. If a wheel with a **diameter** of 16 centimeters is rotating at 10 radians per minute, then the linear speed of a point on the edge of the wheel is equal to

(a) $\frac{4}{3}$ centimeters/second

(b) $\frac{5}{3}$ centimeters/second

(c) $\frac{3}{2}$ centimeters/second

(d) $\frac{2}{3}$ centimeters/second

(e) $\frac{3}{10}$ centimeters/second

22. $\sin 675^\circ + \cos(-405^\circ) + \tan \frac{8\pi}{3} =$

(a) $-\sqrt{3}$

(b) $\sqrt{2} + \sqrt{3}$

(c) $3\sqrt{2} - \sqrt{3}$

(d) $-\sqrt{2} + 2\sqrt{3}$

(e) $2\sqrt{3}$

23. A linear system written in a matrix form

$$\begin{bmatrix} -2 & 6 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ 4 \end{bmatrix}$$

is

- (a) dependent
- (b) inconsistent
- (c) independent
- (d) consistent with only two solutions
- (e) consistent with only three solutions

24. The length of the major axis of the ellipse with center $(-4, 1)$, minor axis parallel to the y -axis and of length 8, and passing through $(0, 3)$ is equal to

- (a) $\frac{16\sqrt{3}}{3}$
- (b) $32\sqrt{3}$
- (c) $\frac{9}{2}\sqrt{3}$
- (d) $\frac{3\sqrt{3}}{2}$
- (e) $\frac{25\sqrt{3}}{3}$

25. Let $A = \begin{bmatrix} -2 & 1 & 0 \\ 0 & -1 & 1 \\ 1 & 0 & 3 \end{bmatrix}$ and I be the 3×3 identity matrix. Then the element in the second row and third column of the matrix $3A^2 - 2I$ is

- (a) 6
- (b) 4
- (c) 7
- (d) 5
- (e) 10

26. The graph of the function

$$y = 3 \cos 3(x - \pi), \quad \pi \leq x \leq \frac{5\pi}{3},$$

lies above the x -axis on the interval

- (a) $\left[\pi, \frac{7\pi}{6}\right) \cup \left(\frac{3\pi}{2}, \frac{5\pi}{3}\right]$
- (b) $\left(\frac{7\pi}{6}, \frac{5\pi}{3}\right)$
- (c) $\left(\frac{7\pi}{6}, \frac{4\pi}{3}\right) \cup \left(\frac{3\pi}{2}, \frac{5\pi}{3}\right)$
- (d) $\left[\pi, \frac{3\pi}{2}\right)$
- (e) $\left[\pi, \frac{7\pi}{6}\right) \cup \left[\frac{4\pi}{3}, \frac{3\pi}{2}\right)$

27. If $A = \begin{bmatrix} 2 & 1 & 4 \\ 1 & 2 & 3 \\ 1 & 5 & 1 \end{bmatrix}$, then the sum of the cofactors of the elements of the **third** row of the matrix A is

(a) -4

(b) 11

(c) 6

(d) -7

(e) 8

28. The number of real solutions of the nonlinear system

$$\begin{cases} x^2 + y^2 = 1 \\ \frac{(x+1)^2}{4} - y^2 = 1 \end{cases}$$

is equal to

(a) 1

(b) 2

(c) 3

(d) 4

(e) 0

29. The exact value of $\tan \left[\sin^{-1} \left(\frac{3}{5} \right) + \cos^{-1} \left(-\frac{5}{13} \right) \right]$ is equal to

(a) $-\frac{33}{56}$

(b) $-\frac{17}{65}$

(c) $\frac{56}{33}$

(d) $-\frac{15}{56}$

(e) $\frac{41}{65}$

30. The graph of $y = 6 \sec \left(\pi x + \frac{\pi}{4} \right)$, $\frac{1}{4} < x < \frac{9}{4}$, is increasing on the interval

(a) $\left(\frac{1}{4}, \frac{3}{4} \right) \cup \left(\frac{7}{4}, \frac{9}{4} \right)$

(b) $\left(\frac{3}{4}, \frac{5}{4} \right) \cup \left(\frac{5}{4}, \frac{7}{4} \right)$

(c) $\left(\frac{1}{4}, \frac{5}{4} \right)$

(d) $\left(\frac{5}{4}, \frac{9}{4} \right)$

(e) $\left(\frac{1}{4}, \frac{5}{4} \right) \cup \left(\frac{5}{4}, \frac{9}{4} \right)$

31. The sum of all solutions of the equation

$$2 \sin x \cos x - \sqrt{3} \sin x + 2\sqrt{2} \cos x + \sqrt{6} = 0, \quad \text{where } 0 \leq x < 2\pi$$

is equal to

(a) 2π

(b) $\frac{9\pi}{4}$

(c) $\frac{5\pi}{6}$

(d) $\frac{11\pi}{4}$

(e) $\frac{17\pi}{6}$

32. Let $A = \begin{bmatrix} 1 & 0 & 1 \\ 2 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}$. Then the sum of the elements of the second row of A^{-1} is equal to

(a) $-\frac{3}{2}$

(b) -4

(c) $\frac{5}{2}$

(d) $\frac{7}{2}$

(e) $-\frac{1}{2}$

33. Which one of the following statements is **TRUE**?

(a) $\tan(\tan^{-1} 100) = 100$

(b) $\sin\left(\sin^{-1} \frac{3}{2}\right) = \frac{3}{2}$

(c) $\cos\left(\cos^{-1}\left(-\frac{3}{2}\right)\right) = \frac{3}{2}$

(d) $\sin\left(\sin\left(-\frac{3\pi}{4}\right)\right) = \frac{\pi}{4}$

(e) $\tan^{-1}\left(\tan\left(-\frac{\pi}{4}\right)\right) = \frac{3\pi}{4}$

34. If the **echelon** form of the augmented matrix for the linear system

$$\begin{cases} x - 3y + z = 8 \\ 2x - 5y - 3z = 6 \\ x - 6y + 7z = -7 \end{cases}$$

is $\left[\begin{array}{ccc|c} 1 & -3 & 1 & 8 \\ 0 & 1 & m & n \\ 0 & 0 & 1 & p \end{array} \right]$, then

(a) $m = -5$, $n = -10$, and $p = 5$

(b) $m = 3$, $n = -6$, and $p = -3$

(c) $m = -5$, $n = 10$, and $p = -3$

(d) $m = -2$, $n = 7$, and $p = -1$

(e) $m = -3$, $n = 6$, and $p = -2$

35) If $\tan \frac{\theta}{2} = -\frac{4}{3}$, $\pi < \theta < \frac{3\pi}{2}$, then $\csc \theta =$

(a) $-\frac{25}{24}$

(b) $\frac{5}{3}$

(c) $-\frac{25}{7}$

(d) $\frac{10}{3}$

(e) $-\frac{25}{12}$