

1. Which one of the following statements is TRUE about the graph of

$$f(x) = 1 - \left(\frac{1}{2}\right)^{x+2}?$$

- (a) The graph of  $f$  decreases on the interval  $(-\infty, \infty)$ .  
 (b) The graph of  $f$  increases on the interval  $(-\infty, \infty)$ .  
 (c) The graph of  $f$  decreases on the interval  $(-\infty, 2)$  and increases on the interval  $(2, \infty)$ .  
 (d) The line  $y = 0$  is a horizontal asymptote for the graph of  $f$ .  
 (e) The graph of  $f$  lies above the line  $y = 1$ .

see #37, 38 p. 317

2. Let  $W$  be the wrapping function. If  $W\left(-\frac{25\pi}{6}\right) = (x, y)$ , then  $x - y =$

- (a)  $\frac{1}{2}(\sqrt{3} + 1)$   
 (b)  $-\frac{1}{2}(\sqrt{3} + 1)$   
 (c)  $\frac{1}{2}(-\sqrt{3} + 1)$   
 (d)  $\frac{1}{2}(\sqrt{3} + \sqrt{2})$   
 (e)  $\frac{1}{2}(\sqrt{3} - \sqrt{2})$

see example 2 p. 425, and # 1 to 11 p. 43

3. The adjacent figure represents a part of the graph of


(a)  $y = -2 \cos \frac{\pi}{2}x$

(b)  $y = -2 \sin \frac{\pi}{2}x$

(c)  $y = 2 \cos \frac{\pi}{2}x$

(d)  $y = 2 \sin \frac{\pi}{2}x$

(e)  $y = 2 \sin \left( \frac{\pi}{2}x - 1 \right)$



See example 5 p.440, Problem #39 p.441  
and many others

4. If  $f(x) = -3 \cos x$  and  $g(x) = \sin x + \tan x$ , then

(a)  $f(x)$  is an even function and  $g(x)$  is an odd function

(b) both  $f(x)$  and  $g(x)$  are even functions

(c)  $f(x)$  is an odd function and  $g(x)$  is an even function

(d)  $f(x)$  is an even function and  $g(x)$  is neither an odd nor an even function

(e) both  $f(x)$  and  $g(x)$  are odd functions

See example 3 p.427 and #33 to 40 p.432

5. The exact value of  $\log_{0.3} \frac{1000}{27}$  is equal to

(a) 3

(b)  $\frac{1}{3}$

(c)  $\frac{1}{9}$

(d) 0.9

(e) -0.9

Similar to #25 to 28 p.327

6. The period  $P$ , amplitude  $A$ , and phase shift  $S$  of the graph of  $y = -8 \sin\left(\frac{3\pi}{2}x - \frac{\pi}{2}\right)$  are:

(a)  $P = \frac{4}{3}$ ,  $A = 8$ ,  $S = \frac{1}{3}$

(b)  $P = 2\pi$ ,  $A = 8$ ,  $S = -\frac{1}{3}$

(c)  $P = \frac{4}{3}$ ,  $A = 8$ ,  $S = \frac{1}{3}$

(d)  $P = \frac{1}{3}$ ,  $A = 8$ ,  $S = \frac{\pi}{3}$

(e)  $P = \frac{4}{3}$ ,  $A = -8$ ,  $S = -3$

See example 6 p.453, problems #39 to 44 p.456

7. Given  $\tan \alpha = -\frac{4}{3}$  where  $\alpha$  terminates in quadrant II and  $\cos \beta = \frac{12}{13}$  where  $\beta$  terminates in quadrant IV, then  $\csc(\alpha + \beta) =$

(a)  $\frac{65}{63}$

(b)  $\frac{65}{33}$

(c)  $-\frac{63}{65}$

(d)  $-\frac{65}{33}$

(e)  $-\frac{65}{63}$

Similar to example 2 p-482 and Problems 31 to 42 p-485

8. The adjacent figure represents the graph of


(a)  $y = \log_{\frac{1}{3}}(x - 2)$

(b)  $y = 2 - \log_2 x$

(c)  $y = \log_{\frac{1}{2}}(x - 3)$

(d)  $y = \log_2(x - 2)$

(e)  $y = \log_{\frac{1}{2}} x - 2$



Similar to example 7 p-325

and # 51, 52 p-328

9. If a wheel rotates at 750 revolutions per minute, then the angular speed of the wheel in radians per second is equal to

- (a)  $25\pi$   
(b)  $1500\pi$   
(c) 750  
(d)  $750\pi$   
(e)  $\frac{375}{\pi}$

See example 7 p.397 and #69, 70

10. The arc length that subtends a central angle  $50^\circ$  in a circle of diameter 10 cm is equal to

- (a)  $\frac{25\pi}{18}$  cm  
(b) 250 cm  
(c) 500 cm  
(d)  $\frac{9\pi}{25}$  cm  
(e) 5 cm

See example 5 p.396 and #59 to 62 p.399

11. The  $x$ - and  $y$ -intercepts of the graph of  $f(x) = 2^{x+1} - 4$  are

- (a)  $(1, 0)$  and  $(0, -2)$
- (b)  $(2, 0)$  and  $(0, 1)$
- (c)  $(0, 0)$  and  $(0, -2)$
- (d)  $(1, 0)$  and  $(0, -1)$
- (e)  $(-1, 0)$  and  $(0, 2)$

12. The smallest positive angle coterminal with the angle  $\theta = -870^\circ$  is equal to

- (a)  $210^\circ$
- (b)  $150^\circ$
- (c)  $30^\circ$
- (d)  $-150^\circ$
- (e)  $-210^\circ$

See example 2 p. 391 and #13 to 18 p. 399

13. The value of  $\csc \frac{5\pi}{3} + \tan \pi + \cot \frac{7\pi}{6}$  is

(a)  $\frac{\sqrt{3}}{3}$

(b)  $-\frac{\sqrt{3}}{2}$

(c)  $\frac{2\sqrt{3}}{3}$

(d)  $-\frac{4\sqrt{3}}{3}$

(e)  $\frac{5\sqrt{3}}{3}$

See example 4 p-430 and #37 to 48 p-432

14. The range of the function

$$y = \frac{1}{3} \csc(2x + 9) - 7$$

is

(a)  $\left(-\infty, -\frac{22}{3}\right] \cup \left[\frac{20}{3}, \infty\right)$

(b)  $\left(-\infty, -\frac{1}{3}\right] \cup \left[\frac{1}{3}, \infty\right)$

(c)  $\left[-\frac{22}{3}, -\frac{20}{3}\right]$

(d)  $\left(-\infty, -\frac{7}{3}\right] \cup \left[\frac{7}{3}, \infty\right)$

(e)  $\left(-\infty, -\frac{20}{3}\right] \cup \left[\frac{22}{3}, \infty\right)$

See example 6 p-453, problems 47 to 49 p-456

15. If the terminal side of an angle  $\theta$  passes through the point  $(-12, 5)$ , then  $\cot \theta + \csc \theta$  is equal to

(a)  $-\frac{1}{5}$

(b)  $-\frac{4}{5}$

(c)  $\frac{4}{5}$

(d)  $\frac{3}{5}$

(e)  $-\frac{3}{5}$

See example 1 p. 416 and # 1 to 6 p. 421

16. The solution set of the equation

$$\log_3(-x) + \log_3(6 - x) = 3$$

contains

- (a) one negative integer only
- (b) two negative integers
- (c) one positive integer and one negative integer
- (d) one positive integer only
- (e) no real numbers

See examples 6, 7 p. 346 and # 25 to 30 p. 349



17. Which one of the following statements is TRUE for the graph of  $y = \left| 3 \sec \frac{2x}{3} \right|$  on the interval  $\left[ 0, \frac{3\pi}{2} \right]$ ?

(a) The graph is symmetric about the line  $x = \frac{3\pi}{4}$ .

(b) The graph is decreasing on  $\left( 0, \frac{3\pi}{4} \right)$ .

(c) The graph is increasing on  $\left( \frac{3\pi}{4}, \frac{3\pi}{2} \right)$ .

(d) The graph has no  $y$ -intercepts.

(e) The graph has only one  $x$ -intercept.

18. The number of the vertical asymptotes for the graph of  $y = 8 \tan \frac{x}{5}$  on the interval  $[0, 10\pi]$  is equal to

(a) 2

(b) 1

(c) 0

(d) 3

(e) 4

See example 2 p-444 and the problems on tangents  
from 21 to 40 p-448

19. The expression  $\frac{1 + \sin \theta}{1 - \sin \theta}$  is identical to

(a)  $(\sec \theta + \tan \theta)^2$

(b)  $(\csc \theta + \cot \theta)^2$

(c)  $(\sec \theta + \sin \theta)^2$

(d)  $(\cos \theta + \tan \theta)^2$

(e)  $(\sec \theta + \csc \theta)^2$

See example 4 p. 474 and problem #57 p. 476

20. The sum of all solutions of the equation

$$e^x + 6e^{-x} = 5$$

is equal to

(a)  $\ln 6$

(b) 1

(c)  $\ln 5$

(d)  $\ln \frac{3}{2}$

(e)  $\ln \frac{2}{3}$

See example 4 p. 345 and #39 to 46 p. 349

21. The product  $\cos 22^\circ \cdot \cot 68^\circ \cdot \csc(-22^\circ)$  simplifies to

(a)  $-1$

(b)  $\cos^2 22^\circ$

(c)  $\sin^2 22^\circ$

(d)  $1$

(e)  $-\cos^2 22^\circ$

A direct application of the Cofunction identities

22. An observer notes that the angle of elevation from point  $A$  to the top of a tower is  $45^\circ$ . From a point 10 m further from point  $A$ , the angle of elevation is  $30^\circ$ . Then the height of the tower is equal to

(a)  $5\sqrt{3} + 5$  m

(b) 5 m

(c) 10 m

(d)  $10\sqrt{3}$  m

(e)  $5\sqrt{3}$  m

See example 6 p. 469 and #72 p. 413

23. The expression  $(\log_3 x)(\ln 9) - \frac{2 \log_3 y}{\log_3 e}$  simplifies to

(a)  $\ln \left( \frac{x^2}{y^2} \right)$

(b)  $\ln(x^2 - y^2)$

(c)  $\ln(x^2 y^2)$

(d)  $\ln(27xy)$

(e)  $\ln \left( \frac{3x}{y^2} \right)$

A direct application of change of base formula and the properties of logarithmic functions

24.  $\frac{\tan \frac{31\pi}{12} + \tan \left( -\frac{\pi}{4} \right)}{1 + \tan \frac{7\pi}{12} \tan \frac{\pi}{4}} =$

(a)  $\tan \frac{\pi}{3}$

(b)  $\tan \frac{\pi}{6}$

(c)  $-\tan \frac{\pi}{3}$

(d)  $-\tan \frac{\pi}{6}$

(e)  $\tan \frac{5\pi}{12}$

See problem 17, 18 p-484

25. Which one of the following statements is TRUE?

- (a) The graphs of  $y = \cos x$  and  $y = \cos |x|$  are the same.
- (b) The function  $y = \sec x$  has infinite number of zeros.
- (c) The phase shift of  $y = \tan\left(3x - \frac{\pi}{6}\right)$  is  $\frac{\pi}{6}$ .
- (d) The maximum value of  $y = -3 \sin(2x + 1)$  is  $-3$ .
- (e) The minimum value of  $y = 9 \csc \frac{2\pi x}{5}$  is  $-9$ .