

**Section 4.1 Linear differential equations: Basic theory****Learning Outcomes**

After completing this section, you will inshaAllah be able to

1. explain what are **initial value problems & boundary value problems** of  $n^{\text{th}}$  order linear differential equations, and their basic properties,
2. identify the **homogeneous and non-homogeneous** ODEs
3. write an **ODE in operator notation**
4. explain about **general solutions of homogeneous linear differential equations**
  - a. basic facts about solutions
  - b. linear independence of solutions
5. **check linear independence** using Wronskians
6. tell about **solutions of non-homogeneous linear differential equations**
  - a. complimentary solution or function
  - b. particular solution
  - c. general solution

### nth order linear differential equations: IVPs

Standard form of  $n^{\text{th}}$  order linear differential equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (*)$$

- Homogeneous if  $g(x) = 0$
- Non-homogeneous if  $g(x) \neq 0$

The corresponding IVP

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1} \quad (2)$$

n conditions

#### Example 5.2.1

- 1)  $2e^x y''' + (2 \sin x) y'' - 4x^2 y' - 5y = 3$  linear, 3<sup>rd</sup> order
- 2)  $xy^{(5)} + 4y^{(4)} + xy''' = x \sin x$  linear, 5<sup>th</sup> order

#### Existence of a unique solution of IVPs

- If  $a_n(x), a_{n-1}(x), \dots, a_0(x)$  and  $g(x)$  are continuous on an interval  $I$  and  $a_n(x) \neq 0$  for all  $x$  in  $I$ .
- Then IVP (1-2) has a unique solution.

See Examples 1, 2 done in class

## Boundary value problems (BVPs)

Conditions specified at different points

Example:

$$a_2(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x) y = g(x)$$

$$y(a) = y_0, y(b) = y_1$$

Solution

- satisfies ODE
- satisfies boundary conditions

Boundary conditions

Can have one or many or no solutions

See Example 3 done in class

## Differential operator notation for ODEs

- Setting


$$D \text{ as } \frac{d}{dx}, D^2 \text{ as } \frac{d^2}{dx^2}, \dots, D^n \text{ as } \frac{d^n}{dx^n}.$$

- An nth order differential operator can be defined as

$$L = a_n(x)D^n + a_{n-1}(x)D^{n-1} + \dots + a_1(x)D + a_0(x).$$

- This can be used to write ODEs in compact form.

See Example 4 done in class



- Here we will only do introduction and basic facts.
- Some methods of finding solutions will be done in subsequent sections.

### Facts about solutions of homogeneous linear equations

- Given  $a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$  (\*)

#### Facts about solution

- If  $y_1(x), y_2(x), \dots, y_k(x)$  are solutions of (\*) then their linear combination

$$y = c_1 y_1(x) + c_2 y_2(x) + \dots + c_k y_k(x)$$

is also a solution of (\*)

- If  $y_1(x), y_2(x), \dots, y_n(x)$  are  $n$  linearly independent solutions of (\*), then the general solution of (\*) is

$$y = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x).$$

If you know few solutions, how many solutions can you generate?

**Superposition Principle**

**General Solution**

So the question of finding all solutions of Eq. (\*) reduces to the **question of finding  $n$  linearly independent solutions**

see next page for the **meaning of linearly independent solutions**

See Example 5 done in class

## Meaning of linearly independent functions

### Definition:

The functions  $y_1(x), y_2(x), \dots, y_n(x)$  are **linearly independent if the equation**

$$c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x) = 0$$

**has only solution**  $c_1 = 0, c_2 = 0, \dots, c_n = 0$ .

Otherwise  $y_1(x), y_2(x), \dots, y_n(x)$  are linearly dependent.

See Examples 6, 7, 8, 9 done in class

The examples show that checking linear independence using definition is not straightforward.

Below we see a practical method of checking linear independence

**Checking linear independence of solutions of ODEs**  
(Practical method)

**Definition:** The **Wronskian** of functions  $y_1(x), y_2(x), \dots, y_n(x)$  is defined as

$$W(y_1, y_2, \dots, y_n) = \begin{vmatrix} y_1 & y_2 & \dots & y_n \\ y_1' & y_2' & \dots & y_n' \\ \vdots & \vdots & \dots & \vdots \\ y_1^{(n-1)} & y_2^{(n-1)} & \dots & y_n^{(n-1)} \end{vmatrix}$$

The solutions  $y_1(x), y_2(x), \dots, y_n(x)$  of  $n^{\text{th}}$  order homogeneous linear differential equation are linearly independent on interval  $I$

$\Leftrightarrow$

the **Wronskian**  $W(y_1, y_2, \dots, y_n) \neq 0$  for some point  $x_0 \in I$ .

See Examples 10, 11, 12, 13 done in class

## Solutions of homogeneous linear equations (Wrap Up)

- Given  $a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$  (\*)
- We have seen that we need to find n linearly independent solutions of (\*).
- Also learnt how to check linear independence of solutions.

We will learn in remaining chapter how to find independent solutions.

- Here we wrap up the facts covered upto now.

A set  $y_1(x), y_2(x), \dots, y_n(x)$  of linearly independent solutions of (\*) is called **fundamental set of solutions** of (\*)

Checking linear independence

- use Wronskian

If  $y_1(x), y_2(x), \dots, y_n(x)$  is a fundamental set of solutions of (\*) then  
 $y = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x)$   
 is a **general solution** of (\*)

Homogeneous linear equations always have fundamental set of solutions

See Example 14 done in class



Next we focus on  
**general solutions** of  $n^{\text{th}}$  order  
**non-homogeneous linear** differential equations

- Only introduction: what is the structure of the general solution of non-homogeneous linear equations
- Methods of finding solutions will be done later

## General solution of non-homogeneous linear differential equations

- Given  $a_n(x)\frac{d^n y}{dx^n} + a_{n-1}(x)\frac{d^{n-1}y}{dx^{n-1}} + \dots + a_1(x)\frac{dy}{dx} + a_0(x)y = g(x)$  (\*)
- The associated homogeneous equation is

$$a_n(x)\frac{d^n y}{dx^n} + a_{n-1}(x)\frac{d^{n-1}y}{dx^{n-1}} + \dots + a_1(x)\frac{dy}{dx} + a_0(x)y = 0 \quad (**)$$

### Determining general solution of (\*) requires

- Find the general solution of associated homogeneous equation (\*\*); denoted as  $y_c$
- Find a particular solution of non-homogeneous equation (\*); denoted as  $y_p$
- The general solution of (\*) is given by  $y = y_c + y_p$ .

Complementary solution or Complimentary function

Particular solution

General Solution

See Example 15 done in class

It can be shown that all solutions of (\*) can be obtained from  $y = y_c + y_p$

### Remember

#### To find general solution of non-homogeneous linear equation

- we need to find general solution of associated homogeneous equation
- we need to find only one particular solution of non-homogeneous equation

**Superposition principle for non-homogeneous linear equations**

Let  $y_{p_1}, y_{p_2}, \dots, y_{p_k}$  be particular solutions of

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y = g_i(x) \text{ for } i = 1, 2, \dots, k.$$

Then

$y_p = c_1 y_{p_1} + c_2 y_{p_2} + \dots + c_k y_{p_k}$  is a particular solution of

$$\begin{aligned} a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y &= \\ &= c_1 g_1(x) + c_2 g_2(x) + \dots + c_k g_k(x) \end{aligned}$$

See Example 16 done in class

*End of 4.1*

*Do Qs. 1-36*