

Exam 1, 101 (012)

Q1 Q0 Speed of sound is 340 m/s. Express this in millimeters
ch Q0 per nanosecond[1 ns = $10^{**}(-9)$ s].

1. Q0

A1 $3.40 * 10^{**}(-4)$ mm/ns

A2 $3.40 * 10^{**}(-6)$ mm/ns

A3 $3.40 * 10^{**}(-3)$ mm/ns

A4 $3.40 * 10^{**}(+3)$ mm/ns

A5 $3.40 * 10^{**}(+6)$ mm/ns

Q0

Q2 Q0 The position of an object moving along an X-axis is
ch Q0 given by $x = 3 + 12*t - t^{**3}$, where x is in meters

2 Q0 and t is in seconds. At what time is the particle
Q0 momentarily at rest?

Q0

A1 2 s

A2 4 s

A3 3 s

A4 1 s

A5 0 s

Q0

Q3 Q0 A rock is dropped ($V_0 = 0$) from a 100-m high cliff. It
ch Q0 takes the rock 3.2 s to fall the first 50 m. How long

2 Q0 does it take to fall the second 50 m?

Q0

A1 1.3 s

A2 1.6 s

A3 4.8 s

A4 3.2 s

A5 0.0 s

Q0

Q4 Q0 The position-time graph for an object is a straight line
Ch Q0 with a positive slope. The object has

2 Q0

A1 a constant velocity

A2 a decreasing acceleration

A3 an increasing velocity

A4 an increasing acceleration

A5 a decreasing velocity

Q0

Q5 Q0 A balloon is going up with a speed of 10 m/s and is
ch Q0 100 m above the ground when a package is dropped from

2 Q0 the balloon. How long does the package take to reach
Q0 the ground?

Q0

A1 5.7 s

A2 4.0 s

A3 3.7 s

A4 2.0 s

A5 6.0 s

Q6 Q0

ch Q0 The two vectors A and B shown in Fig. 1 have equal
3 Q0 magnitudes of 10.0 m. Find the magnitude of the

Q0 resultant, R, of these vectors and the angle theta
Q0 it makes with the positive x-axis.

Q0

A1 $R = 14.1$ m, THETA = 75 degrees

A2 $R = 10.0$ m, THETA = 90 degrees

A3 $R = 12.0$ m, THETA = 60 degrees

A4 $R = 16.0$ m, THETA = 30 degrees

A5 $R = 20.0$ m, THETA = 45 degrees

Q0

Q7 Q0 A vector in the xy-plane has a magnitude of 25.0 and
ch Q0 an x-component of 12.0. The angle that it makes with
Q0 the positive x-axis is:

3 Q0

A1 61.3 degrees

A2 25.6 degrees

A3 28.7 degrees

A4 64.3 degrees

A5 95.3 degrees

Q0

Q8 Q0 The unit vectors in the positive directions of the x,
ch Q0 y, and z axes are labeled i, j, and k. The value of

3 Q0 $[i \cdot (j \times k)]$ is:

Q0

A1 +1

A2 -1

A3 0

A4 -i

A5 +j

Q0

Q9 Q0 Car A is moving with a speed of 30 km/h along the
ch Q0 positive x-axis and car B is moving with a speed of

4 Q0 40 km/h along the positive y-axis. What is the

Q0 velocity of car B with respect to car A?

Q0

A1 $(-30i + 40j)$ km/hA2 $(30i + 40j)$ km/hA3 $(-30i - 40j)$ km/hA4 $(40i + 30j)$ km/hA5 $(40i - 30j)$ km/h

Q0

Q10 Q0 A ball leaves the ground with a speed of 50 m/s at
ch Q0 an angle of 60 degrees with the horizontal. Find its

4 Q0 speed at its highest point.

Q0

A1 25 m/s

A2 50 m/s

A3 0.0 m/s

A4 43 m/s

A5 10 m/s

Q0

Q11 Q0 A stone is thrown from the ground into the air with
ch Q0 an initial velocity $V = (5.0i + 9.0j)$ m/s. To what

4 Q0 maximum height will the stone rise?

Q0

A1 4.1 m

A2 1.3 m

A3 9.0 m

A4 5.0 m

A5 7.0 m

Q0

Q12 Q0 The airplane shown in Fig. 2 is in level flight at an
ch Q0 altitude of 500 m and a speed of 41.7 m/s. At what

4 Q0 distance d should it release a bomb to hit the target
q0 at point A?

Q0

A1 421 m

A2 150 m

A3 300 m

A4 590 m

A5 832 m

Q0

Q13 Q0 A constant force, F, acts on a 19-kg particle. The particle,
ch Q0 initially at rest, moves a distance of 22 m in 3.8 s. Find

5 Q0 the magnitude of the force F.

Q0

A1 58 N

A2 86 N

A3 50 N

A4 41 N

A5 12 N

Q0

Q14 Q0 In Fig. 3, $m_1 = 22$ kg and $m_2 = 37$ kg. The masses are connected
ch Q0 by a light, horizontal cord and are being pulled across a

5 Q0 smooth level surface by a horizontal force $F = 46$ N. Find the

Q0 tension in the cord.

Q0

A1 17 N

A2 29 N

A3 46 N

A4 31 N

A5 63 N

Q0

Q15Q0 Three books (X, Y, and Z) rest on a table as shown in Fig. 4.
ch Q0 The weight of each book is also indicated in the Figure. The
5 Q0 magnitude of the force of book Z on book Y is:

Q0

A1 9.0 N

A2 4.0 N

A3 5.0 N

A4 14 N

A5 19 N

Q0

Q16Q0 Two blocks weighing 25 kg and 35 kg respectively, are
Q0 connected by a string that passes over a massless pulley
Q0 as shown in Fig. 5. The tension in the string is:

Q0

A1 286 N

A2 210 N

A3 500 N

A4 350 N

A5 250 N

Q0

Q17Q0 A 90-kg man stands in an elevator that is moving up at
ch Q0 a constant speed of 5.0 m/s. The magnitude of the force
5 Q0 exerted by him on the floor is:

Q0

A1 882 N

A2 0 N

A3 94 N

A4 450 N

A5 49 N

Q0

Q18Q0 A 3.5-kg block is pulled at constant velocity along a
ch Q0 horizontal floor by a force $F = 15$ N that makes an angle
6 Q0 of 40 degrees with the horizontal (Fig.6). Find the
Q0 magnitude of the force of friction between the block and
Q0 the floor

Q0

A1 11 N

A2 15 N

A3 34 N

A4 0.0 N

A5 26 N

Q0

Q19Q0 Find the minimum coefficient of static friction between
ch Q0 the tyres of a car and a level road if the car is to make
6 Q0 a circular turn of radius 90 m at a speed of 60 km/h.

Q0

A1 0.315

A2 0.521

A3 0.423

A4 0.214

A5 0.125

Q0

Q20Q0 One end of a 1.0-m string is fixed, the other end is attached
ch Q0 to a 1.0-kg stone. The stone swings in a vertical circle,
6 Q0 and has a speed of 5.0 m/s at the top of the circle.

Q0 The tension in the string at this point is approximately:

Q0

A1 15 N

A2 11 N

A3 28 N

A4 31 N
A5 10 N