

First Major Exam 031

Q1 Q0 An empty fuel tank of a car needs 50 liters of gasoline

Q0 to fill up. Find the volume of the fuel tank in m^3 .

Q0 (1 milliliter = 1 cm^3)

Q0

A1 0.050

A2 50 000

A3 50

A4 500

A5 0.50

Q0

Q2 Q0 Fig. 1 shows a graph of position versus time for a particle

Q0 moving along the x axis. What is the total distance travelled

Q0 by the particle in 15 s?

Q0

A1 12.5 m

A2 7.5 m

A3 10 m

A4 5.0 m

A5 22.5 m

Q0

Q3 Q0 An object starts from rest at the origin and moves along the

Q0 x-axis with a constant acceleration of 5.0 m/s^2 . Find its

Q0 average velocity as it goes from $x = 0 \text{ m}$ to $x = 10 \text{ m}$.

Q0

A1 5.0 m/s

A2 10 m/s

A3 17 m/s

A4 3.0 m/s

A5 8.0 m/s

Q0

Q4.Q0 Starting at time $t = 0$, an object moves along a straight line

Q0 with a velocity in m/s given by $v = 72 - 2t^2$,

Q0 where t is in seconds. Find its acceleration when it stops

Q0 momentarily.

Q0

A1 -24 m/s^2

A2 0

A3 -4.0 m/s^2

A4 -9.8 m/s^2

A5 -4.9 m/s^2

Q0

Q5 Q0 A stone is thrown vertically upward with an initial speed of

Q0 15 m/s. What is its speed at a height of 10 m from its release

Q0 point?

Q0

A1 5.4 m/s

A2 0

A3 It will not reach the height of 10 m.

A4 9.8 m/s

A5 12 m/s

Q0

Q6 Q0 The angle between the two vectors $A = 2i + 4j$ and

Q0 $B = 4i - 2j$ is:

Q0

A1 90 degrees

A2 27 degrees

A3 39 degrees

A4 180 degrees

A5 0 degrees

Q0

Q7 Q0 As shown in Fig. 3, a block moves down on a 45-degree inclined

Q0 plane of 2.5 m length, then horizontally for another 2.5 m, and

Q0 then falls down vertically a height of 2.5 m. Find the magnitude

Q0 and direction of the resultant displacement vector of the block.

Q0

A1 6.0 m and 45 degrees below horizontal axis

A2 3.5 m and 30 degrees below horizontal axis

A3 6.0 m and 30 degrees below horizontal axis

A4 3.5 m and 45 degrees below horizontal axis

A5 5.5 m and 60 degrees below horizontal axis

Q0

Q8 Q0 Given the vectors $A = 3j + 6k$, $B = 15i + 21k$. Find the

Q0 magnitude of vector C that satisfies equation $2A + 3C - B = 0$.

Q0

A1 6.16

A2 5.48

A3 18.5

A4 6.71

A5 8.60

Q0

9 Q0 At $t=0$, a particle moving in the xy plane with a constant

Q0 acceleration of $a=(2i + 4j)$ m/s^2 has a velocity $V_0=(-4j)$ m/s

Q0 at the origin. Find the speed of the particle at $t=3$ s.

Q0

A1 10 m/s

A2 0

A3 4 m/s

A4 24 m/s

A5 20 m/s

Q0

10 Q0 A ball is projected from the ground into the air with velocity

Q0 V_0 . At a height of 10.0 m the velocity is observed to be

Q0 $V = 8.5i + 9.1j$ in m/s . Find V_0 .

Q0

A1 $(8.5i + 16.7j)$ m/s

A2 $(16.7i + 9.1j)$ m/s

A3 $(8.5i + 9.1j)$ m/s

A4 $(2.5i + 3.1j)$ m/s

A5 $(6.2i + 1.1j)$ m/s

Q0

11 Q0 Rain is falling vertically at constant speed of 6.0 m/s .

Q0 At what angle from the vertical do the rain appear to be falling

Q0 as viewed by the driver of a car traveling on a straight, level

Q0 road with a speed of 8.0 m/s ?

Q0

A1 53 degrees

A2 37 degrees

A3 49 degrees

A4 41 degrees

A5 0 degree

Q0

12 Q0 The speed of a particle moving in uniform circular motion is

Q0 doubled while the radius of the path of the particle is

Q0 increased by a factor of 4. The new centripetal force needed

Q0 will be :

Q0

A1 the same as before

A2 half as great as before

A3 twice as great as before

A4 1/4 of its original value

A5 four times as great as before

Q0

13 Q0 A ball is thrown horizontally with speed V_0 from the edge of

Q0 a cliff 35 m high. The ball strikes the ground at a point 80 m

Q0 from the base of the cliff. Find V_0 .

Q0

A1 30 m/s

A2 9.8 m/s

A3 2.5 m/s

A4 22 m/s

A5 45 m/s

Q0

14 Q0 As shown in Fig. 7, a 25-kg box is pushed across a frictionless

Q0 horizontal floor with a force of 20 N, directed at an angle of

Q0 20 degrees below the horizontal. The magnitude of the

Q0 acceleration of the box is :

Q0

A1 0.75 m/s^2

A2 0.27 m/s^2

A3 17 m/s^2

A4 21 m/s^2

A5 0.82 m/s^2

Q0

15 Q0 An object of mass $M = 10 \text{ kg}$ moving on frictionless horizontal

Q0 surface is subjected to two applied forces as shown in Fig. 2.

Q0 In which situation is the object accelerating to the right?

Q0

A1 (d)

A2 (a)

A3 (c)

A4 (b)

A5 (e)

Q0

16 Q0 Two blocks A ($M_A = 4 \text{ kg}$) and B ($M_B = 20 \text{ kg}$) are in contact with

Q0 each other and are placed on a horizontal frictionless surface.

Q0 A 36-N constant force is applied to A as shown in Fig. 4. The

Q0 magnitude of the force exerted on A by B is

Q0

A1 30 N

A2 0 N

A3 36 N

A4 15 N

A5 3.6 N

Q0

17 Q0 Two masses $m_1 = 2\text{kg}$, $m_2 = 4\text{kg}$ are connected by a light string

Q0 that passes over a frictionless and massless pulley (see Fig. 5).

Q0 Find the magnitude of the acceleration of the masses.

Q0

A1 3.27 m/s^2

A2 2.15 m/s^2

A3 10.5 m/s^2

A4 0.75 m/s^2

A5 1.23 m/s^2

Q0

18 Q0 A stone, of mass m , is attached to a strong string and rotates

Q0 in a vertical circle of radius R . At the bottom of the path the

Q0 tension in the string is 3 times the weight of the stone. The

Q0 speed of the stone at this point is given by .

Q0

A1 $\sqrt{2gR}$.

A2 $2\sqrt{gR}$

A3 $2gR$

A4 $\sqrt{3gR}$

A5 $\sqrt{gR/2}$

Q0

19 Q0 A block attached to a string, rotates counter-clockwise in a

Q0 circle on a smooth horizontal surface. The string breaks at

Q0 point P (Fig. 6). What path will the block follow?

Q0

A1 path B

A2 path A

A3 path C

A4 path D

A5 path E

Q0

20 Q0 A box slides down a 30 degree incline with an acceleration =

Q0 3.2 m/s^2 . Find the coefficient of kinetic friction between

Q0 the box and the incline.

Q0

A1 0.20

A2 0.25

A3 0.15

A4 0.30

A5 0.62

PHYS101 - FIRST MAJOR EXAM – FIGURES
Term -031




FIGURE-3




FIGURE-2




FIGURE-4




FIGURE-5




FIGURE-6




FIGURE-7

