

# King Fahd University of Petroleum and Minerals

## Department of Physics



PHYS101-051  
MAJOR 1 EXAM


**Test Code: 000**

*("A" is the correct answer in all questions)*

17 October 2005 in Building 54  
Exam Duration: 2hrs (from 9:00pm to 11:00pm)

Name:	
Student Number:	
Section Number:	


- The mass of  $1.0 \text{ cm}^3$  of gold is 19.3 g. What is the mass of a solid cube of gold having a side of 0.70 cm?
  - $6.6 \times 10^{-3} \text{ kg}$
  - $9.1 \times 10^{-2} \text{ kg}$
  - $3.6 \times 10^{-3} \text{ kg}$
  - 0.11 kg
  - 21 kg
- A helicopter at height  $h$  (m) from the surface of the sea is descending at a CONSTANT SPEED  $v$  (m/s). The time it takes to reach the surface of the sea can be found from:
  - $-h = -v t$
  - $h = \frac{1}{2} g t^2$
  - $-h = \frac{1}{2} g t^2$
  - $h = v t - \frac{1}{2} g t^2$
  - $-h = -v t - \frac{1}{2} g t^2$
- A particle starts from rest at  $t = 0$  s. Its acceleration as a function of time is shown in Fig. 1. What is its speed at the end of the 6.0 s?



- 4.0 m/s
  - 0 m/s
  - 12 m/s
  - 2.0 m/s
  - 12 m/s
- The position of a particle  $x(t)$  as a function of time ( $t$ ) is described by the equation:  $x(t) = 2.0 + 3.0 t - t^3$ , where  $x$  is in m and  $t$  is in s. What is the maximum positive position of the particle on the  $x$  axis?
    - 4.0 m
    - 2.0 m
    - 3.0 m
    - 1.0 m
    - 5.0 m


5. A stone is thrown vertically downward from a building with an initial speed of 2.0 m/s. It reaches the ground after 5.0 s. What is the height of the building?
- A) 130 m  
 B) 60 m  
 C) 180 m  
 D) 120 m  
 E) 140 m
6. Three vectors  $\vec{A}$ ,  $\vec{B}$ , and  $\vec{C}$  are such that:  $\vec{C} = \vec{A} + \vec{B}$ ,  $\vec{B} = 5\hat{i}$  and  $\vec{C} = 5\hat{j}$ . Find the angle between  $\vec{A}$  and  $\vec{B}$ .
- A)  $135^\circ$   
 B)  $120^\circ$   
 C)  $270^\circ$   
 D)  $150^\circ$   
 E)  $45^\circ$
7. A man walks 4.65 km West, then 12.7 km in the direction  $30^\circ$  West of North and finally 11.0 km due East. The man is now at
- A) 11.0 km due North  
 B) 12.7 km due West  
 C) 4.65 km due South  
 D) 15.6 km in the direction  $45^\circ$  West of North  
 E) back to where he started
8. If vector  $\vec{A}$  has the magnitude of 3.0 m and makes an angle  $30^\circ$  with the +x-axis, then the vector  $\vec{B} = -2\vec{A}$  is:
- A)  $\vec{B} = -5.2\hat{i} - 3.0\hat{j}$  (m)  
 B)  $\vec{B} = 5.2\hat{i} + 3.0\hat{j}$  (m)  
 C)  $\vec{B} = -5.2\hat{i} + 3.0\hat{j}$  (m)  
 D)  $\vec{B} = 5.2\hat{i} - 3.0\hat{j}$  (m)  
 E)  $\vec{B} = -3.0\hat{i} - 5.2\hat{j}$  (m)
9. A ball is thrown with a velocity  $\vec{v}_o = 3.0\hat{i} + 5.0\hat{j}$  (m/s) from the ground. Its velocity just before it strikes the ground is:
- A)  $\vec{v} = 3.0\hat{i} - 5.0\hat{j}$  (m/s)  
 B)  $\vec{v} = 3.0\hat{i} + 5.0\hat{j}$  (m/s)  
 C)  $\vec{v} = 3.0\hat{i}$  (m/s)  
 D)  $\vec{v} = 5.0\hat{j}$  (m/s)  
 E)  $\vec{v} = -5.0\hat{j}$  (m/s)

10. A ball is kicked from the ground with an initial speed of 20 m/s at an angle of  $45^\circ$ . A player 60 m away starts running to catch the ball at that instant (see Fig 2). What must be his average speed ( $v$ ) if he has to catch the ball just before it hits the ground?




- A) 6.6 m/s  
 B) 10 m/s  
 C) 20 m/s  
 D) 2.0 m/s  
 E) 15 m/s
11. The position of a particle as a function of time is given by  $\vec{r} = 3.0t\hat{i} + 2.0t^2\hat{j}$ . Find the angle between the velocity and acceleration of the particle at  $t = 5.0 \text{ s}$ .
- A)  $8.5^\circ$   
 B)  $0^\circ$   
 C)  $90^\circ$   
 D)  $45^\circ$   
 E)  $78^\circ$
12. Car A is moving towards East with speed 15.0 m/s and car B is moving towards West with speed 25.0 m/s, both relative to the ground. Find the velocity of car B relative to car A.
- A) 40.0 m/s towards West  
 B) 5.00 m/s towards West  
 C) 5.00 m/s towards North  
 D) 40.0 m/s towards East  
 E) 40.0 m./s towards South

13. Two forces  $\vec{F}_1$  &  $\vec{F}_2$  are acting on a 3.0 kg box in the x-y plane. Fig. 3 shows only  $\vec{F}_1$  and the acceleration  $\vec{a}$  of the box. Find  $\vec{F}_2$




- A)  $(-72\hat{i} + 24\hat{j})N$   
 B)  $(-72\hat{i} - 24\hat{j})N$   
 C)  $(72\hat{i} + 24\hat{j})N$   
 D)  $(-36\hat{i} - 48\hat{j})N$   
 E)  $(-36\hat{i} + 48\hat{j})N$
14. A block of mass  $m_1=5.7$  kg on a frictionless  $30^\circ$  inclined plane is connected by a cord over a massless, frictionless pulley to a second block of mass  $m_2=3.5$  kg hanging vertically as shown in Fig 4. The acceleration of  $m_2$  is:




- A)  $0.69 \text{ m/s}^2$  downward  
 B)  $0.54 \text{ m/s}^2$  upward  
 C)  $0.36 \text{ m/s}^2$  downward  
 D)  $0.78 \text{ m/s}^2$  upward  
 E)  $0.93 \text{ m/s}^2$  downward


15. Fig.5 shows a block A of mass 6.0 kg and block B of 8.0 kg connected by a rigid rod of negligible mass. Force  $\vec{F}_a = (16\text{N})\hat{i}$  acts on block A; force  $\vec{F}_b = -(30\text{N})\hat{i}$  acts on block B. The tension (T) in the rod is: (Neglect friction)



- A) 22 N  
 B) 30 N  
 C) 16 N  
 D) 46 N  
 E) 14 N
16. A 5.0-kg mass is held at an angle  $\theta$  from the vertical by a horizontal force  $F=15\text{ N}$  as shown in Fig 6. The tension (T) in the string supporting the mass (in Newton) is:




- A) 51  
 B)  $2/\cos\theta$   
 C)  $\cos\theta/2$   
 D)  $\cos\theta$   
 E) 0
17. A 0.20-kg stone is attached to a string and whirled in a circle of radius  $r = 0.60\text{ m}$  on a horizontal frictionless surface as shown in Fig. 7. If the stone makes 150 revolutions per minute, the tension (T) in the string is:



- A) 30 N  
 B) 0.20 N  
 C) 0.90 N  
 D) 1.96 N  
 E) 0.03 N

18. A block of mass  $M$  slides on a horizontal surface. Which of the following would increase the magnitude of the frictional force on the block?
- A) Increasing  $M$
  - B) Keeping  $M$  constant but decreasing the surface area of contact
  - C) Keeping  $M$  constant but increasing the surface area of contact
  - D) Decreasing  $M$
  - E) None of the other answers
19. A box of mass  $m$  is sliding down a rough inclined plane (which makes an angle of  $30^\circ$  with the horizontal and has a coefficient of kinetic friction  $= \mu_k$ ) at a constant acceleration  $g/4$  (where  $g = 9.8 \text{ m/s}^2$ ). Find  $\mu_k$ .
- A) 0.29
  - B) 0.16
  - C) 2.15
  - D) 0.11
  - E) 0.64
20. A 5.0 kg block is sliding on a rough horizontal plane ( $\mu_k=0.10$ ) under the effect of a horizontal force  $F$ . Fig. 8 shows the velocity ( $v$ ) of the block as a function of time ( $t$ ). Calculate  $F$ .



- A) 15 N
- B) 5.0 N
- C) 10 N
- D) 1.0 N
- E) 30 N

# PHYS101 First Major Exam Formula Sheet

$$y = cx^n; \quad \frac{dy}{dx} = cnx^{n-1}$$

## Motion in One Dimension

$$v = \frac{dx}{dt}; \quad a = \frac{dv}{dt}; \quad v_{avg} = \frac{\Delta x}{\Delta t}; \quad a_{avg} = \frac{\Delta v}{\Delta t}$$

## Motion with Constant Acceleration

$v = v_o + at$	$x - x_o = v_o t + \frac{1}{2}at^2$
$v^2 = v_o^2 + 2a(x - x_o)$	$x - x_o = \frac{1}{2}(v + v_o)t$
	$x - x_o = v t - \frac{1}{2}at^2$

## Free Fall

$$a = -g; \quad g = 9.8m/s^2$$

## Vectors

$$\vec{a} \cdot \vec{b} = ab \cos \phi \quad \left| \vec{a} \times \vec{b} \right| = ab \sin \phi$$

## Motion in Two Dimensions

$$\vec{v} = \frac{d\vec{r}}{dt}; \quad \vec{a} = \frac{d\vec{v}}{dt}$$

$$\vec{r} - \vec{r}_o = \vec{v}_o t + \frac{1}{2}\vec{a}t^2; \quad \vec{v} = \vec{v}_o + \vec{a}t$$

## Projectile Motion

$a_x = 0$	$x - x_o = v_o \cos \theta_o t$
$a_y = -g = -9.80 \text{ m/s}^2$	$y - y_o = v_o \sin \theta_o t - \frac{1}{2}gt^2$
$H = v_o^2 \sin^2 \theta_o / 2g$	$R = v_o^2 \sin 2\theta_o / g$

## Uniform Circular Motion

$$a = \frac{v^2}{r}$$

$$T = \frac{2\pi r}{v}$$

## Relative Motion

$$\vec{v}_{PA} = \vec{v}_{PB} + \vec{v}_{BA}$$

$$\vec{v}_{AB} = \text{velocity of A relative to B} = -\vec{v}_{BA}$$

## Newton's Second Law

$$\sum \vec{F} = m\vec{a} \Rightarrow \sum F_x = ma_x; \quad \sum F_y = ma_y$$

## Friction

$$f_{s,max} = \mu_s N; \quad f_k = \mu_k N$$



## Answer Key

1. A
2. A
3. A
4. A
5. A
6. A
7. A
8. A
9. A
10. A
11. A
12. A
13. A
14. A
15. A
16. A
17. A
18. A
19. A
20. A