

# King Fahd University of Petroleum and Minerals

## Department of Physics



PHYS101-052  
FINAL EXAM  
**Test Code: 100**

Tuesday, 29 May 2006  
Exam Duration: 3 hrs (from 12:30pm to 3:30pm)

Name:	
Student Number:	
Section Number:	

1. The graph of Fig 1 represents the straight line motion of a car that starts at  $t = 0, x = 0$ . What is the position of the car at  $t = 4$  s?
  - A) 30 m
  - B) 0
  - C) -40 m
  - D) 20 m
  - E) -20 m
  
2. As shown in Fig 2, vector  $\vec{A}$  has magnitude of 12 m and vector  $\vec{B}$  has magnitude of 8 m.  $\vec{A} - \vec{B}$  in vector notation is:
  - A)  $(10.3\text{m}) \mathbf{i} + (1.90\text{m}) \mathbf{j}$
  - B)  $(8.40\text{m}) \mathbf{i} + (2.90\text{m}) \mathbf{j}$
  - C)  $(4.49\text{m}) \mathbf{i} + (15.4\text{m}) \mathbf{j}$
  - D)  $0 \mathbf{i} + 0 \mathbf{j}$
  - E)  $(14.4\text{m}) \mathbf{i} + (2.50\text{m}) \mathbf{j}$
  
3. A boy on the edge of a vertical building 19.6 m high throws a stone horizontally outward with a speed of 20.0 m/s. It strikes the ground at a horizontal distance ( $x$ ) from the foot of the building. Find the value of  $x$ .
  - A) 10.0 m
  - B) 9.80 m
  - C) 50.0 m
  - D) 19.6 m
  - E) 40.0 m
  
4. A stone is tied to a 0.50-m string and moves in a vertical circle at a constant speed of 4.0 m/s. Its acceleration at the top of the circle is:
  - A)  $9.8 \text{ m/s}^2$ , up
  - B)  $9.8 \text{ m/s}^2$ , down
  - C)  $0.0 \text{ m/s}^2$
  - D)  $32 \text{ m/s}^2$ , up
  - E)  $32 \text{ m/s}^2$ , down
  
5. A 2.0-kg block slides on a horizontal surface. Part of the surface is smooth and the other part is rough. A horizontal force is applied to the block. On the smooth part, the acceleration of the block is  $3.0 \text{ m/s}^2$ , while it is  $2.0 \text{ m/s}^2$  on the rough part. What is the magnitude of the frictional force on the rough part?
  - A) 8.0 N
  - B) 6.0 N
  - C) 4.0 N
  - D) 2.0 N
  - E) 10 N
  
6. A mass is suspended by a string from the ceiling of a train accelerating horizontally at  $2.5 \text{ m/s}^2$ . The angle that the string makes with the vertical is:
  - A)  $10^\circ$
  - B)  $16^\circ$
  - C)  $14^\circ$
  - D)  $12^\circ$
  - E)  $30^\circ$

7. At time  $t = 0$ , a 2.0-kg particle has a velocity of  $\vec{v}_i = (8.0 \text{ m/s})\hat{i} - (6.0 \text{ m/s})\hat{j}$ . At time  $t = 3.0$  s its velocity is  $\vec{v}_f = (3.0 \text{ m/s})\hat{i} + (4.0 \text{ m/s})\hat{j}$ . During this time interval the net work done on it is:
- 100 J
  - 25 J
  - 75 J
  - 50 J
  - $(-5.0 \text{ J})\hat{i} + (10 \text{ J})\hat{j}$
8. A 6.0-kg block is released from rest 80 m above the ground. When it reaches the ground its kinetic energy is:
- 3500 J
  - 4700 J
  - 1200 J
  - 120 J
  - 640 J
9. A 0.50-kg block attached to an ideal spring with a spring constant of 80 N/m oscillates on a horizontal frictionless surface. The total mechanical energy is 0.12 J. The greatest extension of the spring from its equilibrium length is:
- 0.015 m
  - 0.030 m
  - 0.039 m
  - 0.055 m
  - 18 m
10. A 2.2-kg block starts from rest on a rough inclined plane ( $\mu_k = 0.25$ ) that makes an angle of  $25^\circ$  with the horizontal. As the block goes 2.0 m down the plane, the change in mechanical energy of the block is:
- 0
  - 9.8 J
  - 9.8 J
  - 18 J
  - 18 J
11. A 0.20-kg rubber ball is dropped from the window of a building. It strikes the sidewalk below at 30 m/s and rebounds up at 20 m/s. The impulse on the ball during collision is:
- 9.8 N·s upward
  - 10 N·s downward
  - 2 N·s upward
  - 2 N·s downward
  - 10 N·s upward
12. Blocks A and B are moving toward each other. Block A has a mass of 2.00 kg and a velocity of 50.0 m/s, while block B has a mass of 4.00 kg and a velocity of -25.0 m/s. They suffer a completely inelastic collision. The kinetic energy lost during the collision is:
- 0
  - 1250 J
  - 3750 J
  - 5000 J
  - 5600 J

13. A solid ball of mass  $M = 400 \text{ g}$  and radius  $R = 5.0 \text{ cm}$  is rotating about its fixed central axis with angular speed of  $3.0 \text{ rad/s}$ . It was brought to a stop in  $6.0 \text{ s}$ . The work done to stop the ball is:
- $-1.8 \times 10^{-3} \text{ J}$
  - $-3.0 \times 10^{-4} \text{ J}$
  - $-4.8 \times 10^{-4} \text{ J}$
  - $-3.6 \times 10^{-3} \text{ J}$
  - $-9.0 \times 10^{-3} \text{ J}$
14. Two identical thin (negligible radius) rods are joined together to form the shape shown in Fig 3. Each rod has a mass  $M$  and length  $L$ . The rotational inertia of the assembly about the  $y$  axis is:
- $(1/12) ML^2$
  - $(1/6) ML^2$
  - $ML^2$
  - $(1/2) ML^2$
  - $(1/3) ML^2$
15. A projectile of mass  $m=0.50 \text{ kg}$  moves to the right with speed  $v_0=8.0 \text{ m/s}$  (see Fig 4). The projectile strikes and sticks to the end of a stationary thin rod of mass  $M=6.0 \text{ kg}$  and length  $L=1.0 \text{ m}$  that is pivoted about a frictionless vertical axle through its center ( $O$ ). The final angular velocity ( $\omega$ ) of the (projectile + rod) after collision is:
- $1.2 \text{ rad/s}$  clockwise
  - $1.0 \text{ rad/s}$  clockwise
  - $3.2 \text{ rad/s}$  clockwise
  - $4.0 \text{ rad/s}$  counterclockwise
  - $2.4 \text{ rad/s}$  clockwise
16. A  $5.0\text{-m}$  weightless rod ( $AC$ ), hinged to a wall at  $A$ , is used to support an  $800\text{-N}$  block as shown in Fig 5. The horizontal and vertical components of the force ( $F_H$ ,  $F_V$ ) of the hinge on the rod are:
- $F_H = 800 \text{ N}$ ,  $F_V = 800\text{N}$
  - $F_H = 800 \text{ N}$ ,  $F_V = 600\text{N}$
  - $F_H = 0$ ,  $F_V = 800\text{N}$
  - $F_H = 1200 \text{ N}$ ,  $F_V = 800\text{N}$
  - $F_H = 600 \text{ N}$ ,  $F_V = 800\text{N}$
17. A shearing force  $F = 50 \text{ N}$  is applied to an aluminum rod with a length of  $L = 10 \text{ m}$ , a cross-sectional area  $A=1.0 \times 10^{-5} \text{ m}^2$ , and a shear modulus  $G = 2.5 \times 10^{10} \text{ N/m}^2$ . As a result the rod is sheared through a distance ( $\Delta x$ ) of:
- $0.10 \text{ cm}$
  - $0.30 \text{ cm}$
  - $0.20 \text{ cm}$
  - $0.40 \text{ cm}$
  - $0.50 \text{ cm}$
18. A man holds a rod  $AB$  of length  $= 6.0 \text{ m}$  and weight  $= 30 \text{ N}$  in equilibrium, by exerting an upward force  $F_1$ , with one hand, and a downward force  $F_2$ , with the other hand as shown in Fig 6. What are the magnitude of the forces  $F_1$  and  $F_2$ ?
- $F_1 = 90 \text{ N}$ ,  $F_2 = 60 \text{ N}$
  - $F_1 = 30 \text{ N}$ ,  $F_2 = 30 \text{ N}$
  - $F_1 = 60 \text{ N}$ ,  $F_2 = 40 \text{ N}$
  - $F_1 = 40 \text{ N}$ ,  $F_2 = 50 \text{ N}$
  - $F_1 = 20 \text{ N}$ ,  $F_2 = 60 \text{ N}$

19. Three identical particles each of mass  $m$  are distributed along the circumference of a circle of radius  $R$  as shown in Fig 7. The gravitational force of  $m_2$  on  $m_1$  is  $1.0 \times 10^{-6}$  N. The magnitude of the net gravitational force on  $m_1$  due to  $m_2, m_3$  is:
- $3.0 \times 10^{-6}$  N
  - $1.4 \times 10^{-6}$  N
  - $2.0 \times 10^{-6}$  N
  - 0
  - $2.5 \times 10^{-6}$  N
20. Calculate the mass of the Sun using the fact that Earth is rotating around the Sun in a circular orbit of radius  $1.496 \times 10^{11}$  m with a period of one year (1 year =  $3.156 \times 10^7$  s).
- $1.99 \times 10^{30}$  kg
  - $6.42 \times 10^{32}$  kg
  - $4.88 \times 10^{28}$  kg
  - $3.18 \times 10^{26}$  kg
  - $5.98 \times 10^{24}$  kg
21. Calculate the work require to move an Earth satellite of mass  $m$  from a circular orbit of radius  $2R_E$  to one of radius  $3R_E$ . (Consider  $M_E$  = mass of Earth,  $R_E$  = radius of Earth,  $G$  = universal Gravitational constant)
- $G M_E m / (6 R_E)$
  - $G M_E m / (8 R_E)$
  - $G M_E m / (4 R_E)$
  - $G M_E m / (12 R_E)$
  - $G M_E m / (3 R_E)$
22. A 100-kg rock from outer space is heading directly toward Earth. When the rock is at a distance of  $(9R_E)$  from the Earth's surface, its speed is 12 km/s. Neglecting the effects of the Earth's atmosphere on the rock, find the speed of the rock just before it hits the surface of Earth.
- 12 km/s
  - 16 km/s
  - 0
  - 20 km/s
  - 18 km/s
23. A uniform U-tube is partially filled with water. Oil, of density  $0.75 \text{ g/cm}^3$ , is poured into the right arm until the water level in the left arm rises 3.0 cm (see Fig 8). The length of the oil column ( $h$ ) is then:
- 2.2 cm
  - 8.0 cm
  - 3.0 cm
  - 4.0 cm
  - 10 cm
24. The dimensions of a boat ( $\rho_{\text{boat}} = 150 \text{ kg/m}^3$ ) is  $3.00 \text{ m} \times 3.00 \text{ m} \times 1.00 \text{ m}$ . What maximum load can it carry in sea water ( $\rho_{\text{sea water}} = 1020 \text{ kg/m}^3$ ) without sinking?
- 1350 kg
  - 7830 kg
  - 9200 kg
  - 19500 kg
  - 24300 kg

25. A water line enters a house 2.0 m below ground. A smaller diameter pipe carries water to a faucet 5.0 m above ground, on the second floor. Water flows at 2.0 m/s in the main line and at 7.0 m/s on the second floor. If the pressure in the main line is  $3.0 \times 10^5$  Pa, then the pressure on the second floor is:  
(Take the density of water to be  $1.0 \times 10^3$  kg/m<sup>3</sup>)
- $5.3 \times 10^4$  Pa
  - $4.5 \times 10^5$  Pa
  - $1.1 \times 10^5$  Pa
  - $2.1 \times 10^5$  Pa
  - $3.4 \times 10^5$  Pa
26. The rate of flow of water through a horizontal pipe is 2.00 m<sup>3</sup>/min. Calculate the speed of flow at a point where the diameter of the pipe is 10.0 cm.
- 4.24 m/s
  - 2.00 m/s
  - 0.20 m/s
  - 20.0 m/s
  - 2.12 m/s
27. A mass  $m_1 = 1.0$  kg is connected to a spring (with spring constant equal to  $k$ ) and oscillates on a horizontal frictionless table with a period of 1.0 s. When  $m_1$  is replaced with another unknown mass  $m_2$ , the period changes to 2.0 s. Find the value of  $m_2$ .
- 4.0 kg
  - 2.0 kg
  - 1.0 kg
  - 0.5 kg
  - 0.25 kg
28. A 0.500 kg block is connected to a spring ( $k = 20.0$  N/m) and oscillates on a horizontal frictionless table. Calculate the maximum kinetic energy of the block if the amplitude of the simple harmonic motion is 3.00 cm.
- $4.00 \times 10^{-4}$  J
  - $8.00 \times 10^{-2}$  J
  - $3.00 \times 10^{-1}$  J
  - $9.00 \times 10^{-3}$  J
  - $5.00 \times 10^{-5}$  J
29. If the displacement of a block-spring system is described by the following equation  $x(t) = 0.2 \cos(10t)$  where  $x$  is in m, and  $t$  is in s. What is the speed of the block when its displacement is  $x = 0.1$  m?
- 10.0 m/s
  - 1.73 m/s
  - 0.30 m/s
  - 2.00 m/s
  - 1.00 m/s
30. A simple pendulum has a period of 10.0 s if the free fall acceleration is  $g$ . What would its period be if the free fall acceleration is  $g/2$ ?
- 14.1 s
  - 20.0 s
  - 5.00 s
  - 10.0 s
  - 7.07 s




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

## Answer Key

1. E
2. C
3. E
4. E
5. D
6. C
7. C
8. B
9. D
10. B
11. E
12. C
13. A
14. E
15. C
16. E
17. C
18. A
19. B
20. A
21. D
22. B
23. D
24. B
25. D
26. A
27. A
28. D
29. B
30. A