

Second Major (T031)

- (1)Q0 Five moles of an ideal gas undergo a reversible isothermal
21Q0 compression from volume V to volume $V/2$ at temperature
Q0 30 degrees C. What is the change in the entropy of the gas?
Q0
A1 -29 J/K.
A2 29 J/K.
A3 18 J/K.
A4 -81 J/K.
A5 -18 J/K.
Q0
- (2)Q0 An automobile engine operates with an overall efficiency
21Q0 of 20%. How many gallons of gasoline is wasted for each
Q0 10 gallons burned?
Q0
A1 8.
A2 12.
A3 6.
A4 10.
A5 2.
Q0
- (3)Q0 A heat engine operates between 600 K and 300 K. In each cycle
21Q0 it takes 100 J from the hot reservoir, loses 25 J to the cold
Q0 reservoir, and does 75 J of work. This heat engine violates:
Q0
A1 The second law but not the first law of thermodynamics.
A2 Both, the first law and the second law of thermodynamics.
A3 The first law but not the second law of the thermodynamics.
A4 Neither the first law nor the second law.
A5 Conservation of energy.
Q0
- (4)Q0 As in figure (1), a charge Q is fixed at each of two opposite
22Q0 corners of a square. A charge q is fixed at each of the other
Q0 two corners. If the resultant electrical force on Q is zero,
Q0 then Q and q are related as:
Q0
A1 $Q = - 2 \sqrt{2} q$
A2 $Q = - 4 q$
A3 $Q = q$
A4 $Q = q^2$
A5 $Q = - 2 \sqrt{2} q^2$
Q0
- (5)Q0 Consider two identical conductor spheres, A and B.
22 Q0 Initially, sphere A has a charge of $-80 Q$ and Sphere B
Q0 has a charge of $+20 Q$. If the spheres touched and then
Q0 are separated by a distance of 0.3 m, what is the
Q0 resultant force between them? [Take $Q = 5.7 \times 10^{-8} \text{ C}$]
Q0
A1 0.3 N.
A2 0.2 N.
A3 0.4 N.
A4 0.6 N.
A5 0.9 N.
Q0
- (6)Q0 A particle, of mass m and charge q , is released from rest
23 Q0 at point A in a uniform electric field, see figure (2).
Q0 The kinetic energy, due to the electric field, it attains
Q0 after moving a distance y is:
Q0

- A1 $q \cdot E \cdot y$.
 A2 $m \cdot q \cdot E \cdot y$.
 A3 $q \cdot E \cdot y / 2$.
 A4 $q \cdot E \cdot y^{**2}$.
 A5 $E \cdot y$.
 Q0
- 7 Q0 Which of the following statements are CORRECT:
 Q0
- 23 Q0 1. Electric charge is quantized.
 Q0 2. The potential at the center of a charged conductor is zero.
 Q0 ->
 Q0 3. If $E = 0$ at a point P then V must be zero at P.
 Q0 4. The electric field inside a charged conductor is zero.
 Q0 ->
 Q0 5. If $V = 0$ at a point P then E must be zero at P.
 Q0
- A1 1 and 4.
 A2 2 and 4.
 A3 1, 2 and 3.
 A4 1, 2, and 5.
 A5 3 and 5.
 Q0
- (8) Q0 A long nonconducting cylinder (radius 12.0 cm) has a charge
 24 Q0 of uniform density 5.0 nano-C/m**3 distributed through
 Q0 its column. Determin the magnitude of the electric field
 Q0 5.0 cm from the axis of the cylinder. [See figure (3)].
 Q0
- A1 14 N/C.
 A2 22 N/C.
 A3 31 N/C.
 A4 34 N/C.
 A5 4 N/C.
 Q0
- (9) Q0 In figure (4), what is the magnitude of the electric field
 23 Q0 at point P, center of the equilateral triangle?
 Q0 [take $d = 2$ m, $q = 10^{**(-9)}$ C]
 Q0
- A1 Zero.
 A2 11 N/C.
 A3 9 N/C.
 A4 22 N/C.
 A5 18 N/C.
 Q0
- 10 Q0 For the two infinite dielectric sheets, see figure (5), find
 24 Q0 the magnitude of the electric field at a point P. Consider
 Q0 that each sheet has a positive surface charge density of
 Q0 10^{**2} C/m**2.
 Q0
- A1 $1.1 \cdot 10^{**13}$ N/C.
 A2 $2.2 \cdot 10^{**13}$ N/C.
 A3 $0.5 \cdot 10^{**13}$ N/C.
 A4 $1.7 \cdot 10^{**13}$ N/C.
 A5 Zero.
 Q0
- 11 Q0 A point charge of +4.0 micro-C lies at the center of a hollow
 24 Q0 spherical conducting shell that has a net charge of -13.0
 Q0 micro-C. If the inner radius of the shell is 2.0 cm and the
 Q0 outer radius is 3.0 cm, then the ratio between the charge
 Q0 density on the inner surface to the charge density on the
 Q0 outer surface is:
 Q0

A1 1 : 1.
 A2 -1 : 1.
 A3 1 : 2.
 A4 -1 : 2.
 A5 4 : 1.
 Q0
 12 Q0 A cube, as in figure (6), has an edge length of 3.00 m in a
 24 Q0 region of a uniform electric field given by the equation:
 Q0

$$\vec{E} = (-5.00 \hat{j} + 6.00 \hat{k}) \text{ N/C},$$
 Q0
 Q0 where \hat{i} , \hat{j} , and \hat{k} are the unit vectors in the directions of
 Q0 x , y , and z respectively.
 Q0 Find the electric flux through the top face (shaded).
 Q0
 A1 - 45 N*m**2/C.
 A2 45 N*m**2/C.
 A3 - 30 N*m**2/C.
 A4 30 N*m**2/C.
 A5 Zero.
 Q0
 13 Q0 The electric potential at points in the xy -plane is given by:
 25 Q0 $V = (x^3 - 2xy)$ Volts,
 Q0 where x and y are in meters. The magnitude of the electric
 Q0 field at the point with the coordinates $x = 1$ m and $y = 2$ m is:
 Q0
 A1 $\sqrt{5}$ V/m.
 A2 $\sqrt{8}$ V/m.
 A3 $\sqrt{2}$ V/m.
 A4 $\sqrt{3}$ V/m.
 A5 Zero.
 Q0
 14 Q0 In figure (7), what is the net potential at point P due to the
 25 Q0 four point charges if $V = 0$ at infinity? [take $d = 2$ cm,
 Q0 $q = 1.0$ micro-C].
 Q0
 A1 9.0×10^5 V.
 A2 -9.0×10^5 V.
 A3 4.6×10^7 V.
 A4 -4.6×10^7 V.
 A5 Zero.
 Q0
 15 Q0 Which one of the following statements is true?
 25 Q0
 Q0
 A1 The electric field lines are perpendicular to the equipotential
 A1 surfaces.
 A2 We have to do work to move a charged particle along an
 A2 equipotential surface.
 A3 The electric field is a scalar quantity.
 A4 The electric potential is a vector quantity.
 A5 Any two equipotential surfaces are always parallel.
 Q0
 16 Q0 Two balls with charges 5.0 micro-C and 10 micro-C are at a
 25 Q0 distance of 1.0 m from each other. In order to reduce the
 Q0 distance between them to 0.5 m the amount of work to be
 Q0 performed is:
 Q0
 A1 0.45 J.

- A2 45.0 J.
 A3 1.2×10^{-4} J.
 A4 4.5×10^{-4} J.
 A5 0.23 J.
 Q0
- 17 Q0 Find the equivalent capacitance of three capacitors
 Q0 connected in series. Assume the three capacitors are:
 26 Q0 $C_1 = 2.00$ micro-F, $C_2 = 4.00$ micro-F and
 Q0 $C_3 = 8.00$ micro-F.
 Q0
 A1 1.14 micro-F.
 A2 0.88 micro-F.
 A3 3.01 micro-F.
 A4 26.1 micro-F.
 A5 15.4 micro-F.
 Q0
- 18 Q0 In figure (8), find the total charge stored by the three
 Q0 capacitors if the potential difference "V" is 10.0 volts.
 26 Q0 Assume $C_1 = 10.0$ micro-F, $C_2 = 5.00$ micro-F and
 Q0 $C_3 = 4.00$ micro-F.
 Q0
 A1 31.6 micro-C.
 A2 22.1 micro-C.
 A3 61.3 micro-C.
 A4 26.1 micro-C.
 A5 63.4 micro-C.
 Q0
- 19 Q0 An air filled parallel-plate capacitor has a capacitance of
 26 Q0 1.00×10^{-12} F. The plate separation is then doubled and a
 Q0 wax dielectric is inserted, completely filling the space
 Q0 between the plates. As a result the, capacitance becomes
 Q0 2.00×10^{-12} F. The dielectric constant of the wax is:
 Q0
 A1 4.00.
 A2 0.25.
 A3 2.00.
 A4 0.50.
 A5 8.00.
 Q0
- 20 Q0 Two capacitors, C_1 and C_2 , are connected in series and a
 26 Q0 potential difference is applied to the combination. If the
 Q0 capacitor that is equivalent to the combination has the same
 Q0 potential difference, then the charge on the equivalent
 Q0 capacitors is the same as:
 Q0
 A1 The charge on C_1 or C_2 .
 A2 The sum of the charges on C_1 and C_2 .
 A3 The difference of the charges on C_1 and C_2 .
 A4 The product of the charges on C_1 and C_2 .
 A5 The ratio of the charges on C_1 and C_2 .




Figure (1)




Figure (2)




Figure (3)




Figure (4)




Figure (5)




Figure (6)




Figure (7)




Figure (8)

COURSE: PH102-2

UPM/DPC TESTING SERVICE SYSTEM

SECTION: ALL ITEM ANALYSIS ----- PHYS102 2ND MAJOR EXAM - 031

QUESTION NUMBER	CHOICE NUMBER					DISC	DIFF	EASE
	1 NOS. %	2 NOS. %	3 NOS. %	4 NOS. %	5 NOS. %			
1	123 54.9	35 15.6	29 12.9	11 04.9	26 11.6	0.44	45.08	54.92
2	130 58.0	11 04.9	5 02.2	2 00.8	76 33.9	0.36	41.96	58.04
3	52 23.2	44 19.6	35 15.6	38 16.9	55 24.5	0.19	76.78	23.22
4	53 23.6	47 20.9	82 36.6	16 07.1	26 11.6	0.39	76.33	23.67
5	67 29.9	35 15.6	22 09.8	68 30.3	32 14.2	0.48	70.08	29.92
6	80 35.8	41 18.3	52 23.3	35 15.6	15 06.7	0.60	64.12	35.88
7	136 60.7	30 13.3	19 08.4	25 11.1	14 06.2	0.47	39.28	60.72
8	55 24.6	44 19.7	46 20.6	47 21.0	31 13.9	0.11	75.33	24.67
9	109 48.6	25 11.1	46 20.5	14 06.2	30 13.3	0.34	51.33	48.67
10	79 35.4	37 16.5	39 17.4	33 14.7	35 15.6	0.34	64.57	35.43
11	49 21.8	35 15.6	47 20.9	51 22.7	42 18.7	0.29	78.12	21.88
12	82 36.6	49 21.8	19 08.4	31 13.8	43 19.1	0.34	63.39	36.61
13	87 38.8	23 10.2	31 13.8	61 27.2	22 09.8	0.56	61.16	38.84
14	95 42.4	8 03.5	25 11.1	18 08.0	78 34.8	0.49	57.58	42.42
15	109 48.6	38 16.9	24 10.7	19 08.4	34 15.1	0.46	51.33	48.67
16	100 44.6	29 12.9	20 08.9	26 11.6	49 21.8	0.39	55.35	44.65
17	161 71.8	50 22.3	7 03.1	1 00.4	5 02.2	0.46	28.12	71.88
18	125 56.3	22 09.9	14 06.3	20 09.0	41 18.4	0.31	43.69	56.31
19	47 21.0	19 08.5	105 47.0	46 20.6	6 02.6	0.14	78.92	21.08
20	105 46.8	62 27.6	11 04.9	14 06.2	32 14.2	0.42	53.12	46.88

63 <= A ACHEIVED BY 19 STUDENTS
 55 <= B ACHEIVED BY 48 STUDENTS
 45 <= C ACHEIVED BY 92 STUDENTS
 38 <= D ACHEIVED BY 37 STUDENTS
 F < 38 ACHEIVED BY 28 STUDENTS

average 4/1/16
ST.05 16.02