

- 1 Q0 Two positively charged particles q_1 and q_2 (with $q_2 > q_1$)
 22 Q0 are fixed in place on the x-axis at the positions shown
 Q0 in figure 1. A third charge q_3 is to be placed
 Q0 somewhere on the x-axis such that the net electrostatic
 Q0 force on q_3 is zero. Which one of the following
 Q0 statements is TRUE?
 Q0
 A1 q_3 should be placed at a point between q_1 and q_2 but
 A1 closer to q_1
 A2 q_3 should be placed at the mid point between q_1 and q_2 .
 A3 q_3 should be placed at a point between q_1 and q_2 but
 A3 closer to q_2 .
 A4 q_3 should be placed to the left of q_1 .
 A5 q_3 should be placed to the right of q_2 .
 Q0
- 2 Q0 Two 1.0 g spheres are charged equally and placed 2.0 cm apart.
 22 Q0 When released, each one begins to accelerate at 225 m/s^2 .
 Q0 What is the magnitude of the charge on each sphere?
 Q0
 A1 $1.0 \times 10^{-7} \text{ C}$.
 A2 $2.0 \times 10^{-7} \text{ C}$.
 A3 $3.0 \times 10^{-7} \text{ C}$.
 A4 $0.5 \times 10^{-14} \text{ C}$.
 A5 $8.0 \times 10^{-9} \text{ C}$.
 Q0
- 3 Q0 Three charges $+2.00 \times 10^{-8} \text{ C}$, $+2.00 \times 10^{-8} \text{ C}$, and
 23 Q0 $-4.00 \times 10^{-8} \text{ C}$ are respectively arranged at the
 Q0 corners F, G, and H of a right-angle triangle as shown
 Q0 in figure 2. Find the magnitude and direction of the
 Q0 resultant electric field at point P due to the three charges.
 Q0
 A1 $2.88 \times 10^3 \text{ N/C}$ towards H.
 A2 $5.37 \times 10^3 \text{ N/C}$ towards H.
 A3 $5.37 \times 10^3 \text{ N/C}$ away from H.
 A4 $1.09 \times 10^5 \text{ N/C}$ towards F.
 A5 $2.88 \times 10^3 \text{ N/C}$ away from H.
 Q0
- 4 Q0 In figure 9, a small ball of mass $m = 2.0 \text{ g}$ is hanging from
 23 Q0 a fixed point by a non-conducting string of length 1.00 m.
 Q0 The ball carries a charge $q = 25.0 \times 10^{-9} \text{ C}$. The mass of
 Q0 the string is negligible. An electric field E with magnitude
 Q0 $E = 2.0 \times 10^5 \text{ N/C}$, in the positive x-direction, causes the
 Q0 ball to be in an equilibrium position with an angle θ .
 Q0 Find the angle θ . [Take $g = 9.80 \text{ m/s}^2$].
 Q0
 A1 14.3 degrees.
 A2 10.0 degrees.
 A3 7.1 degrees.
 A4 0.2 degrees.
 A5 75.7 degrees.
 Q0
- 5 Q0 A uniform electric field is set up between two large
 23 Q0 charged plates, see Figure 3. An electron is released
 Q0 from the negatively charged plate, and at the same time,
 Q0 a proton is released from the positively charged plate.
 Q0 They cross each other at a distance of $5.00 \times 10^{-6} \text{ m}$
 Q0 from the positively charged plate. If only the field due
 Q0 to the charged plates is considered, find the distance

- Q0 between the two plates. [Take the ratio
 Q0 mass of the electron : mass of the proton = 1 : 1833]
 Q0
 A1 9.19 mm.
 A2 11.3 mm.
 A3 2.34 mm.
 A4 7.77 mm.
 A5 14.6 mm.
 Q0
- 6 Q0 A very long uniform line of charge having a linear charge
 24 Q0 density of 6.8 micro-C/m lies along x-axis. A second line
 Q0 of charge has a linear charge density of -3.40 micro-C/m
 Q0 and is parallel to x-axis at $y = 0.5$ m. What is the net
 Q0 electric field at point where $y = 0.25$ m on y-axis?
 Q0
 A1 7.3×10^5 N/C along +y-axis.
 A2 4.8×10^6 N/C along +y-axis.
 A3 4.8×10^4 N/C along -y-axis.
 A4 3.4×10^6 N/C along +y-axis.
 A5 7.3×10^2 N/C along -y-axis.
 Q0
- 7 Q0 Which of the following statements are CORRECT:
 24 Q0
 Q0 (1) The electric flux through a Gaussian surface depends on
 Q0 the shape of the surface.
 Q0 (2) The electric flux through a closed surface depends on
 Q0 the net charge enclosed by the surface.
 Q0 (3) The electric field inside a uniformly charged solid
 Q0 conducting sphere in electrostatic equilibrium is zero.
 Q0 (4) The electric potential inside a uniformly charged solid
 Q0 conducting sphere in electrostatic equilibrium is zero.
 Q0
 A1 2 and 3 only.
 A2 1 and 2 only.
 A3 1, 2, 3, and 4.
 A4 3 and 4 only.
 A5 4 only.
 Q0
- 8 Q0 The net electric flux passing through a closed surface
 24 Q0 is -4.00×10^2 N·m²/C. What is net electric charge
 Q0 contained inside the surface if the surface is a cylinder
 Q0 of height 3.52 cm and radius 1.12 cm.
 Q0
 A1 $-3.54 \times 10^{(-9)}$ C.
 A2 $-1.00 \times 10^{(-2)}$ C.
 A3 $3.54 \times 10^{(-9)}$ C.
 A4 $1.00 \times 10^{(-2)}$ C.
 A5 zero.
 Q0
- 9 Q0 A positive point charge q sits at the center of a hollow
 Q0 spherical shell. The shell, with radius R and negligible
 24 Q0 thickness, has net charge $-2q$. The electric field strength
 Q0 outside the spherical shell (at $r > R$) will be:
 Q0
 A1 $k \cdot q / r^2$ radially inwards.
 A2 $k \cdot q / r^2$ radially outwards.
 A3 $3 \cdot k \cdot q / r^2$ radially inwards.
 A4 $3 \cdot k \cdot q / r^2$ radially outwards.

- A5 zero.
Q0
- 10 Q0 A charged, isolated, large non-conducting plate is placed
24 Q0 on the XY-plane. At 1.5 m from the plate, on Z-axis, the
Q0 electric field measured was 10^{*4} N/C and directed into
Q0 the plate. What is the charge density on the plate?
Q0
- A1 $-1.8*10^{*(-7)}$ C/m**2.
A2 $1.8*10^{*(-7)}$ C/m**2.
A3 $-3.2*10^{*(-7)}$ C/m**2.
A4 $3.2*10^{*(-7)}$ C/m**2.
A5 zero.
Q0
- 11 Q0 Two oppositely charged parallel plates, 0.02 m apart, produce
Q0 a uniform electric field between the plates. The potential
Q0 energy U(J) of an electron in the field varies with
25 Q0 displacement x(m) from one of the plates as shown in figure 5.
Q0 What is the magnitude of the force on the electron?
Q0
- A1 $7.5*10^{*(-15)}$ N.
A2 $3.0*10^{*(-18)}$ N.
A3 $6.0*10^{*(-20)}$ N.
A4 $1.5*10^{*(-15)}$ N.
A5 zero.
Q0
- 12 Q0 A point charge Q, at the center of a circle, is surrounded
Q0 by six charges each of magnitude q at a distance r as shown
Q0 in figure 4. How much work is done by an external agent to
25 Q0 remove the charge Q from the center to infinity?
Q0 [Consider the electrostatic potential at infinity = 0]
Q0
- A1 zero.
A2 $k*6*Q*q/r^{*2}$.
A3 $k*6*q/r$.
A4 $k*6*q/r^{*2}$.
A5 $k*3*Q*q/r$.
Q0
- 13 Q0 Two protons, P, are fixed 6.0 m apart, as shown in
25 Q0 figure 7. An electron, e, is released from point A. Find
Q0 its speed at point O, midway between the protons.
Q0
- A1 11.6 m/s.
A2 24.0 m/s.
A3 121 m/s.
A4 2.4 m/s.
A5 0.1 m/s.
Q0
- 14 Q0 Figure 6 shows three points X, Y and Z forming an equilateral
Q0 triangle of side S in a uniform electric field of strength E.
25 Q0 A unit positive test charge is moved from X to Y, then from
Q0 Y to Z, and from Z back to X. Which one of the following
Q0 correctly gives the work done by an external agent in
Q0 moving the charge along the various parts of the path?
Q0
- A1 0, $-E*S*\sin(60 \text{ degrees})$, + $E*S*\sin(60 \text{ degrees})$.
A2 0, $-E*S*\cos(60 \text{ degrees})$, + $E*S*\cos(60 \text{ degrees})$.
A3 $E*S$, $-E*S*\sin(60 \text{ degrees})$, + $E*S*\cos(60 \text{ degrees})$.
A4 0, $-E*S*\cos(60 \text{ degrees})$, + $E*S*\sin(60 \text{ degrees})$.

A5 $-E*S, -E*S*\tan(60 \text{ degrees}), + E*S*\sin(60 \text{ degrees})$.
 Q0

15 Q0 Over a certain region of space, the electric potential
 25 Q0 is give by:
 Q0 $V(x,y) = x**2 + y**2 + 2*x*y$.
 Q0 Find the angle that the electric field vector makes with
 Q0 Z-axis at the point P(1.0,2.0,0.0)
 Q0

A1 90 degrees.
 A2 0 degrees.
 A3 45 degrees.
 A4 75 degrees.
 A5 60 degrees.
 Q0

16 Q0 Consider two separate capacitors: $c_1=30$ micro-F carries a
 Q0 charge of $q_1=6.0*10**2$ micro-C and $c_2=50$ micro-F, carries
 26 Q0 a charge of $q_2=1.0*10**3$ micro-C. If the opposite polarity
 Q0 terminals of the two capacitors are connected together as
 Q0 shown in figure 10, find the new voltage across c_1 .
 Q0

A1 5.0 Volts.
 A2 10 Volts.
 A3 15 Volts.
 A4 3.8 Volts.
 A5 2.2 Volts.
 Q0

17 Q0 A 25 micro-F parallel plates capacitor is constructed using
 Q0 Pyrex glass as a dielectric. If the thickness of the Pyrex
 26 Q0 glass sheet is doubled, calculate the new capacitance of the
 Q0 capacitor. (Dielectric constant of Pyrex Glass = 5.6)
 Q0

A1 12.5 micro-F.
 A2 30.2 micro-F.
 A3 100 micro-F.
 A4 50.0 micro-F.
 A5 6.25 micro-F.
 Q0

18 Q0 Three capacitors $C_1=5$ micro-F, $C_2=10$ micro-F and $C_3= 3$ micro-F
 26 Q0 are connected to a 20 V battery as shown in Figure 8. Find
 Q0 the stored electric energy in C_2 .
 Q0

A1 $2.2*10**(-4)$ J.
 A2 $0.3*10**(-4)$ J.
 A3 $4.0*10**(-6)$ J.
 A4 $1.3*10**(-4)$ J.
 A5 $1.0*10**(-5)$ J.
 Q0

19 Q0 A 500 W electric heater is designed to operate from a 120-V
 27 Q0 power supply. The line voltage decreases and the heater takes
 Q0 only 459 W. Find the voltage drop in the line voltage
 Q0 (Assuming the resistance is constant).
 Q0

A1 5 Volts.
 A2 10 Volts.
 A3 15 Volts.
 A4 3 Volts.
 A5 2 Volts.
 Q0

20 Q0 What diameter must a copper wire have if its resistance is
Q0 to be the same as that of an equal length of an aluminum wire
27 Q0 with 3.26 mm diameter?

Q0 [Resistivity of aluminum = 2.75×10^{-8} Ohm.m;

Q0 Resistivity of copper = 1.69×10^{-8} Ohm.m;

Q0

A1 2.6 mm.

A2 8.3 mm.

A3 10 mm.

A4 4.0 mm.

A5 3.3 mm.

Figures of Phys 102 Second Major 042













Figure 10