

**Q1.**

When a large object moves in air, there is a resistive force on it whose magnitude is given by:  $F = 0.5 D \rho B v^2$ , where  $D$  is a dimensionless constant,  $\rho$  is the density of the air and  $v$  is the speed of the object. What are the dimensions of  $B$  ?

- A)  $L^2$
- B)  $M^2$
- C)  $T^2$
- D)  $ML^2$
- E)  $TM^2$

**Ans:**  $[B] = \left[ \frac{F}{\rho v^2} \right] = \left( \text{kg} \cdot \frac{\text{m}}{\text{s}^2} \right) \cdot \left( \frac{\text{kg}}{\text{m}^3} \right)^{-1} \cdot \left( \frac{\text{m}^2}{\text{s}^2} \right)^{-1}$

$$= \frac{\cancel{\text{kg}} \cdot \text{m} \cdot \cancel{\text{m}^3} \cdot \cancel{\text{s}^2}}{\cancel{\text{s}^2} \cdot \cancel{\text{kg}} \cdot \text{m}^2} = \text{m}^2 \rightarrow L^2$$

**Q2.**

A cubic box has a side of length 1.00 ft. What is the volume of the box in cubic meters? (1 ft = 12.0 inch, 1 inch = 2.54 cm)

- A) 0.0283
- B) 0.843
- C) 0.759
- D) 0.227
- E) 0.00100

**Ans:**

$$l = 1 \text{ ft} \cdot \frac{12 \cancel{\text{in}}}{1 \cancel{\text{ft}}} \cdot \frac{2.54 \cancel{\text{cm}}}{1 \cancel{\text{in}}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} = 0.3048 \text{ m}$$

$$V = l^3 = (0.3048)^3 = 0.0283 \text{ m}^3$$

**Q3.**

A car is travelling along a straight line. It travels at 40.0 km/h for 2.00 h, then at 50.0 km/h for 1.00 h, and finally at 20.0 km/h for 0.500 h. What is the average speed of the car?

- A) 40.0 km/h
- B) 36.7 km/h
- C) 55.0 km/h
- D) 45.0 km/h
- E) 31.6 km/h

**Ans:**

$$\left. \begin{aligned} d_1 &= 40 \times 2 = 80 \text{ km} \\ d_2 &= 50 \times 1 = 50 \text{ km} \\ d_3 &= 20 \times \frac{1}{2} = 10 \text{ km} \end{aligned} \right\} d = 80 + 50 + 10 = 140 \text{ km}$$

$$t = 2 + 1 + \frac{1}{2} = 3.5 \text{ h}$$

$$S_{avg} = \frac{d}{t} = \frac{140}{3.5} = 40 \text{ km/h}$$

**Q4.**

In one dimensional motion, a particle is 5.00 m east of the origin and is moving west with a speed of 2.00 m/s. Five seconds later, it is 11.0 m east of the origin. What is its acceleration? Assume that the acceleration is constant.

- A) 1.28 m/s<sup>2</sup>
- B) -0.320 m/s<sup>2</sup>
- C) 2.08 m/s<sup>2</sup>
- D) 1.68 m/s<sup>2</sup>
- E) 0.781 m/s<sup>2</sup>

**Ans:**  $x_1 = +5 \text{ m}; v_1 = -2 \text{ m/s}$

$x_2 = +11 \text{ m}; t = 5.0 \text{ s}$

$$x_2 - x_1 = v_1 t + \frac{1}{2} a t^2$$

$$11 - 5 = -10 + 12.5 a \Rightarrow a = \frac{16}{12.5} = 1.28 \text{ m/s}^2$$

**Q5.**

An object is thrown vertically upward from the roof of a building that is 50 m high. It rises to a maximum height of 10 m above the roof (**Figure 1**). When is it 20 m below the roof?

- A) 3.9 s
- B) 5.2 s
- C) 4.9 s
- D) 3.2 s
- E) 6.1 s

**Ans:** Throw  $\rightarrow$  max height:  $v^2 = v_0^2 - 2g(y - y_0)$

$$0 = v_0^2 - 19.6(60 - 50) \Rightarrow v_0 = 14 \text{ m/s}$$


Throw  $\rightarrow$  requested point:

$$y - y_0 = v_0 t - \frac{1}{2} g t^2$$

$$30 - 50 = 14t - 4.9t^2$$

$$4.9 t^2 - 14t - 20 = 0$$

$$t = \frac{14 \pm \sqrt{196 + (4 \times 20 \times 4.9)}}{9.8} = 3.9 \text{ s}$$



**Q6.**

At time  $t = 3.0$  s, the velocity of a particle, moving along the  $x$ -axis with constant acceleration, is  $v = +4.0$  m/s. At  $t = 7.0$  s, its velocity is  $v = -12$  m/s. Find the velocity at  $t = 0$ .

- A) + 16 m/s
- B) + 20 m/s
- C) - 16 m/s
- D) - 20 m/s
- E) + 5.0 m/s

**Ans:**  $v = v_0 + at$


$$\left. \begin{array}{l} \text{(i)} \quad 4 = v_0 + 3a \\ \text{(ii)} \quad -12 = v_0 + 7a \end{array} \right\}$$

$$\left. \begin{array}{l} \text{(i)} \times -7: -28 = -7v_0 - 21a \\ \text{(ii)} \times 3: -36 = 3v_0 + 21a \end{array} \right\} \rightarrow \overset{\oplus}{-64} = -4v_0 \Rightarrow v_0 = +16 \text{ m/s}$$

**Q7.**

For the three vectors ( $\vec{A}, \vec{B}, \vec{C}$ ) shown in **Figure 2**, find  $\vec{C} \times (\vec{A} \times \vec{B})$ .

- A)  $-3.19 \times 10^4 \hat{i}$
- B)  $-6.27 \times 10^4 \hat{i}$
- C)  $+2.67 \times 10^4 \hat{i}$
- D)  $+3.67 \times 10^4 \hat{i}$
- E) zero



**Ans:**  $\phi$  angle between  $\vec{A}, \vec{B} = 118^\circ$

$$|\vec{A} \times \vec{B}| = 44 \times 26.5 \times \sin 118^\circ = 1029.5 \rightarrow \hat{k}$$

$\vec{C} \times (\vec{A} \times \vec{B})$  is along  $(-x)$  by right-hand rule

$$|\vec{C} \times (\vec{A} \times \vec{B})| = 31.0 \times 1029.5 \times \sin 90^\circ = 3.19 \times 10^4$$

**Q8.**

What is the angle between two vectors  $\vec{A} = 20\hat{i}$  and  $\vec{B} = -25\hat{i} + 30\hat{j}$ ?

- A) 130°
- B) 150°
- C) 160°
- D) 140°
- E) 110°

**Ans:**  $\vec{A}$  is along the  $x$  – axis

Angle between  $\vec{B}$  and is (+)  $x$  axis is  $\phi = 180 - \tan^{-1}\left(\frac{30}{25}\right) = 130^\circ$

**Q9.**

A body moves from a position with coordinates (1.0, 2.0) m to (– 4.0, 2.0) m. Its displacement vector is given by

- A) 5.0 m at 180°
- B) 5.0 m at 135°
- C) 1.7 m at 297°
- D) 5.0 m at 0°
- E) 5.2 m at 108°

**Ans:**

$$\left. \begin{array}{l} \vec{r}_1 = \hat{i} + 2\hat{j} \\ \vec{r}_2 = -4\hat{i} + 2\hat{j} \end{array} \right\} \vec{d} = \vec{r}_2 - \vec{r}_1 = -5\hat{i} \text{ (m)}$$

$\Rightarrow \vec{d}$  makes an angle of 180° and has a magnitude of 5.0 m.

**Q10.**

At time  $t = 0$ , a particle leaves the origin with a velocity of 9.0 m/s in the positive  $y$ -direction and moves in the  $xy$  plane with a constant acceleration of  $(2.0\hat{i} - 4.0\hat{j}) \text{ m/s}^2$ .

What is the speed of the particle when its  $x$ -coordinate is + 15 m?

- A) 10 m/s
- B) 16 m/s
- C) 12 m/s
- D) 14 m/s
- E) 26 m/s

**Ans:**

$$x = v_{0x}t + \frac{1}{2}a_x t^2$$

$$15 = 0 + \left(\frac{1}{2} \times 2\right)t^2 \Rightarrow t = \sqrt{15} = 3.87 \text{ s}$$

$$v_x = v_{0x} + a_x t = 0 + (2)(\sqrt{15}) = 7.75 \text{ m/s}$$

$$v_y = v_{0y} + a_y t = 9 + (-4)(\sqrt{15}) = -6.49 \text{ m/s}$$

$$\text{Speed} = \sqrt{v_x^2 + v_y^2} = 10 \text{ m/s}$$

**Q11.**

A projectile is launched from the ground at time  $t = 0$ . When  $t = 5.00$  s, its velocity is given as:  $\vec{v} = 25.0\hat{i} - 30.0\hat{j}$  (m/s). Find the maximum height of the projectile.

A) 18.4 m

B) 15.6 m

C) 32.9 m

D) 65.8 m

E) 49.4 m

**Ans:**

$$v_y = v_{0y} - gt$$

$$-30 = v_{0y} - (9.8 \times 5) \Rightarrow v_{0y} = 19 \text{ m/s}$$

launch  $\rightarrow$  max. height:

$$v_y^2 = v_{0y}^2 - 2g(y - y_0)$$

$$0 = (19)^2 - (19.6)(H - 0)$$

$$\Rightarrow H = 18.4 \text{ m}$$

**Q12.**

A river, of width 150 m, flows with a uniform speed of 4.0 m/s toward the east. It takes 20 s for a boat to cross the river from a point on the south side to the opposite point on the north side. What is the speed of the boat relative to the water?

A) 8.5 m/s

B) 9.1 m/s

C) 5.7 m/s

D) 7.0 m/s

E) 6.4 m/s


**Ans:**

$r \rightarrow$  river,  $g \rightarrow$  ground,  $b \rightarrow$  boat

$$v_{rg} = 4.0 \text{ m/s}$$

$$v_{bg} = \frac{150}{20} = 7.5 \text{ m/s}$$

$$v_{br} = \sqrt{v_{bg}^2 + v_{rg}^2} = \sqrt{(4)^2 + (7.5)^2} = 8.5 \text{ m/s}$$



**Q13.**

A particle undergoes counterclockwise uniform circular motion around a circle of radius 5.0 m with a period of 3.2 s, as shown in **Figure 3**. In a quarter of a period, as the particle moves from A to B, what is the magnitude of the average velocity of the particle?

- A) 8.8 m/s
- B) 9.8 m/s
- C) 4.9 m/s
- D) 7.8 m/s
- E) zero


**Ans:**

$$\left. \begin{aligned} \vec{r}_i &= 5.0\hat{i} \\ \vec{r}_f &= 5.0\hat{j} \end{aligned} \right\} \Delta\vec{r} = 5.0\hat{j} - 5.0\hat{i} \text{ (m)}$$

$$\Delta t = \frac{T}{4} = \frac{3.2}{4} = 0.8 \text{ s}$$

$$\vec{v}_{avg} = \frac{\Delta\vec{r}}{\Delta t} = \frac{5.0\hat{j} - 5.0\hat{i}}{0.8} = -6.25\hat{i} + 6.25\hat{j} \text{ (m/s)}$$

$$v_{avg} = \sqrt{2} \times 6.25 = 8.8 \text{ m/s}$$



**Q14.**

In **Figure 4**, the force  $\vec{F}$  acts to move the two blocks on a horizontal frictionless surface. Find the magnitude of the tension in the massless connecting string. Take  $F = 15 \text{ N}$ ,  $M = 1.5 \text{ kg}$ .

- A) 4.3 N
- B) 5.9 N
- C) 5.1 N
- D) 3.8 N
- E) 6.2 N

**Ans:**

$$2M: 2Ma = F\cos\theta - T \rightarrow (1)$$

$$M: Ma = T \rightarrow (2)$$


Adding(1) and (2) we get,  $3Ma = F\cos\theta$

$$\Rightarrow a = \frac{F\cos\theta}{3M}$$

Back to (1):

$$T = F\cos\theta - 2Ma = F\cos\theta - (2M)\left(\frac{F\cos\theta}{3M}\right)$$


$$= \frac{1}{3}F\cos\theta = \frac{1}{3} \times 15 \times \cos 30^\circ = 4.3 \text{ N}$$



**Q15.**


The two blocks shown in **Figure 5** are released from rest and are observed to have an acceleration of  $1.5 \text{ m/s}^2$ . What is the magnitude of the frictional force on the block that slides horizontally? Assume the pulleys and strings are massless, and take  $M = 1.4 \text{ kg}$ .

- A) 21 N
- B) 46 N
- C) 11 N
- D) 37 N
- E) 29 N

**Figure 5****Ans:**

$$\left. \begin{array}{l} Ma = T - f_k \\ 2Ma = 2Mg - T \end{array} \right\} \oplus \rightarrow 3Ma = 2Mg - f_k$$

$$\Rightarrow f_k = (2g - 3a)M = (19.6 - 4.5)(1.4) = 21 \text{ N}$$

**Q16.**

An 80-kg man stands in an elevator that has a downward acceleration of  $1.5 \text{ m/s}^2$ . The magnitude of the force exerted by the man on the floor of the elevator is

- A) 664 N
- B) 784 N
- C) 904 N
- D) 388 N
- E) zero

**Ans:**


$$-Ma = F_N - Mg$$

$$\Rightarrow F_N = M(g - a) = (80)(9.8 - 1.5) = 664 \text{ N}$$

**Q17.**

Two blocks with masses  $M_1 = 3.0 \text{ kg}$  and  $M_2 = 5.0 \text{ kg}$  are connected by a light rope and move on a frictionless surface, as shown in **Figure 6**. A force  $F = 20 \text{ N}$  acts on  $M_2$  as shown in the figure. Find the magnitude of the acceleration of the system.

- A)  $0.66 \text{ m/s}^2$
- B)  $2.6 \text{ m/s}^2$
- C)  $1.8 \text{ m/s}^2$
- D)  $4.3 \text{ m/s}^2$
- E)  $0.86 \text{ m/s}^2$



**Ans:**

$$M_1 a = T - M_1 g \sin \theta \rightarrow (1)$$

$$M_2 a = F - T \rightarrow (2)$$

Adding (1) and (2) we get

$$(M_1 + M_2) a = F - M_1 g \sin \theta$$

$$a = \frac{F - M_1 g \sin \theta}{M_1 + M_2} = \frac{20 - \left(3 \times 9.8 \times \frac{1}{2}\right)}{8} = 0.66 \text{ m/s}^2$$

**Q18.**

A car moves on a level horizontal road in a circle of radius  $40 \text{ m}$ . The coefficient of static friction between tires and road is  $0.50$ . The maximum speed with which this car can round this curve without sliding is

- A)  $14 \text{ m/s}$
- B)  $12 \text{ m/s}$
- C)  $16 \text{ m/s}$
- D)  $10 \text{ m/s}$
- E)  $18 \text{ m/s}$

**Ans:**

$$m a = F_{\text{net}}$$

$$\frac{m v^2}{R} = f_s$$

$$\frac{m v_{\text{max}}^2}{R} = f_{s, \text{max}} = \mu_s F_N = \mu_s m g$$

$$\Rightarrow v_{\text{max}} = \sqrt{\mu_s \cdot R \cdot g} = \sqrt{0.5 \times 40 \times 9.8} = 14 \text{ m/s}$$



**Q19.**Which of the following statements is **TRUE**?

- A) A particle can be in equilibrium and yet moving.
- B) A stone that has been thrown vertically upward reverses its acceleration as it reaches the top of its trajectory.
- C) Two vectors of unequal magnitudes can add up to zero.
- D) On a displacement-time graph, a straight line with positive slope indicates motion at increasing speed.
- E) The action and reaction forces act on the same object.

**Ans:****A**

---

**Q20.**Which of the following statements is **TRUE**?

- A) A car can be accelerating while moving at constant speed.
- B) If an object is released from rest, it falls 9.8 m during the first second of its motion.
- C) The velocity of a projectile equals its initial velocity added to a constant horizontal velocity.
- D) A particle can move with uniform velocity along a circular path.
- E) The velocity of a projectile at the top of its trajectory is zero.

**Ans:****A**

---