

# Chapter 1: Equations and Inequalities

## Section 1.1 Linear and Absolute Value Equations

### Linear Equations

Solve.

- $3x - 5 = x + 9$  (a)  $\frac{1}{7}$  (b)  $-7$  (c)  $-\frac{1}{7}$  (d)  $7$
- $-\frac{8}{7}x = 7$  (a)  $0$  (b)  $-8$  (c)  $-\frac{49}{8}$  (d)  $\frac{1}{7}$
- $x + \frac{2}{11} = \frac{5}{8}$  (a)  $\frac{3}{88}$  (b)  $\frac{39}{88}$  (c)  $\frac{39}{8}$  (d)  $\frac{39}{11}$
- $1 = 3(x - 1) + 2 - 2x$  (a)  $0$  (b)  $6$  (c)  $2$  (d)  $4$

### Contradictions, Conditional Equations, and Identities

5. Which of the following equations is not an identity?

- (a)  $\frac{5x - 15}{5} = x - 3$  (b)  $5(2x - 3) - x = 9x - 15$  (c)  $5x + 2 = -3$  (d) All of the above are identities

6. Which of the following equations is an identity?

- (a)  $4(x - 2) = 4x - 8$  (b)  $4(x - 2) = 8 - 4x$  (c)  $4(x - 2) = 4x - 2$  (d)  $4(x - 2) = x - 8$

7. Which of the following equations is not a conditional equation?

- (a)  $-3x + 8(x + 5) = 5x + 5$  (b)  $-15x + 5 = -3(x + 8) + 8x$  (c)  $-15x - 3 = 8(x + 5)$   
(d)  $-3x + 5 = 5 + 5x$  (e) None of these

8. Find the values of  $a$  and  $c$  that make the equation an identity.

$$9x - 12 = 3(ax + c) - 6x$$

- (a)  $a = -5, c = -13$  (b)  $a = 5, c = -4$  (c)  $a = 12, c = -12$  (d)  $a = 12, c = -4$  (e) None of these

### Absolute Value Equations

Solve.

- $|3x - 1| = 2$  (a)  $\frac{4}{3}, \frac{2}{3}$  (b)  $1, -\frac{1}{3}$  (c)  $-\frac{4}{3}, \frac{2}{3}$  (d)  $-1, -\frac{1}{3}$
- $\left|\frac{5}{3}x + 7\right| + 3 = 13$  (a)  $\frac{69}{5}, \frac{51}{5}$  (b)  $\frac{9}{5}, -\frac{51}{5}$  (c)  $\frac{51}{5}, -\frac{85}{3}$  (d)  $5, -\frac{69}{5}$
- $|4x + 4| = 3$  (a)  $-\frac{7}{4}, -\frac{1}{4}$  (b)  $-\frac{7}{4}, \frac{7}{4}$  (c)  $\frac{3}{2}, \frac{7}{4}$  (d)  $-\frac{7}{4}, \frac{1}{4}$

12. What is the solution to the equation  $2|x| + 1 = 10$ ?

- (a)  $x = \pm 6$                       (b)  $x = \pm 6\frac{1}{2}$                       (c)  $x = \pm 5\frac{1}{2}$                       (d)  $x = \pm 4\frac{1}{2}$

### Applications

13. The formula

$$C = \frac{5}{9}(F - 32)$$

is used to convert degrees Fahrenheit,  $F$ , to degrees Celsius,  $C$ . Convert  $-1^\circ C$  to degrees Fahrenheit.

- (a)  $55.8^\circ F$                       (b)  $-37.0^\circ F$                       (c)  $30.2^\circ F$                       (d)  $-18.3^\circ F$

14. Use the formula  $d = rt$ , where  $d$  is the distance,  $r$  is the rate of speed, and  $t$  is the time, to find how long it would take to travel a distance of 200 miles at a speed of 50 miles per hour.

- (a) 4 hr                      (b) 7 hr                      (c) 5 hr                      (d) 16 hr

15. The sum of the angles in a convex polygon can be determined by the formula

$$D = 180(n - 2)$$

where  $D$  is the total number of degrees in the angles of a convex polygon with  $n$  sides. If a convex polygon's angles have a sum of  $720^\circ$ , find the number of sides in the polygon.

- (a) 7                      (b) 4                      (c) 6                      (d) 8                      (e) None of these

16. The charge for mailing a fourth-class package through the U.S. Postal Service is

$$C = 0.05x + 2.57$$

where  $C$  is the charge in dollars and  $x$  is the weight of the package in pounds.

- (a) Find the charge to mail a package that weighs 10 pounds.  
 (b) How many pounds can be mailed for \$2.97?

- (a) (a) \$0.40                      (b) (a) \$2.92                      (c) (a) \$3.07                      (d) (a) \$3.07  
 (b) 10 lb                      (b) 7 lb                      (b) 8 lb                      (b) 9 lb

### Section 1.2 Formulas and Applications

#### Formulas

Solve the formula for the given variable.

17.  $V = \pi r^2 h$  for  $h$                       (a)  $\pi h = Vr^2$                       (b)  $h = \frac{V}{\pi r^2}$                       (c)  $h = V - \pi r^2$                       (d) none of these

18.  $W = p(V_2 - V_1)$  for  $p$                       (a)  $p = \frac{W}{V_2 - V_1}$                       (b)  $pW = V_2 - V_1$                       (c)  $p = \frac{W}{V_2} + V_1$                       (d)  $p = W(V_2 - V_1)$

19.  $M = \frac{mRT}{PV}$  for  $R$                       (a)  $R = MPV - mT$                       (b)  $R = \frac{MPV}{mT}$                       (c)  $R = \frac{mT}{MPV}$                       (d)  $R = \frac{MmT}{PV}$

20. Solve the equation  $A = \frac{1}{2}h(b+c)$  for  $c$ .

- (a)  $c = \frac{A}{b+c}$                       (b)  $c = \frac{h}{2}A - c$                       (c)  $c = \frac{2}{A}h + b$                       (d)  $c = \frac{2}{h}A - b$

**Applications**

21. The perimeter of a triangle is 60 centimeters. One side is 6 centimeters shorter than the second side. The third side is 6 centimeters shorter than triple the length of the first side. Find the length of each side.
- (a) 12 cm, 12 cm, 12 cm    (b) 6 cm, 12 cm, 35 cm    (c) 12 cm, 18 cm, 30 cm    (d) 6 cm, 12 cm, 42 cm
22. Jacob has \$6.00 in dimes and quarters. He has twice as many quarters as dimes. How many of each coin does he have?
- (a) 10 dimes and 20 quarters    (b) 8 dimes and 56 quarters  
(c) 8 dimes and 16 quarters    (d) 20 dimes and 10 quarters
23. The daily cost of renting a car is \$30 plus \$0.40 per mile. If Jane paid \$115.60 for a 1-day rental, how many miles did Jane travel?
- (a) 364    (b) 289    (c) 86    (d) 214
24. The sum of three consecutive odd integers is 399. What is the largest of the three integers?
- (a) 133    (b) 134    (c) 135    (d) None of these

**Section 1.3 Quadratic Equations****Solve Quadratic Equations by Factoring**

Solve by factoring.

25.  $x^2 - x = 20$     (a) -5, 4    (b) 5, 4    (c) -5, -4    (d) 5, -4
26.  $2x^2 + 13x + 15 = 0$     (a)  $\frac{3}{2}$ , -5    (b)  $-\frac{3}{2}$ , 5    (c)  $\frac{3}{2}$ , 5    (d)  $-\frac{3}{2}$ , -5
27.  $x^2 - 5x - 6 = 0$     (a) 1, -6    (b) 6, -1    (c) 2, -3    (d) 3, -2
28. Solve.    (a)  $-\frac{3}{2}$ , 2    (b)  $-\frac{2}{3}$ ,  $\frac{1}{2}$     (c)  $\frac{2}{3}$ ,  $-\frac{1}{2}$     (d)  $\frac{3}{2}$ , -2
- $$\frac{x^2}{2} + \frac{x}{4} = \frac{3}{2}$$

**Solve Quadratic Equations by Taking Square Roots**

29. Solve.    (a)  $5\sqrt{2}$     (b)  $-5\sqrt{2}$ ,  $5\sqrt{2}$     (c) -25, 25    (d)  $-2\sqrt{5}$ ,  $2\sqrt{5}$
- $$x^2 - 50 = 0$$
30. Solve by taking the square root.  $9x^2 = 900$     (a)  $\pm\sqrt{8100}$     (b)  $\pm 90$     (c)  $\pm 10$     (d)  $\pm\sqrt{891}$
31. Solve.    (a)  $-\frac{5\sqrt{8}}{8}$ ,  $\frac{5\sqrt{8}}{8}$     (b)  $\frac{8\sqrt{5}}{5}$     (c)  $-\frac{8\sqrt{5}}{5}$ ,  $\frac{8\sqrt{5}}{5}$     (d)  $\frac{8}{5}$
- $$5x^2 = 64$$
32. Solve by extracting square roots.
- $$4(x+4)^2 - 108 = 0$$
- (a)  $-4 \pm 3\sqrt{3}$     (b)  $-4 \pm 9\sqrt{27}$     (c)  $-4 \pm 9\sqrt{3}$     (d)  $4 \pm 3\sqrt{3}$

**Solve Quadratic Equations by Completing the Square**

33. Solve by completing the square.  $-9x = 5x^2 - 1$
- (a)  $\frac{-9 \pm \sqrt{101}}{10}$  (b)  $\frac{9 \pm \sqrt{101}}{10}$  (c)  $\frac{-9 \pm \sqrt{61}}{10}$  (d)  $\frac{9 \pm \sqrt{61}}{10}$

Solve by completing the square.

34.  $4x^2 - 6x - 6 = 0$

(a)  $\frac{3+2\sqrt{33}}{4}, \frac{3-2\sqrt{33}}{4}$

(b)  $\frac{3+\sqrt{33}}{4}, \frac{3-\sqrt{33}}{4}$

(c)  $\frac{-3+2\sqrt{33}}{4}, \frac{-3-2\sqrt{33}}{4}$

(d)  $\frac{-3+\sqrt{33}}{4}, \frac{-3-\sqrt{33}}{4}$

35.  $-9x^2 - 90x = 18$

(a)  $5+\sqrt{23}, 5-\sqrt{23}$  (b)  $5+3\sqrt{3}, 5-3\sqrt{3}$  (c)  $-5+3\sqrt{3}, -5-3\sqrt{3}$  (d)  $-5+\sqrt{23}, -5-\sqrt{23}$

36. Solve by completing the square.  $x^2 + 8x - 20 = 0$
- (a)  $-8, 20$  (b)  $10, -2$  (c)  $8, -20$  (d)  $-10, 2$

**Solve Quadratic Equations by Using the Quadratic Formula**

Solve using the quadratic formula.

37.  $2x^2 + 1 = 5x$

(a)  $\frac{5+\sqrt{33}}{4}, \frac{5-\sqrt{33}}{4}$  (b)  $\frac{5+\sqrt{17}}{4}, \frac{5-\sqrt{17}}{4}$  (c)  $\frac{-5+\sqrt{33}}{4}, \frac{-5-\sqrt{33}}{4}$  (d)  $\frac{-5+\sqrt{17}}{4}, \frac{-5-\sqrt{17}}{4}$

38.  $4x^2 - 4x - 5 = 0$

(a)  $\frac{-1+2\sqrt{6}}{2}, \frac{-1-2\sqrt{6}}{2}$  (b)  $\frac{-1+\sqrt{6}}{2}, \frac{-1-\sqrt{6}}{2}$  (c)  $\frac{1+\sqrt{6}}{2}, \frac{1-\sqrt{6}}{2}$  (d)  $\frac{1+2\sqrt{6}}{2}, \frac{1-2\sqrt{6}}{2}$

39. Use the Quadratic Formula to solve.  $2x^2 - 1 = 7x$
- (a)  $\frac{-7 \pm \sqrt{41}}{4}$  (b)  $\frac{7 \pm \sqrt{57}}{4}$  (c)  $\frac{-7 \pm \sqrt{57}}{4}$  (d)  $\frac{7 \pm \sqrt{41}}{4}$

40. Solve using the Quadratic Formula.

$5x^2 + 9x = -11$

(a)  $\frac{9 \pm \sqrt{139}i}{10}$  (b)  $\frac{-9 \pm \sqrt{301}i}{10}$  (c)  $\frac{9 \pm \sqrt{301}i}{10}$  (d)  $\frac{-9 \pm \sqrt{139}i}{10}$

**The Discriminant of a Quadratic Equation**41. Determine the nature of the roots:  $2x^2 + 4x - 5 = 0$ .


- (a) Two equal real roots (b) No real roots (c) Two distinct real roots (d) Cannot be determined

42. Determine the number of real solutions of the equation.  $3x^2 + 6x + 3 = 0$
- (a) 3 (b) 0 (c) 1 (d) 2

43. Which of the following is the discriminant and the nature of the roots for  $2x^2 + 4x + 2 = 0$ ?
- (a) The discriminant is 0 and there are no real roots.      (b) The discriminant is 32 and there are two real roots.  
 (c) The discriminant is 0 and there is one real root.      (d) The discriminant is 32 and there is one real root.
44. Determine the number and nature of the roots of the equation.  $4x^2 - 3x + 1 = 0$
- (a) one real root and one imaginary root      (b) one real root      (c) two imaginary roots      (d) two real roots

### Applications of Quadratic Equations

45. The width of a rectangular carpet is 3 feet less than its length. If the area of the carpet is 70 square feet, find its width.
- (a) 7 ft      (b) 18 ft      (c) 10 ft      (d) 6 ft
46. One integer is eight more than three times another. If the product of the integers is 35, find the two integers.
- (a)  $-5$  and  $-7$       (b) 5 and  $-7$       (c)  $-5$  and 7      (d) 5 and 7
47. A sidewalk was built around a rectangular garden. If the area of the sidewalk is  $224 \text{ m}^2$ , find  $x$ . (Note: Figure not drawn to scale.)



- (a) 7 m      (b) 8 m      (c) 4 m      (d) 6 m
48. The length of a rectangle is 7 cm less than three times its width. If the area is  $40 \text{ cm}^2$ , find the dimensions of the rectangle.
- (a)  $8 \text{ cm} \times 5 \text{ cm}$       (b)  $8 \text{ cm} \times 6 \text{ cm}$       (c)  $47 \text{ cm} \times 33 \text{ cm}$       (d)  $280 \text{ cm} \times 6 \text{ cm}$

### Section 1.4 Other Types of Equations

#### Polynomial Equations

49. Solve.      (a)  $\pm 4, 1$       (b)  $-1, 16$       (c)  $\pm 4, -1$       (d) 1, 4  
 $x^3 + x^2 - 16x - 16 = 0$
50. Solve by factoring.  $4x(3x^2 + 11x + 8) = 0$       (a)  $-\frac{8}{3}$       (b)  $-1, -\frac{8}{3}$       (c) 0      (d)  $0, -1, -\frac{8}{3}$
51. Solve the polynomial equation by factoring.  
 $5x^3 - 10x^2 = 75x$
- (a)  $-4, 0, 2$       (b)  $-5, 0, 3$       (c)  $-3, 0, 5$       (d)  $-2, 0, 4$
52. Solve:      (a)  $0, \pm 36$       (b)  $\pm \sqrt{1296}$       (c)  $0, \pm 6$       (d)  $\pm \sqrt{210}$   
 $6x^4 - 216x^2 = 0$

**Rational Equations**

Solve the equation.

53.  $-\frac{1}{k} - 3 = -\frac{2}{3}$  (a)  $\frac{2}{5}$  (b)  $-\frac{3}{7}$  (c)  $2\frac{1}{2}$  (d)  $-2\frac{1}{3}$

54.  $\frac{x-5}{x+3} = \frac{x-4}{x-8}$  (a)  $-\frac{7}{3}$  (b) 7 (c)  $-\frac{3}{10}$  (d)  $\frac{13}{3}$

55.  $\frac{-3x}{x-4} + 10 = \frac{5x}{x-4}$  (a)  $\frac{1}{20}$  (b) 20 (c) -20 (d)  $-\frac{1}{20}$

56.  $\frac{x}{x^2-81} + \frac{9}{x-9} = \frac{1}{x+9}$  (a) 10 (b) -10 (c) -8 (d) 8

**Radical Equations**

Solve.

57.  $\sqrt{x+9} - 3 = x$  (a) -5 (b) 7 (c) -5, 7 (d) 0

58.  $\sqrt{6x-3} = \sqrt{5x+4}$  (a) 7 (b) 6 (c) 3 (d) No solution

59. Solve.  $\sqrt{x^2+8x+21} = 3$  (a) 2, -6 (b) 2, 6 (c) -2, -6 (d) -2

60. Solve.  $\sqrt{x-2} = x-4$  (a) 3 (b) 6, 3 (c) 6 (d) no solution

**Equations that are Quadratic in Form**

Solve:

61.  $6x^{-2} + x^{-1} + 1 = 0$  (a)  $x = 3, x = -2$  (b)  $x = 3, x = \frac{1}{2}$  (c)  $x = -\frac{1}{3}, x = -\frac{1}{2}$  (d) no solution

62.  $5\sqrt{x} - 2x - 2 = 0$  (a)  $\frac{1}{4}, 5$  (b)  $\frac{1}{4}, 4$  (c)  $-\frac{1}{4}, -4$  (d)  $\frac{1}{6}, 4$

63.  $15x^{-2} - 2x^{-1} + 1 = 0$  (a)  $x = 3, x = -5$  (b)  $x = -\frac{1}{3}, x = -\frac{1}{5}$  (c)  $x = 3, x = \frac{1}{5}$  (d) no solution

64.  $x^{2/3} - 4x^{1/3} - 32 = 0$   
(a)  $x = 512$  or  $x = 16$  (b)  $x = 512$  or  $x = -64$  (c)  $x = 64$  or  $x = 64$  (d)  $x = 8$  or  $x = -4$

**Section 1.5 Inequalities****Properties of Inequalities**

Solve:

65.  $-19 + w < -2$  (a)  $\{w \mid w > -21\}$  (b)  $\{w \mid w < -21\}$  (c)  $\{w \mid w > 17\}$  (d)  $\{w \mid w < 17\}$

Solve:

66.  $20b - 4 \leq 21b + 6$       (a)  $\{b \mid b \leq 10\}$       (b)  $\{b \mid b \geq -10\}$       (c)  $\{b \mid b \geq 24\}$       (d)  $\left\{b \mid b = -\frac{10}{41}\right\}$

67.  $12x - 9x + 18 > 2x - (10 - 2x)$

(a)  $\{x \mid x > 28\}$       (b)  $\{x \mid x < 28\}$       (c)  $\{x \mid x < -4\}$       (d)  $\{x \mid x > -4\}$

68.  $\frac{9}{8} - \frac{1}{2}x + \frac{7}{8} \leq 9x - \frac{3}{2}$       (a)  $\left\{x \mid x \geq \frac{7}{19}\right\}$       (b)  $\left\{x \mid x \geq \frac{7}{17}\right\}$       (c)  $\left\{x \mid x \leq \frac{7}{19}\right\}$       (d) none of these

**Compound Inequalities**

69. Solve.  $x - 4 \leq 0$  or  $x > 7$       (a)  $x \leq 4$  or  $x > 7$       (b)  $4 \leq x < 7$       (c)  $4 < x \leq 7$       (d)  $x < 4$  or  $x \geq 7$

70. Solve.  $3x - 1 > -13$  or  $2x + 3 < 1$       (a)  $x > -4$       (b)  $x < -1$       (c) all real numbers      (d) no solution

71. Solve.  $x + 2 \geq -2$  and  $x < 7$       (a)  $x \leq -4$  or  $x > 7$       (b)  $x < -4$  or  $x \geq 7$       (c)  $-4 < x \leq 7$       (d)  $-4 \leq x < 7$

72. Solve the inequality and give the solution in interval notation.

$2x - 1 > -9$  or  $3x - 3 < -9$

(a)  $(-4, \infty)$       (b)  $(-\infty, -2)$       (c)  $(-\infty, \infty)$       (d)  $\emptyset$

**Absolute Value Inequalities**

73. Solve.  $|3x + 3| > 3$

(a)  $\{x \mid -2 < x < 0\}$       (b)  $\{x \mid x \leq -2$  or  $x \geq 0\}$       (c)  $\{x \mid x < -2$  or  $x > 0\}$       (d) none of these

74. Solve:

$|2x - 3| \geq 2$

(a)  $\left\{x \mid \frac{1}{2} < x < \frac{5}{2}\right\}$       (b)  $\left\{x \mid x < \frac{1}{2}$  or  $x > \frac{5}{2}\right\}$       (c)  $\left\{x \mid x \leq \frac{1}{2}$  or  $x \geq \frac{5}{2}\right\}$       (d)  $\left\{x \mid \frac{1}{2} \leq x \leq \frac{5}{2}\right\}$

75. Solve.  $|x - 2| \geq 2$

(a)  $\{x \mid x \leq 0$  or  $x \geq 4\}$       (b)  $\{x \mid 0 \leq x \leq 4\}$       (c)  $\{x \mid 0 < x < 4\}$       (d)  $\{x \mid x < 0$  or  $x > 4\}$

76. Solve the inequality.

$|x - 6| < 2$

(a)  $x \leq 4, x \geq 8$       (b)  $4 \leq x \leq 8$       (c)  $4 < x < 8$       (d)  $x < 4, x > 8$

**The Critical Value Method**

Solve.

77.  $(x-7)(7x+1) \geq 0$

(a)  $\left\{x \mid -\frac{1}{7} \leq x \leq 7\right\}$  (b)  $\left\{x \mid -7 \leq x \leq \frac{1}{7}\right\}$  (c)  $\left\{x \mid x \leq -\frac{1}{7} \text{ or } x \geq 7\right\}$  (d)  $\left\{x \mid x \leq -7 \text{ or } x \geq \frac{1}{7}\right\}$

78.  $x^2 + 5x \geq 14$

(a)  $\{x \mid x \leq -7 \text{ or } x \geq 2\}$  (b)  $\{x \mid -2 \leq x \leq 7\}$  (c)  $\{x \mid x \leq -2 \text{ or } x \geq 7\}$  (d)  $\{x \mid -7 \leq x \leq 2\}$

79.  $x^2 + 11x + 18 > 0$

(a)  $\{x \mid 2 < x < 9\}$  (b)  $\{x \mid x < -9 \text{ or } x > -2\}$  (c)  $\{x \mid x < 2 \text{ or } x > 9\}$  (d)  $\{x \mid -9 < x < -2\}$

80. Solve the inequality and give the solution in interval notation.

$x^2 - 12x - 5 > 0$

(a)  $(-\infty, 6 - \sqrt{41}) \cup [6 + \sqrt{41}, \infty)$  (b)  $(-\infty, 6 - \sqrt{41}) \cup (6 + \sqrt{41}, \infty)$

(c)  $(6 - \sqrt{41}, 6 + \sqrt{41})$  (d)  $[6 - \sqrt{41}, 6 + \sqrt{41}]$

**Rational Inequalities**81. Solve the inequality: (a)  $-3 < x \leq 2$  (b)  $x < -3, x \geq 2$  (c)  $x < -3, x \geq 23$  (d)  $-3 < x \leq 23$ 

$$\frac{x+23}{x+3} \geq 5$$

82. Solve:

$$\frac{(x-5)(x+3)}{x-3} \geq 0$$

(a)  $3 \leq x \leq -5$  (b)  $x \geq 5 \text{ or } -3 \leq x < 3$  (c)  $x \leq -3 \text{ or } 3 < x \leq 5$  (d)  $x \leq -3 \text{ or } x \geq -5$

83. Identify the solution set of the inequality:  $\frac{5x+1}{x-1} \geq 7$ 

(a)  $1 < x \leq 4$  (b)  $1 \leq x \leq 4$  (c)  $x \leq 4$  (d)  $x < 1 \text{ or } x \geq 4$

84. Solve the equation or inequality.

$$\frac{2x-3}{x^2-36} \leq \frac{1}{x+6}$$

(a)  $x < -6 \text{ or } x \geq 3$ ; undefined at  $x = 6$  (b)  $-3 \leq x < 6$ ;  $x < -6$   
(c)  $-6 < x \leq -3 \text{ or } x > 6$ ; undefined at  $x = -6$  (d)  $-6 < x \leq 3$

**Applications**

85. The daily cost of renting a car is \$30 plus \$0.20 per mile. Joan's budget allows her to spend a maximum of \$42 for a 1-day rental. How many miles may Joan drive the rental car in one day without exceeding her budget of \$42?

(a)  $\leq 60$  mi (b)  $\geq 60$  mi (c)  $\leq 12$  mi (d)  $\geq 12$  mi



# Chapter 1: Equations and Inequalities (Answer Key)

## Section 1.1 Linear and Absolute Value Equations

### Linear Equations

[1] (d) \_\_\_\_\_

[2] (c) \_\_\_\_\_

[3] (b) \_\_\_\_\_

[4] (c) \_\_\_\_\_

### Contradictions, Conditional Equations, and Identities

[5] (c) \_\_\_\_\_

[6] (a) \_\_\_\_\_

[7] (e) \_\_\_\_\_

[8] (b) \_\_\_\_\_

### Absolute Value Equations

[9] (b) \_\_\_\_\_

[10] (b) \_\_\_\_\_

[11] (a) \_\_\_\_\_

[12] (d) \_\_\_\_\_

### Applications

[13] (c) \_\_\_\_\_

[14] (a) \_\_\_\_\_

[15] (c) \_\_\_\_\_

[16] (c) \_\_\_\_\_

## Section 1.2 Formulas and Applications

### Formulas

[17] (b) \_\_\_\_\_

[18] (a) \_\_\_\_\_

[19] (b) \_\_\_\_\_

[20] (d) \_\_\_\_\_

### Applications

[21] (c) \_\_\_\_\_

[22] (a) \_\_\_\_\_

[23] (d) \_\_\_\_\_

[24] (c) \_\_\_\_\_

## Section 1.3 Quadratic Equations

### Solve Quadratic Equations by Factoring

[25] (d) \_\_\_\_\_

[26] (d) \_\_\_\_\_

[27] (b) \_\_\_\_\_

[28] (d) \_\_\_\_\_

### Solve Quadratic Equations by Taking Square Roots

[29] (b) \_\_\_\_\_

[30] (c) \_\_\_\_\_

[31] (c) \_\_\_\_\_

[32] (a) \_\_\_\_\_

### Solve Quadratic Equations by Completing the Square

[33] (a) \_\_\_\_\_

[34] (b) \_\_\_\_\_

[35] (d) \_\_\_\_\_

[36] (d) \_\_\_\_\_

### Solve Quadratic Equations by Using the Quadratic Formula

[37] (b) \_\_\_\_\_

[38] (c) \_\_\_\_\_

[39] (b) \_\_\_\_\_

[40] (d) \_\_\_\_\_

## The Discriminant of a Quadratic Equation

[41] (c) \_\_\_\_\_

[42] (c) \_\_\_\_\_

[43] (c) \_\_\_\_\_

[44] (c) \_\_\_\_\_

## Applications of Quadratic Equations

[45] (a) \_\_\_\_\_

[46] (a) \_\_\_\_\_

[47] (b) \_\_\_\_\_

[48] (a) \_\_\_\_\_

## Section 1.4 Other Types of Equations

### Polynomial Equations

[49] (c) \_\_\_\_\_

[50] (d) \_\_\_\_\_

[51] (c) \_\_\_\_\_

[52] (c) \_\_\_\_\_

### Rational Equations

[53] (b) \_\_\_\_\_

[54] (d) \_\_\_\_\_

[55] (b) \_\_\_\_\_

[56] (b) \_\_\_\_\_

### Radical Equations

[57] (d) \_\_\_\_\_

[58] (a) \_\_\_\_\_

[59] (c) \_\_\_\_\_

[60] (c) \_\_\_\_\_

## Equations that are Quadratic in Form

[61] (d) \_\_\_\_\_

[62] (b) \_\_\_\_\_

[63] (d) \_\_\_\_\_

[64] (b) \_\_\_\_\_

**Section 1.5 Inequalities**

**Properties of Inequalities**

[65] (d) \_\_\_\_\_

[66] (b) \_\_\_\_\_

[67] (b) \_\_\_\_\_

[68] (a) \_\_\_\_\_

**Compound Inequalities**

[69] (a) \_\_\_\_\_

[70] (c) \_\_\_\_\_

[71] (d) \_\_\_\_\_

[72] (c) \_\_\_\_\_

**Absolute Value Inequalities**

[73] (c) \_\_\_\_\_

[74] (c) \_\_\_\_\_

[75] (a) \_\_\_\_\_

[76] (c) \_\_\_\_\_

**The Critical Value Method**

[77] (c) \_\_\_\_\_

[78] (a) \_\_\_\_\_

[79] (b) \_\_\_\_\_

[80] (b) \_\_\_\_\_

**Rational Inequalities**

[81] (a) \_\_\_\_\_

[82] (b) \_\_\_\_\_

[83] (a) \_\_\_\_\_

[84] (b) \_\_\_\_\_

**Applications**

[85] (a) \_\_\_\_\_

[86] (b) \_\_\_\_\_

[87] (c) \_\_\_\_\_

[88] (a) \_\_\_\_\_

**Section 1.6 Variation and Applications**

**Direct Variation**

[89] (d) \_\_\_\_\_

[90] (b) \_\_\_\_\_

[91] (c) \_\_\_\_\_

[92] (a) \_\_\_\_\_

**Inverse Variation**

[93] (b) \_\_\_\_\_

[94] (a) \_\_\_\_\_

[95] (b) \_\_\_\_\_

[96] (d) \_\_\_\_\_

**Joint Variation and Combined Variation**

[97] (b) \_\_\_\_\_

[98] (b) \_\_\_\_\_

[99] (a) \_\_\_\_\_

[100] (b) \_\_\_\_\_