





Chapter 2: Functions and Graphs

Section 2.1 A Two-Dimensional Coordinate System and Graphs

Cartesian Coordinate Systems

1. Plot the point $C(-3, 5)$ on a rectangular coordinate system.




2. Identify the point at $(-5, -2)$.

(a) M

(b) P

(c) O

(d) N



3. Determine the quadrant in which (x, y) is located so that the conditions are satisfied.

$$x > 0 \text{ and } y = 5$$

(a) Quadrant I

(b) Quadrant II

(c) Quadrant III

(d) Quadrant IV

4. Determine the quadrant in which the point $(4, 1)$ is located, or the axis on which it is located.

(a) Quadrant I


(b) Quadrant II

(c) x -axis

(d) y -axis

The Distance and Midpoint Formulas


5. Find the distance between the two plotted points. (a) 9.1 (b) 17.2 (c) 16.2 (d) 8.1




6. Find the possible values of y if the points $(8, 11)$ and $(-7, y)$ are a distance $d = \sqrt{229}$ apart.

- (a) 10, -15 (b) 10, 15 (c) 9, -13 (d) 9, 13

7. Find the midpoint of \overline{QR} . (a) $(1, 2)$ (b) $(2, 1)$ (c) $(-2, -1)$ (d) $(-1, -2)$



8. The midsegment of a trapezoid connects the midpoints of the two non-parallel sides. Find the midpoint of the midsegment of the trapezoid if $A = (4e, 4f)$, $B = (4g, 4f)$, and $C = (4h, 0)$.




- (a) $(2e+2g, 2f+2f)$ (b) $(f+g+h, f)$ (c) $(e+g+h, 2f)$ (d) $(g+2h, 2f)$


Graph of an Equation

9. Which is the graph of $y = -7|x| - 4$?


(a)




(b)



(c)




(d)




Graph:

10. $y = \frac{9}{x}$


(a)




(b)



(c)







(d)





Graph:

11. $y = x^2 + 2$





12. Identify the graph of the quadratic function and list any x -intercepts.

$f(x) = x^2 - 4x + 4$



$(-1, 0), (2, 0)$

$(2, 0)$



$(2, 0)$


$(-1, 0), (3, 0)$

Intercepts

13. Identify the graph of the function and find the y-intercept.


$$f(x) = 0.7x + 5$$

(a)




y-intercept: $(0, -5)$

(b)




y-intercept: $(0, 5)$

(c)




y-intercept: $(0, -5)$

(d)



y-intercept: $(0, 5)$

14. Identify the x- and y-intercepts.



(a) x-intercept: $(0, 4)$

y-intercepts: $(-2, 0)$, $(3, 0)$, $(6, 0)$

(b) x-intercepts: $(-2, 0)$, $(3, 0)$, $(-6, 0)$

y-intercept: $(0, 4)$

(c) x-intercept: $(0, -4)$

y-intercepts: $(-2, 0)$, $(3, 0)$, $(6, 0)$

(d) x-intercepts: $(-2, 0)$, $(3, 0)$, $(6, 0)$

y-intercept: $(0, -4)$

15. Identify the x - and y -intercepts of the graph of the equation.

$$y = x^3 \sqrt{64 - x^2}$$

- (a) x -intercepts: $(0, 0)$, $(-64, 0)$, $(64, 0)$, $(8, 0)$, $(-8, 0)$
 y -intercept: $(0, 0)$
- (b) x -intercepts: $(0, 0)$, $(-64, 0)$, $(64, 0)$
 y -intercept: $(0, 0)$
- (c) x -intercepts: $(8, 0)$, $(-8, 0)$, $(-64, 0)$
 y -intercepts: none
- (d) x -intercepts: $(0, 0)$, $(8, 0)$, $(-8, 0)$
 y -intercept: $(0, 0)$

16. Find the x - and y -intercepts of the graph of the equation. $\frac{1}{2}x = 3$

- (a) 6, none (b) 0, 6 (c) none, 2 (d) 3, 0

Circles, Their Equations, and Their Graphs

17. Find the standard form of the equation of the specified circle.

Center: $(3, -5)$; Radius: $3\sqrt{3}$

- (a) $(x-3)^2 + (y+5)^2 = 27$ (b) $(x+3)^2 + (y-5)^2 = 9\sqrt{3}$
- (c) $(x-3)^2 + (y+5)^2 = 9\sqrt{3}$ (d) None of these

18. Find the center and radius of the circle with the given equation.

$$(x+4)^2 + (y+4)^2 = 9$$

- (a) $(-4, -4)$; 9 (b) $(4, 4)$; 3 (c) $(-4, -4)$; 3 (d) $(-4, 4)$; 3

19. Find the equation of a circle where $C(6, 5)$ and $D(-4, 3)$ are endpoints of a diameter.

- (a) $(x-1)^2 + (y-4)^2 = 28$ (b) $(x-1)^2 + (y-4)^2 = 26$
- (c) $(x-5)^2 + (y-1)^2 = 17$ (d) none of these

20. Find the standard form of the equation of the specified circle.


Center: $(-2, 3)$; Radius: 2

- (a) $(x-2)^2 + (y+3)^2 = 2$ (b) $(x+2)^2 + (y+3)^2 = 4$
- (c) $(x-2)^2 - (y+3)^2 = 2$ (d) $(x+2)^2 + (y-3)^2 = 4$

Section 2.2 Introduction to Functions

Relations

21. Choose a table that represents each relation shown.



(a)

x	y
-3	7
6	5
6	8
-8	-2
-4	7
-4	-9

(b)

x	y
-3	7
6	5
6	8
-8	-9
-4	7
-4	-2


(c)

x	y
-3	7
6	5
6	-2
-8	8
-4	7
-4	-9

(d)

x	y
-3	5
6	7
6	8
-8	-2
-4	7
-4	-9

22. Choose a table that represents each relation shown.



(a)

x	y
9	-8
-3	7
-3	-1
3	4
2	-8
2	-7

(b)

x	y
9	-8
-3	7
-3	-1
3	-7
2	-8
2	4

(c)


x	y
9	-8
-3	7
-3	4
3	-1
2	-8
2	-7

(d)

x	y
9	7
-3	-8
-3	-1
3	4
2	-8
2	-7


23. Construct a model of the relation $\{(-8, -6), (-7, -1), (-8, -3), (-2, 9)\}$. Determine whether the relation is a function.

(a)




Function

(b)




Function

(c)




Not a function

(d)



Not a function

24. Find the domain and range of the relation.



(a) $D = \{0, 7, -7, -5, 5\}$

$R = \{-4, 0, 3, -8\}$

(c) $D = \{-4, 0, 3, -8\}$

$R = \{0, 7, -7, -5, 5\}$

(b) $D = \{-4, 0, 3\}$

$R = \{0, 7, -5, 5\}$

(d) $D = \{0, 7, -5, 5\}$

$R = \{-4, 0, 3\}$

Functions

25. Which set of ordered pairs (x, y) represents y as a function of x ?

(a) $\{(-2, 2), (4, -4), (4, -2), (-4, 4)\}$

(b) $\{(-2, 2), (2, 4), (-2, -4)\}$

(c) $\{-2, 2, 4, -4\}$

(d) $\{(-2, 2), (2, -2), (-4, -4)\}$

26. Which of the following relations is *not* a function?

(a)

x	y
-6	8
-5	6
-4	4

(b) $\{(3, 4), (3, 8), (5, 7), (7, 6)\}$

(c)

x	y
-6	4
-5	1
-4	0

(d) $\{(3, 4), (6, 4), (5, 7), (7, 3)\}$

27. Which of the following relations is *not* a function?

(a)

x	y
-8	3
-7	1
-7	8

(b) $\{(7, 8), (1, 3), (9, 2), (2, 8)\}$

(c)

x	y
-7	1
-6	1
-5	13

(d)

x	y
-7	-1
-6	-4
-5	4

28. Which of the following relations is also a function?

(a)

x	y
-7	6
-7	4
-8	2

(b)

x	y
-9	6
-8	4
-8	2

(c)

x	y
-8	6
-7	4
-8	2

(d)

x	y
-8	4
-7	4
-6	7

Functional Notation

Evaluate the function at the specified value(s) of the independent variable and simplify.

29. $f(x) = |x| + 4$; $f(6)$ (a) 10 (b) -2 (c) 2 (d) -10

30. $f(x) = 9x^2 - \sqrt{-3x}$; $f(-2)$ (a) 39.464 (b) 33.551 (c) 40.243 (d) 30

31. $g(x) = \frac{x^2 - 3}{4x}$; $g(n-5)$ (a) $\frac{n^2 - 3}{4n} - 5$ (b) $\frac{n^2 - 10n + 22}{4n - 5}$ (c) $\frac{n^2 - 8}{4n - 20}$ (d) $\frac{n^2 - 10n + 22}{4n - 20}$

32. Evaluate the function and simplify the results.

$$f(x) = \begin{cases} -\frac{3}{4}x & \text{if } x < -1 \\ 7 - 8x & \text{if } x \geq -1 \end{cases}$$





- a. $f(-1)$ b. $f(0)$
 c. $f(5)$ d. $f(-0.5)$

- (a) a. 15 b. 0 (b) a. $\frac{3}{4}$ b. 0 (c) a. 15 b. 7 (d) a. $\frac{3}{4}$ b. 7
 c. $-3\frac{3}{4}$ d. 11 c. $-3\frac{3}{4}$ d. $\frac{1}{2}$ c. -33 d. 11 c. -33 d. $\frac{1}{2}$

Identifying Functions

33. Which equation defines y as a function of x ? (a) $x = -6y^2$ (b) $x = 3$ (c) $x = 10$ (d) $y = 3$

34. Identify the graph that does *not* represent a function.



35. Which of the following is *not* a polynomial function?

- (a) $f(x) = \frac{3}{5}x^4 + \frac{x^{-3}}{5}$ (b) $f(x) = \frac{x-5}{4} - \frac{x}{5}$ (c) $f(x) = 2x^4 - x^3 + 2$ (d) $f(x) = \frac{1}{3}x^5 - \frac{x}{3}$

36. Determine which relation is a function with the correct domain and range.

(a)

x	y
-7	2
-6	9
-5	7

domain: $\{-7, -6, -5\}$ range: $\{2, 9, 7\}$ (b) $\{(6, 7), (6, 2), (8, 1), (1, 9)\}$ domain: $\{6, 8, 1\}$ range: $\{7, 2, 1, 9\}$

(c)

x	y
-6	2
-6	9
-7	7

domain: $\{2, 9, 7\}$ range: $\{-6, -7\}$

(d)

x	y
-8	2
-7	9
-8	3

domain: $\{-8, -7\}$ range: $\{2, 9, 3\}$


Graphs of Functions

37. Determine the intervals over which the function is increasing, decreasing or constant. $f(x) = \frac{1}{x^2 + 4x}$


- (a) increasing on $(-2, \infty)$; decreasing on $(-\infty, -2)$
- (b) increasing on $(-2, 0)$ and $(0, \infty)$; decreasing on $(-\infty, -4)$ and $(-4, -2)$
- (c) increasing on $(-\infty, -4)$ and $(-4, -2)$; decreasing on $(-2, 0)$ and $(0, \infty)$
- (d) increasing on $(-\infty, -2)$; decreasing on $(-2, \infty)$

38. Use the Vertical Line Test to determine which graph does *not* represent y as a function of x .


(a)




(b)



(c)




(d)




39. Graph: $f(x) = \begin{cases} \sqrt{x} + 1, & x \geq 0 \\ -4x - 3, & x < 0 \end{cases}$


(a)




(b)







(c)



(d)



40. Determine which of the following are one-to-one functions.







- (a) i and iv only (b) i, ii, and iv only (c) iv only (d) ii and iv only

The Greatest Integer Function

41. Identify the graph of the function. (The symbol $\left[\left[\right] \right]$ represents the greatest integer function.)

$$f(x) = \left[\left[\frac{1}{4}x \right] \right] - 3$$



42. A team-building incentive program for workers in an athletic shoe factory gives them a \$10,000 bonus to share for every 45,000 pairs of shoes completed. The basic production cost of a pair of high-quality shoes is \$45 before bonuses. Using $\lceil \cdot \rceil$ to denote the greatest integer function, find the equation that gives the cost, $C(x)$, for producing x high-quality shoes.

$$(a) C(x) = 10,000 \left\lceil \frac{x}{45,000} \right\rceil + 45x$$

$$(b) C(x) = 45,000 \left\lceil \frac{x}{10,000} \right\rceil + 45x$$

$$(c) C(x) = 10,000 \left\lceil \frac{x+1}{45,000} \right\rceil + 45x$$

$$(d) C(x) = 45x + 10,000 \left\lceil \frac{x-44,999}{45,000} \right\rceil$$

43. A taxi company charges \$5.45 for a ride of up to 3 miles, plus \$1.20 for each additional mile or fraction thereof. Using $\lceil \cdot \rceil$ to denote the greatest integer function, find the equation for the cost, $C(x)$, of a ride of x miles, given that $x \geq 3$ miles.

$$(a) C(x) = 5.45 + 1.20 \lceil x - 3 \rceil$$

$$(b) C(x) = 5.45 - 1.20 \lceil 3 - x \rceil$$

$$(c) C(x) = 1.20 - 5.45 \lceil 3 - x \rceil$$

$$(d) C(x) = 1.20 \lceil x + 3 \rceil + 5.45 \lceil x - 3 \rceil$$

44. A wireless telephone company that does not distinguish between local and long-distance calls charges \$26.95 per month and a per minute charge for use beyond 40 minutes. Fractions of a minute count as whole minutes. Using $\lceil \cdot \rceil$ to denote the greatest integer function, the cost, $C(x)$, for x minutes of use can be calculated using the following:

$$C(x) = \begin{cases} 26.95, & x \leq 40 \\ 26.95 - 0.28 \lceil 40 - x \rceil, & x > 40 \end{cases}$$

Find the cost for a person who uses 72.7 minutes in one month.

$$(a) \$36.19$$

$$(b) \$47.11$$

$$(c) \$36.09$$

$$(d) \$37.19$$

Applications

45. A music club membership costs \$21.00 to join and \$6 per CD. Find an equation that shows y , the cost for joining and buying x CDs.

$$(a) y = 21 + 6(x + 1)$$

$$(b) y = 27 + 6(x + 1)$$

$$(c) y = 27 + 6(x - 1)$$

$$(d) y = 21 + 6(x - 1)$$

46. Debbie is 36 years old, which is 12 less than 4 times Minnie's age. Which equation could be used to solve for Minnie's age? How old is Minnie?

$$(a) 4x + 12 = 36; 6 \text{ yr}$$

$$(b) 4x - 12 = 36; 12 \text{ yr}$$

$$(c) 12x - 4 = 36; 3.3 \text{ yr}$$

$$(d) 12x - 4 = 36; 2.7 \text{ yr}$$

47. Ms. Espinoza put away \$269 for a party fund. Each time her class did well on a big project, \$5 was deducted from the fund so that she could buy supplies for a class party. Choose the equation that represents the value remaining in her account after x withdrawals from the fund. Find the value remaining in the fund after 9 parties.

$$(a) V = 269 - 5x; \$224$$

$$(b) V = 5 - 269x; \$224$$

$$(c) V = 269 - 5x; \$1354$$





$$(d) V = 269 - 5x; \$1340$$

48. A sweatshirt printing business spends \$450 on equipment and supplies for each new batch of sweatshirts. In addition to these one time charges, the cost of printing each sweatshirt is \$4.25. Let x be the number of sweatshirts printed.
- Write the total cost C as a function of the total number of sweatshirts printed.
 - Write the average cost per unit, $\bar{C} = \frac{C}{x}$, as a function of x .
 - Find the average cost of printing 9000 sweatshirts.
 - $C = 4.25x - 450$
 - $C = 4.25x + 450$
 - $C = 4.25x + 450$
 - $C = 450x + 4.25$
- $\bar{C} = \frac{4.25x - 450}{x}$
 - $\bar{C} = \frac{4.25x + 450}{x}$
 - $\bar{C} = \frac{4.25x + 450}{x}$
 - $\bar{C} = \frac{450x + 4.25}{x}$
- \$3.50
 - \$4.30
 - \$5.30
 - \$4.50

Section 2.3 Linear Functions

Slopes of Lines

49. Find the slope of the line passing through the pair of points. (a) 0 (b) $\frac{5}{3}$ (c) $\frac{8}{5}$ (d) Undefined
 $(2, -8), (-8, -8)$
50. Find the slope of the line passing through the given points. (a) $\frac{18}{35}$ (b) $\frac{35}{18}$ (c) $\frac{20}{3}$ (d) $\frac{3}{20}$
 $\left(\frac{1}{2}, -3\right)$ and $\left(\frac{4}{5}, -1\right)$
51. Which is the graph of the line with a y -intercept of 4 and slope of 3?



52. Determine the value of r so the line that passes through each pair of points has the given slope.
- $(4, 2), (-2, r); m = -\frac{1}{3}$
- 12
 - 4
 - 20
 - $-6\frac{2}{3}$
 - None of these

Find the Equation of a Line

53. Which shows the equation of a line, in slope-intercept form, that passes through the point $(4, -5)$ with slope 4?


- (a) $y = 4x - 21$ (b) $y = 4x - 2$ (c) $y = -4x - 2$ (d) $y = -4x - 21$

54. Determine the equation of the line that has the given slope and contains the given point.

$$m = -\frac{1}{3}, (5, -3)$$

- (a) $3x + y = 14$ (b) $x + 3y = 14$ (c) $3x + y = -4$ (d) $x + 3y = -4$

55. Determine the slope-intercept form of the equation of the line.



- (a) $y = \frac{1}{2}x - 2$ (b) $y = 2x - 2$ (c) $y = -\frac{1}{2}x - 2$ (d) $y = -2x - 2$

56. Given $f(x) = 6x^2 - 5$, find the value of x which gives $f(x) = 49$.

- (a) $-3, 3$ (b) $-8.49, 3$ (c) $-18, 18$ (d) $4.5, -2.5$

Applications


57. A sloth travels at a rate of 0.4 meter per minute. Let t be the number of minutes spent traveling.

- Find the distance d traveled by the sloth as a function of t .
- Find the distance traveled by the sloth in 2 minutes.
- Construct a graph illustrating the distance traveled by the sloth versus time.

(a) a. $d(t) = t + 0.4$

b. 0.5 m


c.



(b) a. $d(t) = 0.4t + 1$

b. 0.9 m


c.



(c) a. $d(t) = 0.4t$

b. 0.8 m


c.



(d) a. $d(t) = 0.4d$

b. 0.6 m

c.



58. Casey McDonald plans to decorate vases to sell at a crafts fair. The decorations for all of the vases cost \$47.50 and the vases cost \$8.50 each.

- Write a function expressing the cost, $C(x)$ of the project in terms of the number of vases decorated, x .
- Determine the cost of decorating 60 vases.
- How many vases can be decorated with a budget of \$642.50?

(a) a. $C(x) = 8.50x + 47.50$

b. \$557.50

c. 70

(b) a. $C(x) = 47.50x + 8.50$

b. \$2858.50

c. 72

(c) a. $C(x) = 8.50x$

b. \$510.00

c. 71

(d) a. $C(x) = 8.50 + 47.50$

b. \$56.00

c. 73

59. A monthly phone bill, b , in dollars, consists of a \$22 service fee plus \$0.12 per minute, m , of long distance calls. Find an equation that relates the total cost to the number of minutes used. Then plot the total costs for using 8, 28, 45, and 58 minutes.

(a) $b = 0.12 + 22m$

(b) $b = 22 + 0.12m$

(c) $22 = b + 0.12m$

(d) $0.12 = 22 + bm$

60. In 1992 the average price of a house in a small town was \$97,000. By 1998, the average price of a home was \$109,000. Find a linear model for the price of such a house, P , if $t = 0$ represents the year 1992.
- (a) $P = 2000t + 97,000$ (b) $P = 109,000 - 2000t$ (c) $P = 109,000 - 12,000t$
 (d) $P = 12,000t + 97,000$ (e) None of these

Parallel and Perpendicular Lines

61. Determine whether the graphs of the equations are parallel, perpendicular, or neither.
 $-2x + 6y = -3$
 $12x + 4y = 0$
- (a) Parallel (b) Perpendicular (c) Neither
62. Which best describes the relationship between the lines with equations $-4x + 7y = -5$ and $-8x + 14y = 0$?
- (a) Perpendicular (b) Parallel (c) Same line (d) Neither parallel nor perpendicular
63. A woman standing in the middle of a large field was spinning a rock attached at the end of a rope, counterclockwise above her head. When she let go of the rope, the rock traveled along a linear path perpendicular to the radius created by the woman's position and the point of the rock's release. The rock traveled directly over a wall in the field. If the field is represented on a coordinate axis system with the woman at the origin, the point of release at $(6, 3)$, and the wall represented by the line $y = x + 8$, at what point was the ball directly over the wall?
- (a) $\left(\frac{7}{3}, \frac{31}{3}\right)$ (b) $\left(5, \frac{25}{3}\right)$ (c) $\left(23, \frac{31}{3}\right)$ (d) $\left(\frac{7}{3}, -\frac{1}{2}\right)$
64. A boy is playing with a ball suspended from a string, spinning it counterclockwise about the origin. He accidentally lets go of the ball and it hits the trunk of a tree in the adjoining backyard, after travelling along a path perpendicular to its radius of oscillation and the point at which it was released. If the point at which the ball was released is $P(-2, 8)$, and the equation representing the tree is $y = x + 6$, what is the point of contact of the ball and the trunk of the tree?
- (a) $\left(\frac{10}{3}, -40\right)$ (b) $\left(\frac{10}{3}, \frac{28}{3}\right)$ (c) $\left(\frac{34}{3}, -12\right)$ (d) $\left(\frac{29}{2}, \frac{28}{3}\right)$

Section 2.4 Quadratic Functions

Vertex of a Parabola

65. Find the vertex of the graph of the function.
 $f(x) = -(x - 2)^2 - 2$
- (a) $(2, 2)$ (b) $(-2, 2)$ (c) $(2, -2)$ (d) $(-2, -2)$
66. Which is the standard form of the function $f(x) = x^2 + 6x + 15$?
- (a) $f(x) = (x + 3)^2 - 6$ (b) $f(x) = (x + 3)^2 + 6$ (c) $f(x) = (x + 6)^2 - 9$ (d) $f(x) = (x + 6)^2 + 9$

67. Write the equation of the quadratic function in standard form and find the vertex of the graph.

$$f(x) = 30x + 4 + 3x^2$$

(a) $f(x) = 3(x+5)^2 - 71$

Vertex: $(-5, -71)$

(b) $f(x) = 3(x-5)^2 - 71$

Vertex: $(5, -71)$

(c) $f(x) = 3(x-5)^2 - 71$


Vertex: $(71, -5)$

(d) $f(x) = 3(x+5)^2 - 71$


Vertex: $(-71, -5)$

68. Write the equation in standard form. Then graph the equation. $y = x^2 - 2x + 4$


(a) $y = (x+1)^2 + 5$




(b) $y = (x-1)^2 + 3$



(c) $y = (x+1)^2 + 3$



(d) $y = (x-1)^2 + 5$



Maximum and Minimum of a Quadratic Function

69. Find the domain and range of the function.

$$f(x) = 5x^2 + 3$$

(a) Domain: $(-\infty, \infty)$; Range: $[3, \infty)$

(b) Domain: $(-\infty, \infty)$; Range: $(-\infty, 3]$

(c) Domain: $[3, \infty)$; Range: $(-\infty, \infty)$

(d) Domain: $(-\infty, 3]$; Range: $(-\infty, \infty)$

State whether the function has a minimum or a maximum value and find the value.

70. $f(x) = -x^2 + 4x$ (a) minimum: 2 (b) maximum: 2 (c) maximum: 4 (d) minimum: 4

71. $f(x) = x^2 + 2x + 4$ (a) maximum: -1 (b) minimum: 3 (c) minimum: -1 (d) maximum: 3

72. Use a calculator to find the maximum or minimum value of the parabola.

$$y = -x^2 + 5$$

- (a) min at $y = 5$ (b) max at $y = -5$ (c) max at $y = 5$ (d) max at $y = 6$ (e) min at $y = -5$

Applications

73. Find the maximum revenue for the revenue function $R(x) = 450x - 0.05x^2$.

- (a) 5000 (b) 1,012,500 (c) 171,000 (d) 4500

74. Moe estimates that the cost, C , to his company (in thousands of dollars) of producing x boxes of his computer chips can be modeled by $C = 360,000 + 0.01x + 0.0064x^2$. How many boxes should Moe's company produce to pay the minimum cost per box?

- (a) 7500 (b) 75 (c) 230,400 (d) 5625 (e) None of these

75. The area of a rectangle with a perimeter of 64 units is given by

$$A = x(32 - x)$$

where x represents the width of the rectangle. A farmer has available 64 yards of fencing and wishes to enclose a rectangular area. Use a graphing utility to graph the equation for the area of the rectangle and find the value of x that gives the largest area.

- (a) 24 yd (b) 21 yd (c) 16 yd (d) 11 yd


76. Macro Manufacturing estimates that its weekly profit, P , in hundreds of dollars, can be approximated by the formula $P(x) = -5x^2 + 40x + 2$, where x is the number of units produced per week, in thousands. How many units should the company produce per week to earn the maximum profit? Find the maximum weekly profit.

- (a) 3200 units, \$8800 (b) 4000 units, \$8200 (c) 4 units, \$8300 (d) 32 units, \$7800

Section 2.5 Properties of Graphs

Symmetry

77. Use symmetry tests to verify any of the three symmetries (x -axis, y -axis, or origin) the graph suggests.




- (a) The graph is symmetrical with respect to the origin.
 (b) The graph is symmetrical with respect to the x -axis, the y -axis, and the origin.
 (c) The graph is symmetrical with respect to the y -axis.
 (d) The graph has none of the three symmetries.
78. Identify the equation whose graph is symmetric with respect to the the origin.
- (a) $x = |y - 4|$ (b) $y^3 = 9x^4 - 7$ (c) $y = x^3 - 4$ (d) $9x^2 + 16y^2 = 144$

79. Which symmetries does the graph of $-3x^3y^4 - 6xy^6 = -5$ possess?


- (a) x -axis, y -axis, and the origin (b) origin only (c) x -axis only (d) y -axis only

80. Find the equation whose graph is symmetric with respect to the the origin.


(a) $9x^2 + 4y^2 = 36$




(b) $y^3 = 9x^4 - 7$



(c) $x = |y - 5|$




(d) $y = x^3 - 5$







Even and Odd Functions

81. Use the graph to determine if the function is even, odd, or neither. (a) Even (b) Odd (c) Neither



82. Which is the graph of a function that is neither even nor odd?



83. One of the following functions is neither odd nor even. Which one?





- (a) $2x^2 - |x^2| - 3$ (b) $x^5 - 3x^3 + 2x$ (c) $x^4 - 3x + 2 + |x - 2|$ (d) $-11x^3 - 3x$

84. Which of the following functions is even?


- (a) $g(x) = 2x^7 + 5x^3$ (b) $F(x) = -2x^6 + 5x^2 - 5$ (c) $f(x) = |-2x + 2| - 2$ (d) $h(x) = \frac{-2x^3}{-2x^2 + 2}$

Translations of Graphs


85. Use the graph of $f(x) = |x|$ to identify the graph of $f(x) = |x - 2| - 3$.




86. Use transformations of the graph of $y = x^2$ to determine the graph of $y = (x+1)^2 - 9$.




(a)




(b)








(c)







(d)



87. Using the graph of $f(x) = x^2$ as a guide, graph the function $g(x) = (x-3)^2 + 5$.




88. Without using graphing technology, sketch the parent graph and translate it to obtain the graph of $y - 3 = |x + 3|$.




Reflections of Graphs


89. The graph of the function $f(x) = \frac{1}{x}$ is shown below. Find the equation and the graph of the function $g(x)$ which is the reflection of $f(x)$ in the y -axis.




(a) $g(x) = -x$




(b) $g(x) = -\left(\frac{1}{x}\right)$





(c) $g(x) = x$



(d) $g(x) = \frac{-1}{-x}$








92. The graph of the function $f(x) = \frac{3}{4}\sqrt{x}$ is shown on the left below. The graph on the right is the same graph reflected over the y -axis. Find the equation of the graph on the right.



- (a) $f(x) = \frac{3}{4}\sqrt{x}$ (b) $f(x) = -\frac{3}{4}\sqrt{x}$ (c) $f(x) = -\frac{3}{4}\sqrt{-x}$ (d) $f(x) = \frac{3}{4}\sqrt{-x}$

Compressing and Stretching Graphs


93. Given is the graph of a function $l(x)$. Identify the graph of $2l(x)$.




96. Find an equation and graph for the function that is described by the given characteristics.

The shape of $f(x) = x^3$, with a vertical shrink of $\frac{1}{5}$.


(a) $g(x) = \frac{1}{5}x^3$




(b) $g(x) = -5x^3$



(c) $g(x) = 5x^3$



(d) $g(x) = -\frac{1}{5}x^3$



Section 2.6 The Algebra of Functions

Operations on Functions

97. Given $f(x) = \frac{7}{x-5}$ and $g(x) = \frac{9}{x+7}$, find $(f+g)(x)$.

(a) $\frac{-16x-4}{(x-5)(x+7)}$, $x \neq 5, -7$ (b) $-\frac{2}{2x+2}$, $x \neq -1$ (c) $\frac{16x+4}{(x-5)(x+7)}$, $x \neq 5, -7$ (d) $\frac{16}{2x+2}$, $x \neq -1$

98. If $f(x) = 4 - x^2$ and $g(x) = 2 - x$, find $(f \cdot g)(x)$.

(a) $-x^2 - x + 6$ (b) $2 + x$ (c) $x^3 - 2x^2 - 4x + 8$ (d) $-x^2 + x + 2$

99. Find $(f+g)(x)$ for $f(x) = \frac{1}{x^3}$ and $g(x) = x$.

(a) x^4 (b) $\frac{1-x}{x^4}$ (c) $\frac{x^3}{1+x^4}$ (d) $\frac{1+x^4}{x^3}$

100. Given $f(x) = 81 - x^2$, and $g(x) = 9 - x$, find $\frac{f}{g}(x)$.

(a) $x - 9$ (b) $x^2 - 81$ (c) $9 + x$ (d) $-x^2 + x + 72$

116. Do the results of the linear regression shown below seem to indicate a linear correlation between the variables? If they do, is it strong or moderate, positive or negative? If graphed, would the points tend to rise or fall as the value of the x variable increases?

<p>LinReg</p> $y = ax + b$ $a = 0.5369047619$ $b = -1.1047619048$ $r^2 = 0.3563168857$ $r = 0.5969228474$

- (a) There seems to be a moderately strong positive linear relationship between the variables.
 (b) There is a strong positive linear relationship between the variables.
 (c) There is a strong negative linear relationship between the variables.
 (d) A linear model may not be appropriate for the data.

Quadratic Regression Models

Use a graphing utility with a least squares regression program to find the least squares regression quadratic for the given points.

117. $(-3, -4)$, $(-2, 2)$, $(-1, 3)$

(a) $f(x) = -\frac{5}{2}x^2 + x + 1$

(b) $f(x) = -\frac{5}{2}x^2 - \frac{13}{2}x - 1$

(c) $f(x) = -\frac{5}{2}x^2 + x + \frac{1}{10}$

(d) $f(x) = -\frac{5}{2}x^2 - \frac{13}{2}x + \frac{1}{10}$

118. $(-1, 0)$, $(0, -4)$, $(1, -9)$, $(2, -8)$, $(3, -9)$

(a) $f(x) = \frac{6}{7}x^2 - \frac{137}{35}x + \frac{164}{175}$

(b) $f(x) = \frac{6}{7}x^2 + \frac{41}{9}x + \frac{163}{35}$

(c) $f(x) = \frac{6}{7}x^2 - \frac{137}{35}x - \frac{163}{35}$

(d) $f(x) = \frac{6}{7}x^2 + \frac{41}{9}x + \frac{164}{175}$

Chapter 2: Functions and Graphs (Answer Key)

Section 2.1 A Two-Dimensional Coordinate System and Graphs

Cartesian Coordinate Systems

[1] (d) _____

[2] (b) _____

[3] (a) _____

[4] (a) _____

The Distance and Midpoint Formulas

[5] (c) _____

[6] (d) _____

[7] (d) _____

[8] (c) _____

Graph of an Equation

[9] (a) _____

[10] (c) _____

[11] (a) _____

[12] (c) _____

Intercepts

[13] (d) _____

[14] (d) _____

[15] (d) _____

[16] (a) _____

Circles, Their Equations, and Their Graphs

[17] (a) _____

[18] (c) _____

[19] (b) _____

[20] (d) _____

Section 2.2 Introduction to Functions

Relations

[21] (a) _____

[22] (a) _____

[23] (c) _____

[24] (c) _____

Functions

[25] (d) _____

[26] (b) _____

[27] (a) _____

[28] (d) _____

Functional Notation

[29] (a) _____

[30] (b) _____

[31] (d) _____

[32] (c) _____

Identifying Functions

[33] (d) _____

[34] (c) _____

[35] (a) _____

[36] (a) _____

Graphs of Functions

[37] (c) _____

[38] (c) _____

[39] (c) _____

[40] (d) _____

The Greatest Integer Function

[41] (d) _____

[42] (a) _____

[43] (b) _____

[44] (a) _____

Applications

[45] (c) _____

[46] (b) _____

[47] (a) _____

[48] (b) _____

Section 2.3 Linear Functions

Slopes of Lines

[49] (a) _____

[50] (c) _____

[51] (d) _____

[52] (b) _____

Find the Equation of a Line

[53] (a) _____

[54] (d) _____

[55] (a) _____

[56] (a) _____

Applications

[57] (c) _____

[58] (a) _____

[59] (b) _____

[60] (a) _____

Parallel and Perpendicular Lines

[61] (b) _____

[62] (b) _____

[63] (a) _____

[64] (b) _____

