

Practice Test Math 002- Term 052

1) The domain of the function $f(x) = \ln\left(\frac{x-3}{x}\right)$, in interval notation, is equal to


- a) $(-\infty, 0) \cup (3, \infty)$
- b) $(3, \infty)$
- c) $(-\infty, 0)$
- d) $(0, 3)$
- e) non of the above

2) The range of the function $f(x) = e^{-|x|} - 1$, in interval notation, is equal to

- a) $(-1, 0]$
- b) $(-1, \infty]$
- c) $[0, \infty)$
- d) $(-\infty, 0]$
- e) $[1, 2]$

3) Which one of the following functions corresponds to the adjacent figure?

- a) $f(x) = \log(-x)$
- b) $f(x) = -\log|x|$
- c) $f(x) = -\log(-x)$
- d) $f(x) = \log(x-1)$
- e) $f(x) = \log(1-x)$



4) If $\log_x(\log_2 8) = 2$, the x is equal to

a) $\sqrt{3}$

b) 3

c) $-\sqrt{3}$

d) $\frac{3}{2}$

e) -3

5) The expression $\log_{x^2} b \times \log_{b^3} x$ is equal to

a) $\frac{1}{6}$

b) $\frac{1}{6} \log_x b$

c) $6 \log_x b$

d) 6

e) $\frac{1}{6} \log_b x$

6) If $f(x) = 2^{-x+1} - 1$, then $f^{-1}(x)$ is equal to

a) $1 - \log_2(x+1)$

b) $1 + \log_2(x+1)$

c) $2 - \log_2(2x+1)$

d) $2 + \log_2(2x+1)$

e) $\log_2 x + 2$

7) If $\log 2 = x$, then $\log_5 800$ is equal to

a) $\frac{3x+2}{1-x}$

b) $\frac{3x+2}{2}$

c) $3x$

d) $\frac{3x+2}{x+1}$

e) $\frac{3x}{5}$

8) The length of an arc that subtends a central angle of 120° in a circle of diameter 12 cm is equal to

a) 4π cm

b) $\frac{3\pi}{2}$ cm

c) 720 cm

d) 40 cm

e) $\frac{2\pi}{3}$

9) If a wheel with radius 10 centimeters is rotating at 100 revolutions per minute, then the linear speed of the wheel in centimeters per second is equal to

a) $\frac{100\pi}{3}$

b) 1000

c) $\frac{50\pi}{3}$

d) $\frac{20\pi}{3}$

e) $\frac{1000\pi}{3}$

10) Given the angles $\alpha = 51^\circ 49'$ and $\beta = 38^\circ 11'$. Which one of the following is true?

- a) α and β are complementary angles.
- b) $\alpha + \beta$ is an obtuse angle.
- c) $\alpha + \beta$ is an acute angle.
- d) α and β are supplementary angles.
- e) Non of the above.

11) The graph of $y = 2\sec\frac{1}{2}x$ and $y = 2\sin\frac{1}{2}x$, where $0 \leq x \leq 6\pi$, intersect at :

- a) no point
- b) two points
- c) one point
- d) four points
- e) three points

12) The x - intercept of the graph of the function $f(x) = 3 + 2\log_2(2x - 1)$ is:

- a) $(\frac{9}{16}, 0)$
- b) $(\frac{17}{14}, 0)$
- c) $(-\frac{21}{16}, 0)$
- d) $(\frac{11}{4}, 0)$
- e) $(\frac{23}{16}, 0)$

13) The sum of all solutions of $(\log x)^2 = \log x_{\frac{1}{10}}$ is equal to :

- a) $\frac{11}{10}$
- b) $\frac{101}{100}$
- c) 11
- d) 21
- e) 1

14) If W is the wrapping function with $W(t) = \left(\frac{3}{5}, -\frac{4}{5}\right)$, then the x -coordinate of $W\left(t - \frac{\pi}{2}\right)$ is equal to :

- a) $-\frac{4}{5}$
- b) $\frac{3}{5}$
- c) $\frac{4}{5}$
- d) $-\frac{3}{5}$
- e) $-\frac{3}{4}$

15) The range of the function $y = -\frac{5}{2} + \frac{3}{2}\sec(x + \pi)$ is equal to:

- a) $(-\infty, -4] \cup [-1, \infty)$
- b) $\left(-\infty, -\frac{3}{2}\right] \cup \left[\frac{3}{2}, \infty\right)$
- c) $\left[-\frac{3}{2}, \frac{3}{2}\right]$
- d) $[-4, -1]$
- e) $(-\infty, -1] \cup [1, \infty)$

16) The graph in the figure below is part of the graph of the equation:


a) $y = -3\sin\left(\frac{\pi}{4}x + \frac{\pi}{4}\right)$

b) $y = -3\cos\left(\frac{\pi}{4}x + \frac{\pi}{4}\right)$

c) $y = 3\sin(\pi x + \pi)$

d) $y = 3\sin\left(\frac{\pi}{2}x + \frac{\pi}{2}\right)$

e) $y = -3\cos(\pi x + \pi)$



17) The expression $\frac{\sin x + \csc x \cos^2 x + 1}{\sec x \csc x - \tan x} =$ is identical to

a) $\sec x + \tan x$

b) $\tan x - \csc x$

c) $\cos x + 1$

d) 0

e) $\sec x + \cot x$

18) $\frac{1 - \tan 21^\circ \cot 51^\circ}{\tan 21^\circ + \tan 39^\circ} =$

a) $\frac{\sqrt{3}}{3}$

b) $-\sqrt{3}$

c) $-\frac{\sqrt{5}}{2}$

d) $2\sqrt{2}$

e) $\frac{\sqrt{3}}{2}$

19) If $f(x) = a \cos bx$ with period = 8 and $f(4) = 3$, then $ab =$

a) $-\frac{3\pi}{4}$

b) $\frac{3\pi}{4}$

c) $-\frac{\pi}{4}$

d) $\frac{\pi}{4}$

e) $-\frac{\pi}{2}$

20) If $f(x) = -2 \tan \frac{2x}{3}$, then the number of vertical asymptotes in the interval $[-3\pi, 3\pi]$ is equal to

a) 4

b) 3

c) 2

d) 1

e) 5

21. If $f(x) = -2 \cos\left(\frac{\pi}{2}x + \pi\right) + 1$, then the **period + phase shift + amplitude** is equal to

a) 4

b) -4

c) 6

d) -6

e) 8

22) The expression $\sqrt{\frac{1 - \cos x}{1 + \cos x}}$, $0 < x < \frac{\pi}{2}$ is equal to

a) $\csc x - \cot x$

b) $\csc x + \cot x$

c) $\sec x + \tan x$

d) $\sec x - \tan x$

e) $\tan x + \cot x$

23) Given $\sin \alpha = \frac{4}{5}$, α in quadrant II, and $\sin(\frac{\pi}{2} - \beta) = -\frac{12}{13}$, β in quadrant IV, then

$\sec(\alpha + \beta)$ is equal to

a) $\frac{65}{63}$

b) $\frac{65}{33}$

c) $-\frac{65}{63}$

d) $-\frac{33}{63}$

e) *non of the above*

24) $\frac{\sin 15^\circ}{\sin 75^\circ} =$

a) $2 - \sqrt{3}$

b) $2 + \sqrt{3}$

c) $\frac{\sqrt{3}}{4}$

d) $\frac{\sqrt{3} + \sqrt{2}}{2}$

e) $\frac{\sqrt{3} - \sqrt{2}}{2}$

25) If $\frac{10^x - 10^{-x}}{10^x + 10^{-x}} = \frac{1}{2}$, then $x =$

a) $\frac{\log 3}{2}$

b) $\frac{\log 2}{3}$

c) $\log_2 3$

d) $\log_3 2$

d) $\log 6$