

# Unreliability of Co-Occurrence-Based Sediment Quality Guidelines for Contaminated Sediment Evaluations at Superfund/Hazardous Chemical Sites

Anne Jones-Lee

G. Fred Lee

---

---

---

*Many Superfund/hazardous chemical sites include waterbodies whose sediments contain hazardous chemicals. With the need to assess, rank, and remediate contaminated sediments at such sites, as well as in other waterways, regulators seek a simple, quantitative assessment approach that feeds easily into a decision-making scheme. Numeric, co-occurrence-based "sediment quality guidelines" have emerged with the appearance of administrative simplicity. However, the very foundation of the co-occurrence approach, based on the total concentrations of a chemical(s) in sediment, is technically invalid; its application relies on additional technically invalid presumptions. Use of technically invalid evaluation approaches renders any assessment of the significance of sediment contamination unreliable. This article reviews the technical roots and assumptions of the co-occurrence-based SOGs, the fundamental flaws in the rationale behind their development and application, and their misapplication for sediment quality evaluation. It also reviews concepts and approaches for the more reliable evaluation, ranking, and cleanup assessment of contaminated sediments at Superfund sites and elsewhere. © 2005 Wiley Periodicals, Inc.*

## INTRODUCTION

Many Superfund/hazardous chemical sites include waterbodies with sediments that have been contaminated by hazardous or otherwise deleterious chemicals associated with the site. The complexity of the remediation of these sites is compounded by the fact that the evaluation of the impacts of sediment-associated contaminants is itself complicated and not amenable to simple appraisal. With the need to assess, rank, and remediate contaminated sediments, an almost reflexive approach to the evaluation and management of contaminated sediment has been the quantification of the concentration of selected contaminants in the sediment. While chemical analysis is simple, renders numeric data for comparison, and lends itself to easy decision making, it has been long understood that there is no relationship between the total concentrations of chemical contaminants in sediments and toxicity, bioaccumulation, or other adverse impacts.

Nevertheless, in an attempt to make simple chemical analysis useful, some have developed and advanced the use of co-occurrence-based sediment quality guidelines (SOGs). These are being adopted by state and other regulatory agencies for any number of purposes, including sediment hot-spot identification and remediation, as well as Superfund site remediation. This is a source of mounting concern, however, because of

the inherent unreliability and technical invalidity of this approach. Use of technically invalid evaluation approaches renders any assessment of the significance of contamination invalid. It also renders invalid conclusions regarding the necessity for, specification of the type of, and assessment of the sufficiency of any proposed or undertaken remediation.

This article reviews the fundamental technical failings of the chemical concentration-based and co-occurrence-based sediment quality guideline approaches for assessing the potential impact of sediment-associated contaminants. It also reviews an approach that can be followed to make more technically valid, cost-effective assessments for identification, ranking, remediation, and management of sediment-associated contaminants. While the focus is largely on the misapplication of this approach for evaluating sediment-associated contaminants at federal and state Superfund sites, the information presented is equally applicable to the evaluation, regulation, and remediation of sediment-associated contaminants in any situation.

Use of technically invalid evaluation approaches renders any assessment of the significance of contamination invalid.

## WIDESPREAD PROBLEM

Leah Evison of the U.S. Environmental Protection Agency (US EPA) Office of Emergency and Remedial Response summarized the magnitude of the problem of contaminated sediments at Superfund sites in a presentation entitled "Contaminated Sediment at Superfund Sites: What We Know So Far," at a US EPA and U.S. Army Corps of Engineers (US ACE) national contaminated sediment workshop (Evison, 2003). Also at that workshop, the Sediments Team Leader at the US EPA Office of Emergency and Remedial Response presented a discussion entitled "Superfund Cleanup Issues at Contaminated Sediment Sites," highlighting the breadth of the problem (Ellis, 2003). In their recent summary of the scope of the contaminated sediment problem, Bridges et al. (2004) noted that in 1997, the US EPA concluded that there were 1.2 billion cubic yards of surface sediments in the United States that "pose potential risks." They also indicated that there are approximately 350 contaminated sediment sites within Superfund, about 30 of which are "megsites" (> \$50 M). The megasites include Hudson River, New York (\$460 M); New Bedford Harbor, Massachusetts (\$361 M); Bayou Bonfouca, Louisiana (\$90 M); Marathon Battery, New York (\$84 M); Triana/Tennessee River, Alabama (\$80 M); Fox River, Wisconsin (\$361 M); Silver Bow Creek, Montana (\$97 M); Commencement Bay, Washington (\$197 M); Bunker Hill (Coeur d'Alene Basin); and Housatonic, MA. Bridges et al. expect that other sites will be found as investigations continue.

The US EPA Superfund program has established a "Contaminated Sediments in Superfund" Web site (<http://www.epa.gov/superfund/resources/sediment/>) that contains additional information on the magnitude of the contaminated sediment issue at Superfund sites. The US EPA has also established a Superfund Sediment Resource Center (SSRC) where information is available on various technical aspects of managing contaminated sediments. Information on the SSRC and links to other Web site information sources (such as US EPA programs, the US ACE, the National Oceanic and Atmospheric Administration [NOAA], academia, and consultants) are available at [http://www.epa.gov/superfund/resources/sediment/ssrc\\_resources.htm](http://www.epa.gov/superfund/resources/sediment/ssrc_resources.htm).

Given the magnitude and extent of Superfund-associated and non-Superfund sediment contamination in the United States and the costs associated with their remediation, it is imperative that technically valid assessments be made in the screening of sediments for determining:

- the comparative urgency of attention,
- the environmental significance of the sediment-associated contaminants,
- the effectiveness of remediation alternatives, and
- the sufficiency of remediation accomplished at Superfund and other sites.

## “CO-OCCURRENCE”-BASED SQGS

As noted above, it has been understood for decades that there is no relationship between the total concentrations of chemical contaminants in sediments and toxicity, bioaccumulation, or other adverse impacts that may be caused by those contaminants. Such impacts are controlled by the aqueous environmental chemistry of chemicals in the aquatic sediments, which, in turn, is controlled by the physical, chemical, and biological character of the particular sediment/water environment. There is no simple protocol for making a reliable determination of the potential water/environmental quality significance of contaminants associated with sediment. As discussed in a subsequent section, protocols that have proven reliable require site-specific assessment of the release, behavior, and availability of the sediment-associated contaminants. However, legislative and regulatory imperatives for identifying and ranking contaminated sediment areas and ordering them for remedial attention, as well as for remediation of Superfund sites, beg for a simple, quantitative assessment approach that feeds easily into a decision-making scheme. An illusion of an easy comparative quantitation of sediment contamination has been offered by numeric, co-occurrence-based sediment quality guidelines. Because of its simplicity and ease of application, that approach has been embraced and advanced by some for evaluation and regulation of sediment-associated contaminants, without regard to its foundation and presumptions that render it technically invalid for the evaluation or regulation of contaminants in sediments.

Basically, the co-occurrence-based sediment quality guidelines were developed by examining a group of sediments for biological impact or “effect,” as measured any number of ways without consideration of what conditions or constituents were causing or influencing the effect. The concentrations of a few selected contaminants in those sediments were also measured. For each contaminant considered, the sediments were ordered by concentration, along with their associated assessments of their effects. In this ranked order, the lowest concentration of the contaminant in a sediment that had been identified as having an effect was noted; the concentration of that contaminant being considered became the basis for a “sediment quality guideline” and presumed to be a cause for concern in that, as well as any other, sediment. Fundamental but refutable presumptions made are:

- There is a causal relationship between the concentration of each constituent considered in a sediment and the water quality effect of that sediment.
- The effect reported for each sediment was caused independently by each of the measured chemical constituents in that sediment.
- No other condition(s) or chemical(s) influenced the effect.

Tables of such co-occurrences of chemicals at a given concentration and an effect developed by Long and Morgan (1990), Long et al. (1995, 1998), and MacDonald et al. (1992, 2002) have formed the foundation of what are termed “co-occurrence approaches” for sediment quality evaluation.

An illusion of an easy comparative quantitation of sediment contamination has been offered by numeric, co-occurrence-based sediment quality guidelines.

In regulatory applications, co-occurrence information has been used, albeit incorrectly, to establish various “effects threshold” values based on statistical manipulation. Examples of co-occurrence applications are the Apparent Effects Threshold (AET); the Effects Range Low (ER-L) and Effects Range Median (ER-M) values developed from Long and Morgan’s (1990) data presentation; the Probable Effects Level (PEL) values derived from MacDonald et al.’s (1992, 2002) co-occurrence compilations; and NOAA Screening Quick Reference Tables (SQiRT) values. If a sediment contains a chemical in concentrations above the AET, ER-M, PEL, or similar value, the sediment is considered by some regulators or in proposed regulations to be polluted, triggering special consideration such as remediation, alternative methods of dredged sediment disposal, or additional control of discharges to the waterbody of the chemical that can accumulate in sediments.

No consideration was given to many of the common chemicals and conditions that are, in fact, well understood to cause sediment toxicity.

## SUMMARY OF TECHNICAL DEFICIENCIES

### *Failure to Consider Cause and Effect*

As discussed by Lee and Jones-Lee (1993, 1996b, 2004), in the development of co-occurrence-based SQGs, no consideration has been given to the actual cause of the effect reported for any particular sediment. Independent of the cause of the biological response, the same degree of response was attributed to (co-occurred with) the concentration of each individual chemical constituent measured in the sediment. Further, no consideration was given to the fact that total concentration is not related to impact. No consideration was given to many of the common chemicals and conditions that are, in fact, well understood to cause sediment toxicity. No consideration was given to sediments that contain concentrations of the contaminant higher than the “effects”-associated concentration without exhibiting adverse impacts. The only basis for the so-called “guideline” concentration was the co-occurrence of that concentration of the contaminant in sediment and the exhibiting by that sediment of some biological impact.

While there may be statistical manipulations to create correlations between toxicity and an exceedance of a sediment quality guideline, such relationships are not demonstrations of cause and effect, but rather are coincidental and thus unreliable for assessing the cause of biological impacts. It is entirely expected that sediments in urban/industrial areas contain chemical constituents that cause biological effects, along with elevated concentrations of what, under some circumstances, could be toxic chemicals, which are in nontoxic, unavailable forms. The fact that a chemical constituent exceeds a particular sediment quality guideline does not mean that that constituent is in any way related to biological effects, such as toxicity, bioaccumulation, and/or changes in organism assemblages. The actual cause of the biological response seen could readily be another measured constituent, a constituent or constituents not measured or not considered in the scheme, or a combination of constituents that, while measured, do not, individually or summed, exceed the sediment quality guidelines. Thus, without a cause-and-effect relationship, there can be no expectation that funds spent to achieve SQG values will result in any improvement in sediment/water quality or that sediments targeted by the exceedance of guideline values are, in fact, of the greatest concern.

### ***Failure to Consider Full Range of Potential Pollutants***

The failure to find any exceedances of an ER-M, ER-L, PEL, or SQiRT value in sediment should never be assumed to be a reliable indication of a lack of impact of sediment-associated chemicals on aquatic life. Aside from the fundamental fatal defect in the basic premise of the approach, the co-occurrence-based sediment quality guidelines consider only a small number of the thousands of chemicals that can be present in sediments and can affect aquatic life. For example, Lee and Jones-Lee (2003a) discussed the fact that pyrethroid-based pesticides, one of the most common types of pesticides now used widely in agricultural and urban areas, tend to accumulate in sediments. Weston et al. (2004) reported finding aquatic sediments in the Central Valley of California with measurable toxicity and measurable concentrations of pyrethroid pesticides. The sediment quality guidelines do not include pyrethroid-based pesticides or other similar chemicals.

### ***Failure to Consider Ammonia, Hydrogen Sulfide or Low-Dissolved Oxygen***

As discussed by Lee and Jones-Lee (1996a), ammonia, hydrogen sulfide, and low-dissolved oxygen are, by far, the most common causes of sediment toxicity. However, none of these common causes were included in the co-occurrence-based sediment quality guidelines. In their extensive laboratory and field studies of the release of contaminants from sediment discussed subsequently, Jones and Lee (1988) found that the most common cause of sediment toxicity in their sediments was ammonia. Numerous other investigators have found this as well, making ammonia one of the most common causes of sediment toxicity. Co-occurrence-based sediment quality guidelines do not include consideration of ammonia as a potential cause of sediment toxicity, despite the fact that Long and Morgan (1990) made use of the authors' sediment database. While they used the heavy metal, organochlorine pesticide, and PCB data from the database, Long and Morgan ignored the ammonia and hydrogen sulfide data. Any sediment quality evaluation concerned with assessing toxicity to aquatic life that does not consider ammonia as a potential toxicant can be highly unreliable.

Any sediment quality evaluation concerned with assessing toxicity to aquatic life that does not consider ammonia as a potential toxicant can be highly unreliable.

### ***Failure to Consider Additive and Synergistic Toxicity***

Lee and Jones-Lee (2003a) have also discussed the fact that there is increasing evidence that the toxicities of pesticides and some other potentially toxic constituents, such as heavy metals, are additive and, in some cases, synergistic. This means that sediment that does not exceed any co-occurrence-based sediment quality guideline could be causing or contributing to sediment toxicity through additive or synergistic impacts with other chemicals for which there may or may not be an SQG. This type of chemical interaction intensifies the importance of assessing the potential for sediment impacts through toxicity testing.

### ***Failure to Consider Bioaccumulation as an Effect***

Bioaccumulation of chemicals to excessive levels in fish, causing them to be a threat to those who eat them, can occur at concentrations well below those that are toxic to benthic organisms. Lee and Jones-Lee (2002a) reviewed approaches for evaluating excessive concentra-

tions of organochlorine (OCl) pesticides in aquatic sediments. They also discussed the unreliability of co-occurrence-based approaches, such as Long and Morgan or MacDonald values, for predicting the water quality impact of the OCl's in sediments on the beneficial uses of waterbodies. Evaluation of sediments for these chemicals should be based on the toxicity of the sediment-associated chemicals to aquatic life and whether the sediment-associated OCl's contribute to excessive bioaccumulation in fish and other aquatic life tissue.

### *Use of Co-Occurrence for "Screening"*

As the need for sediment evaluation tools broadens and intensifies, SQGs have received a wider audience and been embraced by regulatory bodies looking for an easily implementable approach.

Some try to skirt the fundamental technical flaws in the co-occurrence-based SQG approach by limiting its use to "screening" sediments. However, using a patently unreliable screening tool can do nothing but provide patently unreliable results, which will serve to misdirect concern, responsibility, and funds for remediation, and leave real problem areas unaddressed. While sediments that exceed one or more sediment quality guidelines may, in fact, merit further investigation or remediation, the guideline values are meaningless for making that assessment. Under no circumstances should anyone assume that the exceedance of a guideline value implicates sediment as a potential, or actual, cause of problems, much less rank sediments for attention. Further, it cannot be assumed that not exceeding an SQG indicates that the sediment should be of diminished water quality concern. A reliable evaluation and regulatory program must be based on reliable assessments of the adverse effect and its cause.

### *Recent Reviews of the Unreliability Chemical Concentration-Based SQGs*

As the need for sediment evaluation tools broadens and intensifies, SQGs have received a wider audience and been embraced by regulatory bodies looking for an easily implementable approach. This has made the technical community more vocal on the unreliability of SQGs. Presented below is a summary of several presentations made at recent conferences that provide additional information on the applicability of SQGs.

*Society of Environmental Toxicology and Chemistry (SETAC) 2004 Conference.* Grapentine et al. (2004) reported on the results of a large study on the relationship between SQGs and sediment toxicity in selected nearshore sediment of the U.S.-Canadian Great Lakes. They found that exceedance of co-occurrence-based sediment quality guidelines was a poor predictor of sediment toxicity.

Bay et al. (2004) reported that exceedance of co-occurrence-based SQGs was a poor predictor of sediment toxicity in southern California coastal bays and nearshore marine sediments, even when applied to limited areas.

MacDonald et al. (2004) presented a paper comparing sediment toxicity with SQG values in a number of areas, including the Grand Calumet River in northern Indiana. That presentation could leave the mistaken impression that those SQG values had some validity for assessing the potential impact of sediment-associated contaminants in those sediments. Lee (2004b) discussed the unreliability of using MacDonald's co-occurrence-based SQGs to predict toxicity in Grand Calumet River sediments.

*Sediment Quality Assessment (SQA5) Conference.* In the fall of 2002, the International Aquatic Ecosystem Health and Management Society held a three-day conference in Chicago (Fifth International Conference on Sediment Quality Assessment [SQA5]) at which several presentations were made (including several invited keynote presentations)

on sediment quality evaluation. A number of the leading authorities on sediment quality evaluation, including DiToro, Chapman, and Burton, discussed the unreliability of co-occurrence-based approaches for evaluating sediment quality. In his presentation, DiToro (2002) characterized any so-called agreement between the results of co-occurrence-based values and observed sediment toxicity as a “coincidence,” and certainly not indicative of a cause-and-effect relationship. While Long, MacDonald, and others claim that their co-occurrence-based values have predictive capability based on a particular data set, a critical review of these data sets shows that they do not form a reliable basis for evaluating the ability of co-occurrence-based values to predict sediment toxicity, and certainly not anything that can be extrapolated to sediments outside the data set upon which they were based.

The focus of Chapman’s SQA5 discussion (Chapman, 2004) was the assessment of bioavailable forms of contaminants in sediments. He pointed out the unreliability of trying to assess bioavailability based on chemical measurements, noting specifically that co-occurrence-based sediment quality guidelines fail to reliably assess bioavailable forms of contaminants.

As discussed by O’Connor (1999a, 1999b; 2002), O’Connor and Paul (2000), O’Connor et al. (1998), Engler (personal communication, 2004), DiToro (2002), Chapman (2004), Burton (2002), Lee and Jones-Lee (1993; 1996a, 1996b, 1996c; 2004), the co-occurrence approaches are obviously technically invalid and unreliable for assessing cause and effect. From these assessments, it may be concluded that flipping a coin is a more reliable method of predicting sediment toxicity than exceedance of the guideline values.

Some, out of unawareness or obdurate advocacy, have claimed that co-occurrence-based SQGs are NOAA-generated or -supported values. Those values and approaches are not NOAA values. In fact, the chief scientist for the NOAA Status and Trends program (Thomas P. O’Connor) has repeatedly pointed out the unreliability of a Long and Morgan sediment quality guideline as an indicator of sediment toxicity. In his paper entitled “The Sediment Quality Guideline, ERL, Is Not a Chemical Concentration at the Threshold of Sediment Toxicity,” O’Connor (2004) has provided additional information on the unreliability of using co-occurrence-based approaches for assessing sediment toxicity. He stated in his abstract, “While it is being used as such, the sediment quality guideline ERL (effects range low) is not a threshold of any chemical concentration in sediment at which the probability of toxicity shows an abrupt increase. Similarly, while it has been done, there is no basis for assuming that multiple concentrations above an ERL increase the probability of toxicity.”

Long (2004) recently commented that it is possible to predict the broad scope of relationships between sediment chemical concentrations and toxicity. However, the error bars are very large, making such predictions worthless for site- or area-specific assessments. He concluded, “The presumption that you can predict benthic impacts with sediment chemistry data alone is very weak.”

Some, out of unawareness or obdurate advocacy, have claimed that co-occurrence-based SQGs are NOAA-generated or -supported values.

## EXAMPLES OF MISUSE OF CO-OCCURRENCE-BASED APPROACHES

Lee and Jones-Lee (2002b) summarized several examples of egregious misuse of co-occurrence-based SQGs in contaminated sediment remediation. The US EPA (2002) Region 9 is using co-occurrence based SQGs as cleanup objectives for organochlorine legacy pesticides in contaminated sediments in a Total Maximum Daily Load (TMDL)

program. The inappropriateness of this application is compounded by the fact that, as discussed above, the Long and Morgan and MacDonald “effects” did not include bioaccumulation as an effects measure, but rather focused on toxicity to aquatic life.

Even though there is substantial literature that shows that this approach is fundamentally unreliable, some state regulatory agencies, environmental groups, some parts of the US EPA, and others have been attempting to develop approaches that utilize the total concentration data that have been developed as part of contaminated sediment reconnaissance studies in regulating contaminated sediments.

SWRCB staff adopted an approach for the use of the available funds that failed to focus on gathering the information needed to develop SQOs.

### *State of California's Attempts to Develop Sediment Quality Objectives*

The California State Water Resources Control Board (SWRCB) is now trying to use a concentration-based approach for developing sediment quality objectives (SQOs), or standards, in its regulatory program. As discussed by Lee and Jones-Lee (2003b), the SWRCB is under a court decree to develop SQOs that can be used in a sediment quality regulatory program by August 2005. The goal is to identify, evaluate, and rank sediments in need of cleanup; National Permit Discharge Elimination System (NPDES) permits from public and private wastewater dischargers and stormwater management agencies could then be altered to reduce the concentrations of constituents that exceed the SQOs.

In the late 1980s, the California legislature adopted the Bay Protection and Toxic Cleanup Program (BPTCP), which required that the SWRCB develop SQOs that could be used to regulate polluted sediments. SWRCB staff adopted an approach for the use of the available funds that failed to focus on gathering the information needed to develop SQOs. This led to litigation by an environmental group. The litigation settlement led to the current effort by the SWRCB to try again to develop SQOs.

At the time this article was written, the renewed effort was in its eighteenth month. While the work plans for developing the SQOs have only recently been released, it appears that the SWRCB staff is making many of the same errors in attempting to develop SQOs as occurred in the BPTCP. Those who developed the work plans assert that the SQOs will be based on a weight-of-evidence approach involving developing numeric ranking of organism assemblage, sediment toxicity, and chemistry information. However, the approach for using the chemical concentration information is based on total concentration of chemicals relative to some co-occurrence-based sediment quality guideline. SWRCB staff states that some yet-to-be-developed total concentration-based value is to be used as a threshold to indicate when the regulatory agency needs to take action. As discussed herein, this approach is not technically valid; its inclusion invalidates the results of the weight-of-evidence assessment.

This approach ignores the substantial literature that exists on the lack of a relationship between the total concentration of a chemical in sediments and its water quality impacts. As discussed above, O'Connor (2004) of NOAA, Chapman (2004), DiToro (2002), and Lee and Jones-Lee (2003a, 2003b) have recently published information on the unreliability of total concentrations of a chemical as an indicator of sediment toxicity. The State Water Board staff and its selected Science Team (many members of which had been part of the failed BPTCP SQO development effort) have determined that the regional water boards' staffs who will be implementing the SQOs into regulatory programs do not have sufficient technical expertise to use aquatic chemistry, toxicology, and water quality information properly in evaluating sediment quality. They have deemed that, therefore, the

SQOs need to include “scaling factors” that would enable the regional board staff to use total concentrations of chemicals information without having to understand the unreliability of their use in evaluating the water quality/beneficial use impacts of the chemical. For the SWRCB staff to now propose total chemical concentration-based “scaling factors” for a chemical or chemicals in sediment, and presume that they are in some way related to sediment quality impacts as measured by aquatic life toxicity, reflects a lack of understanding of, or disregard for, the basic principles of aquatic chemistry/toxicology. As discussed by Lee and Jones-Lee (2003a), the appropriate use of chemical information must be based on toxic/available forms of chemicals and not on the total concentration of a chemical or group of chemicals. As summarized by Lee and Jones-Lee (2004), several internationally recognized authorities have warned against trying to use numeric scaling in weight-of-evidence evaluation of sediment quality.

In order for the SQOs to be scientifically defensible, they must not contain arbitrary components that can be easily shown to be technically invalid, such as the total concentration of a chemical when it is known that much, if not all, of the chemical is in a non-toxic form. It is important to consider the scrutiny that the SQOs will face in the regulatory process and the costs associated with their challenge. Regulatory boards and the courts are not likely to support the use of SQOs that can be shown to be based on invalid premises such as arbitrary scaling factors, inert, nontoxic chemical concentrations’ influencing the ranking of the pollution characteristics of the sediment, and so on. Dischargers or others who could become responsible for paying for sediment cleanup will rightfully challenge technically weak or invalid SQOs. While not included in the literature cited in support of the currently proposed SWRCB SQO work plans, there is a substantial literature that shows that the exceedance of a co-occurrence-based SQG is a technically invalid tool for assessing the impact of sediment-associated contaminants. This makes any SQO that uses an SQG as a trigger value unreliable for regulating or screening sediments. All of these issues were brought to the attention of the SWRCB staff and Board prior to their beginning to develop SQOs.

### ***Application of Co-Occurrence SQGs to NPL Superfund Sites***

Notwithstanding the extensive literature demonstrating the unreliability of the co-occurrence SQG approach for evaluating the potential impact of chemicals in sediments, the site investigators at two different National Priority List (NPL) Superfund sites at which the authors are US EPA Technical Assistance Grant (TAG) advisors to the public have proposed to use and/or have used chemical concentration-based SQGs as pass/fail values in an ecological risk assessment for the pollutants in the sites’ aquatic sediments. One of these locations is the University of California, Davis (UCD)/U.S. Department of Energy Laboratory for Energy-Related Health Resources (LEHR) national Superfund site. UCD is attempting to use Long et al. (1995) and MacDonald et al. (2000) co-occurrence-based guidelines as a technical basis for conducting an ecological risk assessment for contaminated sediments associated with that Superfund site. However, as discussed by Lee (2004a), that approach cannot be relied upon for characterization of LEHR site sediments with respect to their potential threat to aquatic life and, therefore, whether or not they should require remediation. A similar approach had been initiated at the Lava Cap Mine Superfund site near Nevada City, California; the US EPA contractor had proposed to use co-occurrence-based SQGs in an ecological risk assessment for the site. Following Lee’s

Dischargers or others who could become responsible for paying for sediment cleanup will rightfully challenge technically weak or invalid SQOs.

(2001) critique of the proposed approach, the US EPA changed the ecological risk assessment approach to focus on toxicity assessment rather than numeric SQGs.

The US EPA Superfund Program national headquarters' management has been concerned for a number of years about the potential for using co-occurrence-based sediment quality guidelines as a foundation for remediation decisions. The managers of US EPA Superfund have made it clear that that approach should not be followed, because of the unreliability of co-occurrence-based sediment quality guidelines. According to Ellis (2003), the US EPA Superfund Program supports the position that co-occurrence-based sediment quality guidelines are not appropriate for establishing the impacts of chemicals in sediments or to serve as the basis for sediment cleanup objectives.

The work done in the DMRP served as the basis for the US EPA and US ACE to develop criteria for open-water disposal of contaminated dredged sediments.

## BACKGROUND TO DEVELOPING SEDIMENT QUALITY EVALUATION

Numerous approaches have been proposed over the years for determining what constitutes an excessive level of a heavy metal, pesticide, or some other potential pollutant in aquatic sediments. One of the earliest efforts (and the most comprehensive) to develop and evaluate protocols for this purpose was undertaken by the US ACE, which has been mandated by Congress to maintain the navigation depth of U.S. waterways. With only unreliable bulk sediment guidelines for a few parameters available, an understanding that contaminant release and impact could not be determined based on sediment concentrations, and increasing awareness of the environmental impacts of chemical contaminants often associated with sediment, the US ACE recognized the need for developing an expedient, technically sound approach to assessing the potential impact of sediment-associated contaminants.

In the 1970s, Congress authorized and funded the U.S. Army Corps of Engineers Waterways Experiment Station in Vicksburg, Mississippi, to investigate the water quality impacts of dredged sediment disposal. This \$30 million, five-year Dredged Material Research Program (DMRP) focused on evaluating and developing management approaches for the water quality impacts of dredged sediment disposal. The author G. F. Lee served as an advisor to the US ACE to help develop that program, and directed about a million dollars' worth of the studies of the potential impacts of open-water disposal of contaminated dredged sediments.

In his DMRP studies, Lee and his graduate students evaluated the total concentrations of about 30 potential conventional pollutants, including heavy metals, organochlorine pesticides, PCBs, and nutrients in sediments from about 100 sites across the United States. They also examined the behavior of those chemicals in many of those sediments as they were suspended in the water column during the open-water disposal; disposal sites were extensively monitored for total and soluble contaminants and physical characteristics before, during, and following disposal operations. In addition, in conjunction with the field assessments, they worked with the US ACE to develop and evaluate laboratory protocols to assess the release and potential toxicity of sediment-associated contaminants disposed of in open water. This effort generated about 50,000 data points; the findings were published in two reports (Jones & Lee, 1978; Lee et al., 1978). A summary of those studies is published in the *Handbook of Dredging Engineering* (Lee & Jones-Lee, 2000).

The work done in the DMRP served as the basis for the US EPA and US ACE to develop criteria for open-water disposal of contaminated dredged sediments. The results clearly demonstrated that the total concentration of a particular chemical in sediment, such as a heavy metal or the sum of all heavy metals, bears no relationship to the

amount of that contaminant released to the water column during disposal or to the toxicity of that sediment to aquatic life. Many of the sediments collected from urban/industrial waterways caused toxicity to aquatic life in laboratory tests. This toxicity, however, was largely independent of the total concentrations of various conventional pollutants, such as heavy metals and pesticides, as a result of their sediment-binding characteristics. While the identification of the cause of laboratory toxicity was beyond the scope of the studies, in subsequent work, the authors (Jones & Lee, 1988) found a link between toxicity and ammonia released that was consistent with the known toxicity of ammonia. As noted above, ammonia was not even considered in the co-occurrence-based SQGs.

Thus, the results of the DMRP substantiated what was known at the outset of the studies—namely, that the total concentration of a constituent(s) in sediment is not a reliable indicator of the potential impacts of that sediment on the beneficial uses of the waters in which the sediment is located. The results also led the US ACE and the US EPA to develop, in the late 1970s, biological effects-based dredged sediment water quality evaluation procedures for both freshwater and marine dredged systems (US EPA/US ACE, 1991, 1998). Those protocols focused on aquatic life toxicity testing and assessing the potential for excessive bioaccumulation of hazardous chemicals from sediments. The dredging of contaminated Superfund and other sediments and their disposal requires attention to the same water quality/environmental quality and contaminant availability, transport, and impact issues that are considered in dredging and disposal of contaminated sediment associated with channel-depth maintenance.

## BPJ WEIGHT-OF-EVIDENCE APPROACH

Lee and Jones-Lee (2004) presented a paper at the SQA5 conference entitled “Appropriate Incorporation of Chemical Information in a Best Professional Judgment ‘Triad’ Weight of Evidence Evaluation of Sediment Quality.” In addition to updating the review of the unreliability of co-occurrence-based values for evaluating sediment quality, they gave particular attention to how technically valid sediment quality evaluations should be conducted. As they described, a best professional judgment (BPJ) Triad weight-of-evidence approach that integrates information on aquatic toxicity and excessive bioaccumulation, alteration in organism assemblages relative to habitat characteristics, and chemicals as a cause of toxicity/bioavailability should be used. Chemical information is incorporated into the approach through toxicity identification evaluations (TIEs) and sediment bioavailability testing to determine the toxicity/availability of the forms and mixtures of contaminants and characteristics in the sediment.

Regulatory agencies should adopt the technically sound approaches of the US EPA Superfund and US ACE . . .

## CONCLUSION

It will be important that other regulatory agencies not follow the California SWRCB and others in regulating sediment-associated chemicals based on SQGs and SQOs developed through, or influenced by, chemical concentration/co-occurrence approaches. Regulatory agencies should adopt the technically sound approaches of the US EPA Superfund and US ACE; they advocate making sediment quality evaluations through effects-based approaches using aquatic chemistry, aquatic life toxicity, and excessive bioaccumulation of hazardous chemicals in higher trophic-level organisms. Chemical concentration information should focus on the availability/toxicity of the chemical in a

TIE framework. Numeric scaling factors should not be used in sediment quality evaluation. Scaling schemes are arbitrary and can readily lead to a technically invalid assessment of sediment quality. The focus of the SQOs should be the use of effect-based assessment of aquatic toxicity to several types of sensitive aquatic life and whether there is excessive bioaccumulation of hazardous chemicals in edible aquatic life. Chemical studies should be used to identify the constituent(s) responsible for the toxicity through toxicity identification evaluation and the concentration of toxic/available constituents in the sediments. It should be combined with toxicity in forensic studies to identify the source of the toxic constituent(s). Where excessive bioaccumulation is found in edible fish and other aquatic life, the source of the bioaccumulatable chemicals should be determined and sediment uptake studies should be conducted to define the availability of the chemical in the sediments.

Adoption of this effects-based approach will lead to a technically valid regulatory approach for defining the sediments that are having an adverse impact on the beneficial uses of a waterbody.

## REFERENCES

- Bay, S., Vidal, D., Field, L., & Myre, P. (2004). Effect of spatial scale on relationships between sediment quality guidelines and toxicity. Presented at Fourth Society of Environmental Toxicology and Chemistry (SETAC) World Congress, Portland, OR.
- Bridges, T. S., Nadeau, S., & Ells, S. (2004). Addressing uncertainty and managing risk at contaminated sediment sites. Presentation at US ACE/US EPA/SMWG Joint Sediment Conference, St. Louis, MO. Retrieved December 29, 2004, from <http://el.erdc.usace.army.mil/training.cfm?Topic=Workshop&List=04oct-ccs>
- Burton, A. (2002). Presentation at Fifth International Conference on Sediment Quality Assessment. Sponsored by International Aquatic Ecosystem Health and Management Society, Chicago, IL.
- Chapman, P. M. (2004). Paracelsus' dictum for sediment quality (and other) assessments. *Aquatic Ecosystem Health & Management*, 7(3), 369–374.
- DiToro, D. (2002) Presentation at Fifth International Conference on Sediment Quality Assessment. Sponsored by International Aquatic Ecosystem Health and Management Society, Chicago, IL.
- Ellis, S. (2003). Superfund cleanup issues at contaminated sediment sites. Presented at US EPA and US Army Corps of Engineers National Workshop: Environmental Stability of Chemicals in Sediments, San Diego, CA. Retrieved December 29, 2004, from <http://www.sediments.org/sedstab/agenda.pdf>
- Evison, L. (2003). Contaminated sediment at Superfund sites: What we know so far. Presented at US EPA and US Army Corps of Engineers National Workshop: Environmental Stability of Chemicals in Sediments, San Diego, CA. Retrieved December 29, 2004, from <http://www.sediments.org/sedstab/agenda.pdf>
- Grapentine, L., Milani, D., & Reynoldson, T. (2004). Assessment of ecological effects of contaminated sediments in the Laurentian Great Lakes. Presented at Fourth Society of Environmental Toxicology and Chemistry (SETAC) World Congress, Portland, OR.
- Jones, R. A., & Lee, G. F. (1978). Evaluation of the elutriate test as a method of predicting contaminant release during open water disposal of dredged sediment and environmental impact of open water dredged material disposal, Vol. I: Discussion. Tech Report D-78-45. Vicksburg, MS: US ACE Waterway Experiment Station.

- Jones, R. A., & Lee, G. F. (1988). Toxicity of U.S. waterway sediments with particular reference to the New York Harbor Area. In J. J. Lichtenberg, F. A. Winter, C. I. Weber, & L. Fradkin (Eds.), *Chemical and biological characterization of sludges, sediments, dredge spoils and drilling muds* (pp. 403–417). ASTM STP 976. Philadelphia, PA: American Society for Testing Materials.
- Lee, G. F. (2001, August 22). Comments on final draft ecological risk assessment for the Lava Cap Mine Superfund site (undated). Report prepared by G. Fred Lee & Associates, El Macero, CA. Available upon request from [gfredlee@aol.com](mailto:gfredlee@aol.com).
- Lee, G. F. (2004a). Review of the UCD ecological risk assessment revised draft. Report submitted to Davis South Campus Superfund Oversight Committee, Davis, CA. Retrieved December 29, 2004, from <http://www.members.aol.com/dscsoc>
- Lee, G. F. (2004b). Comments on January 13, 2004, draft preliminary problem formulation technical memorandum for the West Branch of the Grand Calumet River, Lake County, Indiana. Prepared by Tetra Tech for the U.S. Fish and Wildlife Service. Comments submitted to the U.S. Fish and Wildlife Service on behalf of the Sanitary District of Hammond, Indiana, by G. Fred Lee & Associates, El Macero, CA. Retrieved December 29, 2004, from <http://www.gfredlee.com/Hammond-GCR-Comments.pdf>
- Lee, G. F., & Jones-Lee, A. (1993). Sediment quality criteria: Numeric chemical- vs. biological effects-based approaches. *Proceedings of the Water Environment Federation National Conference, Surface Water Quality & Ecology*, pp. 389–400.
- Lee, G. F., & Jones-Lee, A. (1996a). "Co-Occurrence" in sediment quality assessment. Report of G. Fred Lee & Associates, El Macero, CA. Retrieved December 29, 2004, from <http://www.members.aol.com/apple27298/COOCCUR2PAP.pdf>
- Lee, G. F., & Jones-Lee, A. (1996b). Can chemically-based sediment quality criteria be used as reliable screening tools for water quality impacts? *Society of Environmental Toxicology and Chemistry News*, 16(3), 14–15.
- Lee, G. F., & Jones-Lee, A. (1996c). Evaluation of the water quality significance of the chemical constituents in aquatic sediments: Coupling sediment quality evaluation results to significant water quality impacts. *Proceedings of the Water Environment Federation National Conference, Surface Water Quality and Ecology I & II*, 4, 317–328.
- Lee, G. F., & Jones-Lee, A. (2000). Water quality aspects of dredging and dredged sediment disposal. In J. B. Herbich (Ed.), *Handbook of dredging engineering* (2nd ed.; pp. 14-1–14-42). New York: McGraw-Hill. Retrieved December 29, 2004, from <http://www.gfredlee.com/dredging.html>
- Lee, G. F., & Jones-Lee, A. (2002a). Organochlorine pesticide, PCB and dioxin/furan excessive bioaccumulation management guidance. California Water Institute Report TP 02-06 to the California Water Resources Control Board/Central Valley Regional Water Quality Control Board, California State University Fresno, Fresno, CA. Retrieved December 29, 2004, from <http://www.gfredlee.com/OCITMDLRpt12-11-02.pdf>
- Lee, G. F., & Jones-Lee, A. (2002b). Unreliability of sediment co-occurrence-based approaches for evaluating aquatic sediment quality. Excerpted from California Water Institute Report TP 02-06 to the California Water Resources Control Board/Central Valley Regional Water Quality Control Board, California State University Fresno, Fresno, CA. Retrieved December 29, 2004, from <http://www.gfredlee.com/UnrelSedCooccur.pdf>
- Lee, G. F., & Jones-Lee, A. (2003a). Regulating water quality impacts of port and harbor stormwater runoff. Presented at the International Symposium on Prevention of Pollution from Ships, Shipyards, Drydocks,

- Ports, and Harbors, New Orleans, LA. Retrieved December 29, 2004, from <http://www.members.aol.com/duklee2307/PHStormwater-papfinal.pdf>
- Lee, G. F., & Jones-Lee, A. (2003b). Development of sediment quality objectives for California. Presented at the American Water Resources Association National Conference, San Diego, CA. Retrieved December 29, 2004, from <http://www.gfredlee.com/sediment-SQO-paperSD.pdf>
- Lee, G. F., & Jones-Lee, A. (2004). Appropriate incorporation of chemical information in a best professional judgment "triad" weight of evidence evaluation of sediment quality. Presented at the 2002 Fifth International Symposium on Sediment Quality Assessment (SQA5). Reprinted in *Aquatic Ecosystem Health & Management*, 7(3), 351–356. Retrieved December 29, 2004, from <http://www.gfredlee.com/BPJWOEpaper-pdf>
- Lee, G. F., Jones, R. A., Saleh, F. Y., Mariani, G. M., Homer, D. H., Butler, J. S., et al. (1978). Evaluation of the elutriate test as a method of predicting contaminant release during open water disposal of dredged sediment and environmental impact of open water dredged materials disposal, Vol. II: Data report. Technical Report D-78-45. Vicksburg, MS: US ACE Waterway Experiment Station.
- Long, E. (2004, October 27). Comments as reported in the minutes from the California Sediment Quality Objectives Advisory Committee Sixth Meeting. State of California Water Resources Control Board, Water Quality Program, Sacramento, CA.
- Long, E., & Morgan, L (1990). The potential for biological effects of sediment-sorbed contaminants tested in the National Status and Trends Program. NOAA Technical Memorandum NOS OMA 52, National Oceanographic and Atmospheric Administration, Seattle, WA.
- Long, E. R., Field, L. J., & MacDonald, D. D. (1998). Predicting toxicity in marine sediments with numerical sediment quality guidelines. *Environmental Toxicology and Chemistry*, 17(4), 714–727.
- Long, E. R., MacDonald, D. D., Smith, S. L., & Calder, F. D. (1995). Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environmental Management*, 19, 81–97.
- MacDonald, D. D., Brydges, K., & Haines, M. L. (1992). Development of an approach to the assessment of sediment quality in Florida coastal waters. Prepared for the Florida Coastal Management Program, Florida Department of Environmental Regulation, MacDonald Environmental Services, Ltd., Ladysmith, British Columbia.
- MacDonald, D. D., Ingersoll, C. G., & Berger, T. A. (2000). Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Archives of Environmental Contamination and Toxicology*, 39, 20–31.
- MacDonald, D., Ingersoll, C., Smorong, D., Sparks, D., Smith, J., Meyer, J., et al. (2004). A comparison between regional and national data sets for freshwater sediments. Presented at Fourth Society of Environmental Toxicology and Chemistry (SETAC) World Congress, Portland, OR.
- O'Connor, T. P. (1999a, January). Sediment quality guidelines do not guide. NOAA National Status and Trends Program, Learned Discourses: Timely Scientific Opinions. SETAC News.
- O'Connor, T. P. (1999b, May). Sediment quality guidelines reply-to-reply. NOAA National Status and Trends Program, Learned Discourses: Timely Scientific Opinions. SETAC News.
- O'Connor, T. P. (2002). Empirical and theoretical shortcomings of sediment-quality guidelines. In R. Whittemore et al. (Eds.), *Handbook on sediment quality* (pp. 317–325). Alexandria, VA: Water Environment Federation.

- O'Connor, T. P. (2004). The sediment quality guideline, ERL, is not a chemical concentration at the threshold of sediment toxicity. *Marine Pollution Bulletin*, 49(5/6), 383–385.
- O'Connor, T. P., Daskalakis, K. D., Hyland, J. L., Paul, J. F., & Summers, J. K. (1998). Comparisons of measured sediment toxicity with predictions based on chemical guidelines. *Environmental Toxicology and Chemistry*, 17, 468–471.
- O'Connor, T. P., & Paul, J. F. (2000). Misfit between sediment toxicity and chemistry. *Marine Pollution Bulletin*, 40(1), 59–64.
- United States Environmental Protection Agency (US EPA). (2002). Total maximum daily loads for toxic pollutants San Diego Creek and Newport Bay, California. U.S. Environmental Protection Agency, Region 9, San Francisco, CA.
- United States Environmental Protection Agency/United States Army Corps of Engineers (US EPA/US ACE). (1991). Evaluation of dredged material proposed for ocean disposal—Testing manual. EPA-503-8-91-001. Washington, DC: U.S. Environmental Protection Agency and U.S. Army Corps of Engineers.
- United States Environmental Protection Agency/United States Army Corps of Engineers (US EPA/US ACE). (1998). Evaluation of dredged material proposed for discharge in waters of the U.S.—Testing manual: Inland testing manual. EPA-823-B-98-004. Washington, DC: U.S. Environmental Protection Agency and U.S. Army Corps of Engineers.
- Weston, D. P., You, J., & Lydy, M. J. (2004). Distribution and toxicity of sediment-associated pesticides in agriculture-dominated water bodies of California's Central Valley. *Environmental Science & Technology*, 38, 2752–2759.

---

**Anne Jones-Lee**, PhD, is vice president of G. Fred Lee & Associates, a specialty environmental consulting firm focusing on water quality and solid and hazardous waste investigation and management. She has a bachelor's degree in biology and master's and PhD degrees in environmental science. For 11 years, she held university graduate-level professorial teaching and research positions in environmental engineering and environmental science.

**G. Fred Lee**, PhD, PE, DEE, is president of G. Fred Lee & Associates. He has a bachelor's degree in environmental health sciences, an MS degree in public health, and a PhD degree from Harvard University in environmental engineering. For 30 years, he taught and conducted research at several major U.S. universities in the areas of aquatic chemistry, water quality, solid and hazardous waste management, and environmental engineering.

Further information about the authors and their work/publications is available on their Web site, [www.gfredlee.com](http://www.gfredlee.com).

---