©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

NUMERICAL FUNCTIONAL ANALYSIS AND OPTIMIZATION Vol. 24, Nos. 3 & 4, pp. 327–338, 2003

Applications of the Best Approximation Operator to *-Nonexpansive Maps in Hilbert Spaces

N. Hussain¹ and A. R. Khan^{2,*}

 ¹Centre for Advanced Studies in Pure and Applied Mathematics, Bahauddin Zakariya University, Multan, Pakistan
²Department of Mathematical Sciences, King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia

ABSTRACT

The notion of a *-nonexpansive multivalued map is different from that of a continuous map. We give some Ky Fan type best approximation theorems for *-nonexpansive mappings defined on closed convex unbounded subsets of a Hilbert space. As applications of our theorems, we derive fixed point results under many boundary conditions. Approximating sequences to the fixed points are also constructed.

Key Words: Best approximation; Fixed point; Hilbert space; *-Nonexpansive map; Weakly inward map.

2000 AMS Subject Classification: 41A65; 46C05; 47H10; 54H25

327

DOI: 10.1081/NFA-120022926 Copyright © 2003 by Marcel Dekker, Inc. 0163-0563 (Print); 1532-2467 (Online) www.dekker.com

^{*}Correspondence A. R. Khan, Department of Mathematical Sciences, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia; E-mail: arahim@fupm.edu.sa.