

King Fahd University of Petroleum and Minerals

Research Project # MS/TRIVIAL/224

May 15, 2001 - May 14, 2002

FINAL REPORT

“TRIVIAL RING EXTENSIONS DEFINED BY
HOMOLOGICAL CONDITIONS”

Salah-Eddine Kabbaj, Professor

[External Investigator: Najib Mahdou, Associate Professor]

(Wednesday, November 13, 2002)

Contents

Abstract and Keywords	2
1 Trivial Extensions of Local Rings and Costa's First Conjecture	3
1.1 Introduction	3
1.2 Result and Example	5
1.3 Discussion	9
Bibliography	12
2 Trivial Extensions Defined by Coherent-like Conditions	14
2.1 Introduction	14
2.2 Ring-theoretic Approach	16
2.2.1 v -Coherent Rings with Zerodivisors	16
2.2.2 Results of Transfer and Examples	20
2.3 Homological Approach	24
2.3.1 n -Coherence and Strong n -Coherence	25
2.3.2 Costa's Second Conjecture	30
Bibliography	32
List of Papers	35
Acknowledgments	35

Chapter 1

Trivial Extensions of Local Rings and Costa's First Conjecture

1.1 Introduction

All rings considered in this paper are commutative with identity elements and all modules are unital. For a nonnegative integer n , an R -module E is n -presented if there is an exact sequence $F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_0 \rightarrow E \rightarrow 0$ in which each F_i is a finitely generated free R -module (In [1], such E is said to have an n -presentation). In particular, “0-presented” means finitely generated and “1-presented” means finitely presented. Also, $pd_R E$ will denote the projective dimension of E as an R -module.

In 1994, Costa [2] introduced a doubly filtered set of classes of rings throwing a brighter light on the structures of non-Noetherian rings. Namely, for nonnegative integers n and d , a ring R is an (n, d) -ring if every n -presented R -module has projective dimension at most d . The Noetherianness deflates the (n, d) -property to the notion of regular ring. However, outside Noetherian settings, the richness of this classification resides in its ability to unify classic concepts such as von Neumann regular, hereditary/Dedekind, and semi-hereditary/Prüfer rings. Costa was motivated by the sake of a deeper understanding of what makes a Prüfer domain Prüfer. In this context, he asked

“what happens if we assume only that every finitely presented (instead of generated) sub-module of a finitely generated free module is projective?” It turned out that a non-Prüfer domain having this property exists, i.e., (In the (n, d) -jargon) a $(2, 1)$ -domain which is not a $(1, 1)$ -domain. This gave rise to the theory of (n, d) -rings. Throughout, we assume familiarity with n -presentation, coherence, and basics of the (n, d) -theory as in [1, 2, 3, 6, 7, 8, 10].

Costa’s paper [2] concludes with a number of open problems and conjectures, including the existence of (n, d) -rings, specifically whether: “*There are examples of (n, d) -rings which are neither $(n, d - 1)$ -rings nor $(n - 1, d)$ -rings, for all nonnegative integers n and d* ”. Some limitations are immediate; for instance, there are no $(n, 0)$ -domains which are not fields. Also, for $d = 0$ or $n = 0$ the conjecture reduces to “ $(n, 0)$ -ring not $(n - 1, 0)$ -ring” or “ $(0, d)$ -ring not $(0, d - 1)$ -ring”, respectively.

Let’s summarize the current situation. So far, solely the cases $n \leq 2$ and d arbitrary were gradually solved in [2], [3], and [14]. These partial results were obtained using various pullbacks. For obvious reasons, these were no longer useful for the specific case $d = 0$. Therefore, in [14], the author appealed to trivial extensions of fields k by infinite-dimensional k -vector spaces, and hence constructed a $(2, 0)$ -ring (also called 2-von Neumann regular ring) which is not a $(1, 0)$ -ring (i.e., not von Neumann regular). This encouraged further work for other trivial extension contexts.

Let A be a ring and E an A -module. The trivial ring extension of A by E is the ring $R = A \ltimes E$ whose underlying group is $A \times E$ with multiplication given by $(a, e)(a', e') = (aa', ae' + a'e)$. An ideal J of R has the form $J = I \ltimes E'$, where I is an ideal of A and E' is an A -submodule of E such that $IE' \subseteq E'$. Considerable work, part of it summarized in Glaz’s book [10] and Huckaba’s book [11], has been concerned with trivial ring extensions. These have proven to be useful in solving many open problems and conjectures for various contexts in (commutative and non-commutative) ring theory. See for instance [4, 5, 9, 12, 13, 15, 16, 17].

Costa’s conjecture is still elusively outstanding. A complete solution (i.e., for all nonnegative integers n and d) would very likely appeal to new techniques and constructions. Our aim in this paper is much more modest. We shall resolve the case “ $n = 3$ and d arbitrary”. For this purpose, Section 1 investigates

the transfer of the (n, d) -property to trivial extensions of local (not necessarily Noetherian) rings by their residue fields. A surprising result establishes such a transfer and hence enables us to construct a class of $(3, d)$ -rings which are neither $(3, d - 1)$ -rings nor $(2, d)$ -rings, for d arbitrary. Section 2 is merely an attempt to show that Theorem 1.1 and hence Example 1.4 are the best results one can get out of trivial extensions of local rings by their residue fields.

1.2 Result and Example

This section develops a result on the transfer of the (n, d) -property for a particular context of trivial ring extensions, namely, those issued from local (not necessarily Noetherian) rings by their residue fields. This will enable us to construct a class of $(3, d)$ -rings which are neither $(3, d - 1)$ -rings nor $(2, d)$ -rings, for d arbitrary.

The next theorem not only serves as a prelude to the construction of examples, but also contributes to the study of the homological algebra of trivial ring extensions.

Theorem 1.2.1 *Let (A, M) be a local ring and let $R = A \rtimes A/M$ be the trivial ring extension of A by A/M . Then*

- 1) *R is a $(3, 0)$ -ring provided M is not finitely generated.*
- 2) *R is not a $(2, d)$ -ring, for each integer $d \geq 0$, provided M contains a regular element.*

The proof of this theorem requires the next preliminary.

Lemma 1.2.2 *Let A be a ring, I a proper ideal of A , and R the trivial ring extension of A by A/I . Then $\text{pd}_R(I \rtimes A/I)$ and hence $\text{pd}_R(0 \rtimes A/I)$ are infinite.*

Proof. Consider the exact sequence of R -modules

$$0 \rightarrow I \rtimes A/I \rightarrow R \rightarrow R/(I \rtimes A/I) \rightarrow 0$$

We claim that $R/(I \times A/I)$ is not projective. Deny. Then the sequence splits. Hence, $I \times A/I$ is generated by an idempotent element $(a, e) = (a, e)(a, e) = (a^2, 0)$. So $I \times A/I = R(a, 0) = Aa \times 0$, the desired contradiction (since $A/I \neq 0$). It follows from the above sequence that

$$pd_R(R/(I \times A/I)) = 1 + pd_R(I \times A/I). \quad (1)$$

Let $(x_i)_{i \in \Delta}$ be a set of generators of I and let $R^{(\Delta)}$ be a free R -module. Consider the exact sequence of R -modules

$$0 \rightarrow Ker(u) \rightarrow R^{(\Delta)} \oplus R \xrightarrow{u} I \times A/I \rightarrow 0$$

where

$$u((a_i, e_i)_{i \in \Delta}, (a_0, e_0)) = \sum_{i \in \Delta} (a_i, e_i)(x_i, 0) + (a_0, e_0)(0, 1) = \left(\sum_{i \in \Delta} a_i x_i, a_0 \right)$$

since $x_i \in I$ for each $i \in \Delta$. Hence,

$$Ker(u) = (U \times (A/I)^{(\Delta)}) \oplus (I \times A/I)$$

where $U = \{(a_i)_{i \in \Delta} \in A^{(\Delta)} / \sum_{i \in \Delta} a_i x_i = 0\}$. Therefore, we have the isomorphism of R -modules $I \times A/I \cong (R^{(\Delta)} / (U \times (A/I)^{(\Delta)})) \oplus (R/(I \times A/I))$. It follows that

$$pd_R(R/(I \times A/I)) \leq pd_R(I \times A/I). \quad (2)$$

Clearly, (1) and (2) force $pd_R(I \times A/I)$ to be infinite.

Now the exact sequence of R -modules

$$0 \rightarrow I \times A/I \rightarrow R \xrightarrow{v} 0 \times A/I \rightarrow 0,$$

where $v(a, e) = (a, e)(0, 1) = (0, a)$, easily yields $pd_R(0 \times A/I) = \infty$, completing the proof of Lemma 1.2.2. \square

Proof of Theorem 1.2.1. 1) Suppose M is not finitely generated. Let $H_0 (\neq 0)$ be a 3-presented R -module and let $(z_i)_{i=1, \dots, n}$ be a minimal set of generators of H_0 (for some positive integer n). Consider the exact sequence of R -modules

$$0 \rightarrow H_1 := Ker(u_0) \rightarrow R^n \xrightarrow{u_0} H_0 \rightarrow 0$$

where $u_0((r_i)_{i=1,\dots,n}) = \sum_{i=1}^n r_i z_i$. Throughout this proof we identify R^n with $A^n \rtimes (A/M)^n$. Our aim is to prove that $H_1 = 0$. Deny. By the above exact sequence, H_1 is a 2-presented R -module. Let $(x_i, y_i)_{i=1,\dots,m}$ be a minimal set of generators of H_1 (for some positive integer m). The minimality of $(z_i)_{i=1,\dots,n}$ implies that $H_1 \subseteq M^n \rtimes (A/M)^n$, whence $x_i \in M^n$ (and $y_i \in (A/M)^n$) for $i = 1, \dots, m$. Consider the exact sequence of R -modules

$$0 \rightarrow H_2 := \text{Ker}(u_1) \rightarrow R^m \xrightarrow{u_1} H_1 \rightarrow 0$$

where $u_1((a_i, e_i)_i) = \sum_{i=1}^m (a_i, e_i)(x_i, y_i) = \sum_{i=1}^m (a_i x_i, a_i y_i)$, since $x_i \in M^n$ for each i . Then, $H_2 = U \rtimes (A/M)^m$, where $U = \{(a_i)_{i=1,\dots,m} \in A^m / \sum_{i=1}^m a_i x_i = 0$

and $\sum_{i=1}^m a_i y_i = 0\}$. By the above exact sequence, H_2 is a finitely presented (hence generated) R -module, so that (via [11, Theorem 25.1]) U is a finitely generated A -module. Further, the minimality of $(x_i, y_i)_{i=1,\dots,m}$ yields $U \subseteq M^m$. Let $(t_i)_{i=1,\dots,p}$ be a set of generators of U and let $(f_i)_{i=p+1,\dots,p+m}$ be a basis of the (A/M) -vector space $(A/M)^m$. Consider the exact sequence of R -modules

$$0 \rightarrow H_3 := \text{Ker}(u_2) \rightarrow R^{p+m} \xrightarrow{u_2} H_2 \rightarrow 0$$

where

$$u_2((a_i, e_i)_i) = \sum_{i=1}^p (a_i, e_i)(t_i, 0) + \sum_{i=p+1}^{p+m} (a_i, e_i)(0, f_i) = \left(\sum_{i=1}^p a_i t_i, \sum_{i=p+1}^{p+m} a_i f_i \right),$$

since $t_i \in M^m$ for each $i = 1, \dots, p$ and $(f_i)_i$ is a basis of the (A/M) -vector space $(A/M)^m$. It follows that $H_3 \cong (V \rtimes (A/M)^p) \oplus (M^m \rtimes (A/M)^m)$, where $V = \{(a_i)_{i=1,\dots,p} \in A^p / \sum_{i=1}^p a_i t_i = 0\}$. By the above sequence, H_3 is a finitely generated R -module. Hence $M \rtimes A/M$ is a finitely generated ideal of R , so M is a finitely generated ideal of A by [11, Theorem 25.1], the desired contradiction.

Consequently, $H_1 = 0$, forcing H_0 to be a free R -module. Therefore, every 3-presented R -module is projective (i.e., R is a $(3, 0)$ -ring).

2) Assume that M contains a regular element m . We must show that R is not a $(2, d)$ -ring, for each integer $d \geq 0$. Let $J = R(m, 0)$ and consider the exact sequence of R -modules

$$0 \rightarrow \text{Ker}(v) \rightarrow R \xrightarrow{v} J \rightarrow 0$$

where $v(a, e) = (a, e)(m, 0) = (am, 0)$. Clearly, $\text{Ker}(v) = 0 \times (A/M) = R(0, 1)$, since m is a regular element. Therefore, $\text{Ker}(v)$ is a finitely generated ideal of R and hence J is a finitely presented ideal of R . On the other hand, $pd_R(\text{Ker}(v)) = pd_R(0 \times A/M) = \infty$ by Lemma 1.2.2, so $pd_R(J) = \infty$. Finally, the exact sequence

$$0 \rightarrow J \rightarrow R \rightarrow R/J \rightarrow 0$$

yields a 2-presented R -module, namely R/J , with infinite projective dimension (i.e., R is not a $(2, d)$ -ring, for each $d \geq 0$), completing the proof. \square

We are now able to construct a class of $(3, d)$ -rings which are neither $(3, d-1)$ -rings nor $(2, d)$ -rings, for d arbitrary. In order to do this, we first recall from [14] an interesting result establishing the transfer of the (n, d) -property to finite direct sums.

Theorem 1.2.3 ([14, Theorem 2.4]) *A finite direct sum $\bigoplus_{1 \leq i \leq n} A_i$ is an (n, d) -ring if and only if so is each A_i . \square*

Example 1.2.4 Let d be a nonnegative integer and B a Noetherian ring of global dimension d . Let (A_0, M) be a nondiscrete valuation domain and $A = A_0 \times (A_0/M)$ the trivial ring extension of A_0 by A_0/M . Let $R = A \times B$ be the direct product of A and B . Then R is a $(3, d)$ -ring which is neither a $(3, d-1)$ -ring nor a $(2, d)$ -ring, for d arbitrary (The case $d = 0$ reduces to “ $(3, 0)$ -ring not $(2, 0)$ -ring”).

Proof. By Theorem 1.2.1, A is a $(3, 0)$ -ring (also called 3-Von Neumann regular ring) which is not a $(2, d')$ -ring for each nonnegative integer d' . Moreover, R is a $(3, d)$ -ring by [14, Theorem 2.4] since both A and B are $(3, d)$ -rings

(by gnomonic theorems of Costa [2]). Further, R is not a $(2, d)$ -ring by [14, Theorem 2.4] (since A is not a $(2, d)$ -ring). Finally, we claim that R is not a $(3, d - 1)$ -ring. Deny. Then B is a $(3, d - 1)$ -ring by [14, Theorem 2.4]. Hence, by [2, Theorem 2.4] B is a $(0, d - 1)$ -ring since B is Noetherian (i.e., 0-coherent). So that $\text{gldim}(B) \leq d - 1$, the desired contradiction. \square

1.3 Discussion

This section consists of a brief discussion of the scopes and limits of our findings. This merely is an attempt to show that Theorem 1.2.1 and hence Example 1.4 are the best results one can get out of trivial extensions of local rings by their residue fields.

Remark 1.3.1 In Theorem 1.2.1, the (n, d) -property holds for a trivial ring extension of a local ring (A, M) by its residue field sans any (n, d) -hypothesis on the basic ring A . This is the first surprise. The second one resides in the narrow scope revealed by this (strong) result, namely $n = 3$ and $d = 0$. Thus, the two assertions of Theorem 1.2.1, put together with Costa's gnomonic theorems, restrict the scope of a possible example to $n = 3$ and d arbitrary.

Furthermore, since in Theorem 1.2.1 the upshot is controlled solely by restrictions on M , the next two examples clearly illustrate its failure in case one denies these restrictions, namely, " M is not finitely generated" and " M contains a regular element", respectively.

Example 1.3.2 Let K be a field and let $A = K[[X]] = K + M$, where $M = XA$. We claim that the trivial ring extension R of A by $A/M (= K)$ is not an (n, d) -ring, for any integers $n, d \geq 0$.

Proof. Let's first show that R is Noetherian. Let $J = I \times E$ be a proper ideal of R , where I is a proper ideal of A and E is a submodule of the simple A -module A/M (i.e., $E = 0$ or $E = A/M$). Since A is a Noetherian valuation ring, $I = Aa$ for some $a \in M$. Let $f \in A$ such that $(a, \bar{f}) \in J$. Without loss of generality, suppose $J \neq R(a, \bar{f})$. Let $(c, \bar{g}) \in J \setminus R(a, \bar{f})$, where $c, g \in A$, and let $c = \lambda a$, for some $\lambda \in A$. Then $(0, \bar{g} - \lambda \bar{f}) = (c, \bar{g}) - (a, \bar{f})(\lambda, \bar{0}) \in$

$J \setminus R(a, \bar{f})$, so that we may assume $c = 0$ and $\bar{g} \neq \bar{0}$, i.e., g is invertible in A . It follows that $(0, \bar{1}) = (0, \bar{g})(g^{-1}, \bar{0}) \in J$ (hence $E = A/M$) and $(a, \bar{0}) = (a, \bar{f}) - (0, \bar{g})(g^{-1}f, \bar{0}) \in J$. Consequently, $J = (a, \bar{0})R + (0, \bar{1})R$, whence J is finitely generated, as desired.

Now, by Lemma 1.2.2, $pd_R(0 \times A/M) = pd_R R(0, 1) = \infty$, hence $gldim(R) = \infty$. Then an application of [2, Theorem 1.3(ix)] completes the proof. \square

Example 1.3.3 Let K be a field and E be a K -vector space with infinite rank. Let $A = K \times E$ be the trivial ring extension of K by E . The ring A is a local $(2, 0)$ -ring by [14, Theorem 3.4]. Clearly, its maximal ideal $M = 0 \times E$ is not finitely generated and consists entirely of zero-divisors since $(0, e)M = 0$, for each $e \in E$. Let $R = A \times (A/M)$ be the trivial ring extension of A by $A/M (\cong K)$. Then R is a $(2, 0)$ -ring (and hence Theorem 1.2.1(2) fails because of the gnomonic property).

Proof. Let H be a 2-presented R -module and let (x_1, \dots, x_n) be a minimal set of generators of H . Our aim is to show that H is a projective R -module. Consider the exact sequence of R -modules

$$0 \rightarrow Ker(u) \rightarrow R^n \xrightarrow{u} H \rightarrow 0$$

where $u((r_i)_{i=1, \dots, n}) = \sum_{i=1}^n r_i x_i$. So, $Ker(u)$ is a finitely presented R -module with $Ker(u) = U \times E'$, where U is a submodule of A^n and E' is a K -vector subspace of K^n . We claim that $Ker(u) = 0$. Deny. The minimality of (x_1, \dots, x_n) yields

$$Ker(u) = U \times E' \subseteq (M \times A/M)R^n = (M \times A/M)^n$$

since R is local with maximal ideal $M \times A/M$. Let $(y_i, f_i)_{i=1, \dots, p}$ be a minimal set of generators of $Ker(u)$, where $y_i \in M^n$ and $f_i \in K^n$. Consider the exact sequence of R -modules

$$0 \rightarrow Ker(v) \rightarrow R^p \xrightarrow{v} Ker(u) (= U \times E') \rightarrow 0$$

where $v((a_i, e_i)_{i=1, \dots, p}) = \sum_{i=1}^p (a_i, e_i)(y_i, f_i) = (\sum_{i=1}^p a_i y_i, \sum_{i=1}^p a_i f_i)$. Here too the minimality of $(y_i, f_i)_{i=1, \dots, p}$ yields $Ker(v) \subseteq (M \times A/M)^p$; whence, $Ker(v) =$

$V \cong (A/M)^p$, where $V = \{(a_i)_{i=1,\dots,p} \in A^p / \sum_{i=1}^p a_i y_i = 0\} (\subseteq M^p)$. By the above exact sequence, $\text{Ker}(v)$ is a finitely generated R -module, so that V is a finitely generated A -module [11, Theorem 25.1]. Now, by the exact sequence

$$0 \rightarrow V \rightarrow A^p \xrightarrow{w} U \rightarrow 0$$

where $w((a_i)_{i=1,\dots,p}) = \sum_{i=1}^p a_i y_i$, U is a finitely presented A -module (since U is generated by $(y_i)_{i=1,\dots,p}$). Further, U is an A -submodule of A^n and A is a $(2,0)$ -ring, then U is projective. In addition, A is local, it follows that U is a finitely generated free A -module. On the other hand, $U \subseteq M^n = (0 \times E)^n$, so $(0, e)U = 0$ for each $e \in E$, the desired contradiction (since U has a basis). \square

Bibliography

- [1] N. Bourbaki, Algèbre commutative, chapitres 1-4, Masson, Paris, 1985.
- [2] D.L. Costa, Parameterizing families of non-Noetherian rings, Comm. Algebra 22 (1994) 3997-4011.
- [3] D.L. Costa, S. Kabbaj, Classes of $D + M$ rings defined by homological conditions, Comm. Algebra 24 (1996) 891-906.
- [4] R. Damiano, J. Shapiro, Commutative torsion stable rings, J. Pure Appl. Algebra 32 (1984) 21-32.
- [5] F. Dischinger, W. Müller, Left PF is not right PF, Comm. Algebra 14 (1986) 1223-1227.
- [6] D.E. Dobbs, On the global dimensions of $D + M$, Canad. Math. Bull. 18 (1975) 657-660.
- [7] D.E. Dobbs, S. Kabbaj, N. Mahdou, M. Sobrani, When is $D + M$ n -coherent and an (n, d) -domain?, Lecture Notes in Pure and Appl. Math., Dekker, 205 (1999), 257-270.
- [8] D.E. Dobbs, I.J. Papick, When is $D + M$ coherent?, Proc. Amer. Math. Soc. 56 (1976) 51-54.
- [9] R. Fossum, Commutative extensions by canonical modules are Gorenstein rings, Proc. Amer. Math. Soc. 40 (1973) 395-400.
- [10] S. Glaz, Commutative coherent rings, Lecture Notes in Mathematics, Vol. 1371, Springer-Verlag, Berlin, 1989.

- [11] J.A. Huckaba, Commutative rings with zero divisors, Marcel Dekker, New York, 1988.
- [12] G. Levin, Modules and Golod homomorphisms, *J. Pure Appl. Algebra* 38 (1985) 299-304.
- [13] C. Löfwall, On the homotopy Lie algebra of a local ring, *J. Pure Appl. Algebra* 38 (1985) 305-312.
- [14] N. Mahdou, On Costa's conjecture, *Comm. Algebra* 29 (7) (2001) 2775-2785.
- [15] I. Palmér, J.-E. Roos, Explicit formulae for the global homological dimensions of trivial extensions of rings, *J. Algebra* 27 (1973) 380-413.
- [16] D. Popescu, General Néron desingularization, *Nagoya Math. J.* 100 (1985) 97-126.
- [17] J.-E. Roos, Finiteness conditions in commutative algebra and solution to a problem of Vasconcelos, *Commutative algebra: Durham 1981* (Durham, 1981), pp. 179-203, *London Math. Soc. Lecture Note Ser.*, 72, Cambridge Univ. Press, Cambridge-New York, 1982.

Chapter 2

Trivial Extensions Defined by Coherent-like Conditions

2.1 Introduction

All rings considered in this paper are commutative with identity elements and all modules are unital. Let A be a ring and E an A -module. The trivial ring extension of A by E is the ring $R := A \ltimes E$ whose underlying group is $A \times E$ with multiplication given by $(a, e)(a', e') = (aa', ae' + a'e)$. Considerable work, part of it summarized in Glaz's book [21] and Huckaba's book (where R is called the idealization of E in A) [22], has been concerned with trivial ring extensions. These have proven to be useful in solving many open problems and conjectures for various contexts in (commutative and non-commutative) ring theory. See for instance [10, 11, 16, 23, 24, 25, 26, 27, 30, 31, 32].

A ring R is coherent if every finitely generated ideal of R is finitely presented; equivalently, if $(0 : a)$ and $I \cap J$ are finitely generated for every $a \in R$ and any two finitely generated ideals I and J of R [21]. Examples of coherent rings are Noetherian rings, Boolean algebras, von Neumann regular rings, valuation rings, and Prüfer/semihereditary rings. The concept of coherence first sprang up from the study of coherent sheaves in algebraic geometry, and then developed, under the influence of Noetherian ring theory and homology, towards a full-fledged topic in algebra. During the past 30 years, several (commutative) coherent-like notions grew out of coherence such as finite con-


ductor, quasi-coherent, v -coherent, n -coherent, and -to some extent- GCD and G-GCD rings (see the respective definitions in the beginning of Sections 2 and 3) . Noteworthy is that both the ring-theoretic and homological aspects of coherence run through most of these generalizations (see for instance [20]).

This paper investigates coherent-like conditions and related properties that might a trivial extension $R := A \times E$ inherit from the ring A for some classes of modules E . It captures previous results dealing primarily with coherence [21, 32], and also establishes satisfactory analogues of well-known coherence-like results on pullback constructions [7, 14, 12, 13, 15, 18]. Our results generate new families of examples of rings (with zerodivisors) subject to a given coherent-like condition.

The second section provides a ring-theoretic approach. We first extend the definition of a v -coherent domain to rings with zerodivisors and develop a theory of these rings parallel to Glaz's study of finite conductor, quasi-coherent, and G-GCD rings [20]. Afterwards, we study the possible transfer of all these notions for various trivial extension contexts. Thereby, new examples are provided which, particularly, enrich the current literature with new classes of coherent-like rings with zerodivisors.

The third section treats the homological aspect. We first study conditions under which trivial extensions yield (strong) n -coherent rings [8, 9, 12, 13]. Due to reciprocal effects [8, Section 2], we also deal with the (n, d) -rings of Costa, i.e., those in which n -presented modules [6] have projective dimension at most d . Particularly, the second part of this section is devoted to Costa's second conjecture which sustains that one may characterize the (n, d) -property intrinsically by ideal-theoretic-conditions [8]. We explore the scope of validity of this conjecture in various trivial extension non-coherent contexts. Recall at this point that Costa's second conjecture is valid in the class of coherent rings [9]. This fact was behind our motivation for studying large classes of coherent-like rings. The paper closes with an independent result showing that this conjecture holds in the class of finite conductor domains (resp., rings) for $n \leq 2$ and $d = 1$ (resp., $d = 0$). The general case is still elusively open.

The following diagram of commutative rings summarizes the relations between the coherent-like notions involved in this paper:



2.2 Ring-theoretic Approach

A ring R is quasi-coherent (resp., finite conductor) if $(0 : a)$ and $a_1R \cap \dots \cap a_nR$ (resp., $bR \cap cR$) are finitely generated ideals of R for any finite set of elements a and a_1, \dots, a_n (resp., b, c) of R [3, 20, 34]. Also, R is called a G-GCD ring if every principal ideal of R is projective and the intersection of any two finitely generated flat ideals of R is a finitely generated flat ideal of R [1, 20].

2.2.1 v -Coherent Rings with Zerodivisors

In view of Glaz’s recent work on finite conductor, quasi-coherent, and G-GCD rings [20], we first extend the definition of a v -coherent domain [15, 18, 28, 29] to rings with zerodivisors. For this purpose, we review some terminology related to basic operations on fractional ideals in an arbitrary ring (i.e., not necessarily a domain). Let R be a commutative ring and let $Q(R)$ denote the total ring of quotients of R . By an ideal of R we mean an integral ideal of R . Let I and J be two nonzero fractional ideals of R . We define the fractional ideal $(I : J) = \{x \in Q(R) \mid xJ \subset I\}$. We denote $(R : I)$ by I^{-1} and $(I^{-1})^{-1}$

by I_v (called the v -closure of I). A nonzero fractional ideal I is said to be invertible if $II^{-1} = R$, divisorial (or a v -ideal) if $I_v = I$, and v -finite if $I_v = J_v$ (or, equivalently, if $I^{-1} = J^{-1}$) for some finitely generated fractional ideal J of A . The v -operation on R is not necessarily a $*$ -operation, since, in general, $(a)_v \neq (a)$ when a is a zerodivisor of R . However, the other basic properties of the v -operation on integral domains [19, (32.1)(2)&(3), (32.2)(a)&(b)] carry over to arbitrary rings.

Definition 2.2.1 *A ring R is v -coherent if $(0 : a)$ and $\bigcap_{1 \leq i \leq n} Ra_i$ are v -finite ideals of R for any finite set of elements a and a_1, \dots, a_n of R .*

Proposition 2.2.2 *Let R be a ring and let's consider the following assertions:*

- (1) I^{-1} is v -finite for any finitely generated ideal I of R .
- (2) $I_v \cap J_v$ is v -finite for any two finitely generated ideals I and J of R .
- (3) $\bigcap_{1 \leq i \leq n} Ra_i$ is v -finite for any finite set of elements a_1, \dots, a_n of R .

Then (1) \implies (2). Moreover, if R is an integral domain, then the three assertions are equivalent, each of which characterizes v -coherence. \square

Proof. Assume that (1) is true and let I and J be any finitely generated ideals of R . Then there exist two finitely generated ideals I_1 and J_1 such that $I_v = I_1^{-1}$ and $J_v = J_1^{-1}$. So, $I^{-1} = (I_1)_v$ and $J^{-1} = (J_1)_v$, hence $I_v \cap J_v = (I^{-1} + J^{-1})^{-1} = ((I_1)_v + (J_1)_v)^{-1} = ((I_1)_v)^{-1} \cap ((J_1)_v)^{-1} = (I_1)^{-1} \cap (J_1)^{-1} = (I_1 + J_1)^{-1}$ which is v -finite by hypothesis since $(I_1 + J_1)^{-1}$ is a finitely generated ideal of R .

Now, assume that R is an integral domain. Then (1) \iff (2) is handled by [15, Proposition 3.6], and (1) \iff (3) always holds since $(\sum_{i=1}^n Ra_i)^{-1} = \bigcap_{1 \leq i \leq n} Ra_i^{-1}$ for each $a_i \in R$ and any integer $n \geq 1$. \square

Clearly, quasi-coherent rings are v -coherent, and if R is a domain, the above definition matches the definition of a v -coherent domain. It is worth recalling that v -coherent domains offer a large context of validity for the so-called Nagata's theorem for the class group [17]. Also, recall from [28] that PVMDs [19, 34] and Mori domains [4] are v -coherent. Moreover, non-Krull integrally closed Mori domains [2] are (v -coherent but) not finite conductor [34].

Let $(R_j)_{1 \leq j \leq m}$ be a family of rings and $R = \prod_{j=1}^m R_j$. For any $C = (c_j)$ and $A_1 = (a_{1j}), \dots, A_n = (a_{nj}) \in R$, we have $(0 : C) = \prod_{j=1}^m (0 : c_j)$ and

$\bigcap_{1 \leq i \leq n} RA_i = \prod_{j=1}^m (\bigcap_{1 \leq i \leq n} Ra_i)$. Further, for any ideals $I = \prod_{j=1}^m I_j$ and $J = \prod_{j=1}^m J_j$ of R , we have $I \cap J = \prod_{j=1}^m (I_j \cap J_j)$ and $I^{-1} = \prod_{j=1}^m I_j^{-1}$. Then $\prod_{j=1}^m R_j$ is v -coherent if and only if so is R_j for each $j = 1, \dots, m$. Thus, finite products (for instance, of Mori domains) may provide us with original examples of v -coherent rings with zerodivisors.

Let's now examine v -coherence for rings of small weak dimension. Recall first that rings of weak dimension 0 are precisely the von Neumann regular rings. Moreover, Glaz showed that for a ring R of weak dimension 1 the finite conductor property, the quasi-coherence, and coherence deflate to the mere fact that $(0 : c)$ is finitely generated for every $c \in R$ [20, Proposition 2.2]. She also proved that the finite conductor and quasi-coherence properties coincide for rings of weak dimension 2 [20, Theorem 2.3]. In contrast with these results, the next example denies any similar effect to the weak dimension on v -coherence.

Example 2.2.3 Let E be a countable direct sum of copies of $\mathbb{Z}/2\mathbb{Z}$ with addition and multiplication defined component wise, where \mathbb{Z} is the ring of integers. Let $R = \mathbb{Z} \times E$ with multiplication defined by $(a, e)(b, f) = (ab, af + be + ef)$. Then:

- (1) $\text{w.dim}(R) = 1$.
- (2) R is not coherent.
- (3) R is a v -coherent ring.

Proof. (1) That $\text{w.dim}(R) = 1$ this is handled in [33, Example 1.3, page 10].
 (2) Let $x = (2, 0) \in R$. Then $(0 : x) = \{(a, e) \in R \mid (a, e)(2, 0) = 0\} = \{(a, e) \in R \mid (2a, 0) = 0\} = 0 \times E$ which is not a finitely generated ideal of R . Therefore, R is not a coherent ring.
 (3) Notice first that an element $s \in R$ is regular if and only if $s = (a, 0)$ with $a \in \mathbb{Z} \setminus 2\mathbb{Z}$. This easily follows from the four basic facts: E is Boolean; $2E = 0$; $ae = e$ for any $a \in \mathbb{Z} \setminus 2\mathbb{Z}$ and $e \in E$; and for any $e \neq 0 \in E$, there exists $f \neq 0 \in E$ such that $ef = 0$.

Next, we wish to show that each ideal of R is v -finite which implies that R is v -coherent. Let J be an ideal of R and let $I = \{a \in \mathbb{Z} \mid (a, e) \in J \text{ for some } e \in E\}$. Assume $I = 0$. Let s be any regular element of R . Clearly, $(0, e) = s(0, e)$ for any $e \in E$. It follows that $sJ = J$ and hence $J^{-1} = Q(R) = (R(0, e))^{-1}$

for any $e \neq 0 \in E$. Now, assume $I = x\mathbb{Z}$, where x is a nonzero integer. We claim that $J^{-1} = (R(x, 0))^{-1}$. Indeed, let $y/s \in Q(R)$, where $y = (a, e) \in R$ and $s = (b, 0)$ is a regular element. It can easily be seen that $sR = b\mathbb{Z} \times E$. Then $y/s \in J^{-1} \Leftrightarrow yJ \subseteq sR \Leftrightarrow (a, e)J \subseteq b\mathbb{Z} \times E \Leftrightarrow aI \subseteq b\mathbb{Z} \Leftrightarrow ax \in b\mathbb{Z} \Leftrightarrow (a, e)(R(x, 0)) \subseteq sR \Leftrightarrow y/s \in (R(x, 0))^{-1}$. Thus, in both cases, J is v -finite, as asserted. \square

While a ring R which is a total ring of quotients trivially is v -coherent, R need not be finite conductor [20, Example 3.5]. The following construction may offer new contexts that illustrate this fact.

Example 2.2.4 Let (R, M) be any local ring with $M^2 = 0$. Then:

- (1) R is a v -coherent ring that is not G-GCD.
- (2) The following conditions are equivalent:
 - (i) R is a coherent ring;
 - (ii) R is a quasi-coherent ring;
 - (iii) R is a finite conductor ring;
 - (iv) $(0 : c)$ is finitely generated for every $c \in R$;
 - (v) M is finitely generated.

Proof. (1) That R is v -coherent this is straightforward since $R = Q(R)$ is a total ring of quotients. Let $c \neq 0 \in M$. Then $\text{Ann}(c) = (0 : c) = M$. Hence Rc is not projective (since not free), so that R is not a G-GCD ring [20].

(2) Clearly, we only need prove $(v) \implies (i)$. Assume that M is finitely generated and let I be a finitely generated proper ideal of R . Let $\{x_1, \dots, x_n\}$ be a minimal generating set of I and consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(u) \rightarrow R^n \xrightarrow{u} I \rightarrow 0$$

where $u(a_1, \dots, a_n) = \sum_{i=1}^n a_i x_i$. We claim that $\text{Ker}(u) = \prod M =: M^n$. Indeed, $M^n \subseteq \text{Ker}(u)$ is clear since $M^2 = 0$ and $x_i \in M$ for each $i = 1, \dots, n$. On the other hand, $\text{Ker}(u) \subseteq M^n$ since $\{x_1, \dots, x_n\}$ is minimal. Therefore, $\text{Ker}(u) = M^n$ is a finitely generated R -module (since M is). Hence, I is finitely presented and thus R is coherent. \square

2.2.2 Results of Transfer and Examples

This subsection investigates the possible transfer of the coherence properties for various trivial extension contexts. Our results generate new families of examples subject to a given coherent-like condition.

For the convenience of the reader, we next discuss some basic facts connected to trivial ring extensions. These will be used frequently in the sequel without explicit mention. Let A be a ring and E an A -module and let $R := A \times E$ be the trivial ring extension of A by E . An ideal of R of the form $I \times IE$, where I is an ideal of A , is finitely generated if and only if so is I [21, page 141]. Also recall that R has always its Krull dimension equal to the Krull dimension of A [22, Theorem 25.1(3)].

For a general description of modules over a trivial ring extension, we refer the reader to [21, pages 140 & 141]. Here, we describe a specific type of R -modules that play a crucial role within the R -module structure, namely, finitely generated free R -modules and their R -submodules. Let n be a positive integer. Define the “multiplication” on E by elements of A^n within E^n through the natural A -bilinear map $\varphi : A^n \times E \rightarrow E^n$ defined by $ae = \varphi(a, e) := (a_i e)_{1 \leq i \leq n}$, for any $a = (a_i)_{1 \leq i \leq n} \in A^n$ and $e \in E$. Now let U be an A -submodule of A^n and E' an A -submodule of E^n such that $UE \subseteq E'$. Let $U \times E'$ denote the set $U \times E'$ with natural addition and scalar multiplication defined by $(a, e)(u, e') = (au, ae' + ue)$. Clearly, $U \times E'$ is an R -module; and, under this notation, the finitely generated free R -module R^n identifies with $A^n \times E^n$. Also, $U \times E'$ is a finitely generated R -module only if U is a finitely generated A -module. Conversely, let M be an R -submodule of R^n . Set $U := \{u \in A^n \mid (u, e') \in M \text{ for some } e' \in E^n\}$ and $E' := \{e' \in E^n \mid (u, e') \in M \text{ for some } u \in A^n\}$. It is easily seen that U and E' are A -modules such that $M \subseteq U \times E'$. The next example illustrates the fact that the equality does not hold in general.

Example 2.2.5 Let (A, M) be a local domain which is not a field, $E := A/M$, and $R := A \times E$ be the trivial ring extension of A by E . Let $J = R(x, 1)$, where $x \neq 0 \in M$. Set $I = \{a \in A \mid (a, e) \in J \text{ for some } e \in E\}$ and $E' = \{e \in E \mid (a, e) \in J \text{ for some } a \in A\}$. Then $J \subsetneq I \times E'$.

Proof. One may easily check that $I = Ax$ and $E' = E$. Further, we claim that $(x, 0) \in I \times E' \setminus J$. Deny. We have $(x, 0) = (a, e)(x, 1)$ for some $(a, e) \in R$ so that $x = ax$. Hence $a = 1 \in M$, the desired contradiction. \square

Nevertheless, it is easily seen that $M = U \times E'$ if and only if $0 \times E' \subseteq M$ if and only if $U \times 0 \subseteq M$.

Example 2.2.5 shows that [22, Theorem 25.1(1)] fails to be true in general. (This was confirmed by the author of [22] through a private e-communication.)

Theorem 2.2.6 *Let (A, M) be a local ring and E an A -module with $ME = 0$. Let $R := A \times E$ be the trivial ring extension of A by E . Then:*

- (1) *R is a v -coherent ring that is not G -GCD.*
- (2) *R is coherent (resp., quasi-coherent, finite conductor) if and only if so is A , M is finitely generated, and E is an (A/M) -vector space of finite rank.*

Before proving Theorem 2.2.6, we establish the following Lemma.

Lemma 2.2.7 *Under the hypotheses of Theorem 2.2.6, $(0 : c)$ is a finitely generated ideal of R for each $c \in R$ if and only if $(0 : a)$ is a finitely generated ideal of A for each $a \in A$, M is finitely generated, and E is an (A/M) -vector space of finite rank.*

Proof. Assume that $(0 : c)$ is a finitely generated ideal of R for each $c \in R$. Let $a \in A$ and set $c := (a, 0) \in R$. Then $(0 : c) = (0 : a) \times E'$, where $E' = \{e \in E | ae = 0\}$. Therefore, $(0 : a)$ is a finitely generated ideal of A . Let $e \neq 0 \in E$ and set $c := (0, e) \in R$. Similar arguments show that M is a finitely generated ideal of A since $(0 : c) = M \times E$. Further, assume that $M \times E = \sum_{i=1}^n R(x_i, e_i)$, where $x_i \in M$ and $e_i \in E$ for each $i = 1, \dots, n$. Then $E \subseteq \sum_{i=1}^n (A/M)e_i$, and hence E is an (A/M) -vector space of finite rank.

Conversely, let $c := (a, e) \neq 0 \in R$. If a is invertible in A , then c is invertible in R . Then, without loss of generality, we may assume that $a \in M$. Hence, $(0 : c) = \{(b, f) \in R | (ab, be) = 0\} = \{(b, f) \in M \times E | ab = 0\}$ (since if b is invertible in A , then (b, f) is invertible in R , and so $c = 0$, a contradiction). It can easily be seen that if $a = 0$ then $(0 : c) = M \times E$, and if $a \neq 0$ then

$(0 : c) = (0 : a) \otimes E$. In both cases, $(0 : c)$ is a finitely generated ideal of R since M and $(0 : a)$ are finitely generated ideals of A and E is an (A/M) -vector space of finite rank, completing the proof of Lemma 2.2.7. \square

Proof of Theorem 2.2.6. (1) Similar arguments as in Example 2.2.4(1) lead to the conclusion.

(2) Assume that R is a coherent ring. By Lemma 2.2.7, it remains to show that A is coherent. Let $I = \sum_{i=1}^n Aa_i$, where $a_i \in M$ and set $J := \sum_{i=1}^n R(a_i, 0)$. Consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(u) \rightarrow R^n = A^n \otimes E^n \xrightarrow{u} J \rightarrow 0$$

where $u((b_i, e_i)_{1 \leq i \leq n}) = \sum_{i=1}^n (b_i, e_i)(a_i, 0) = (\sum_{i=1}^n a_i b_i, 0)$ since $a_i \in M$ for each $i = 1, \dots, n$. On the other hand, consider the exact sequence of A -modules:

$$0 \rightarrow \text{Ker}(v) \rightarrow A^n \xrightarrow{v} I \rightarrow 0$$

where $v((b_i)_{1 \leq i \leq n}) = \sum_{i=1}^n a_i b_i$. Hence, $\text{Ker}(u) = \text{Ker}(v) \otimes E^n$. But J is finitely presented since R is coherent, so $\text{Ker}(u)$ is a finitely generated R -module and hence $\text{Ker}(v)$ is a finitely generated A -module. Therefore, I is a finitely presented ideal of A , so A is coherent.

Conversely, let J be a finitely generated ideal of R and let $S := \{(a_i, e_i)\}_{1 \leq i \leq n}$ be a minimal generating set of J , where $a_i \in M$ and $e_i \in E$. Consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(u) \rightarrow R^n \xrightarrow{u} J \rightarrow 0$$

where $u((b_i, f_i)_{1 \leq i \leq n}) = \sum_{i=1}^n (a_i, e_i)(b_i, f_i) = (\sum_{i=1}^n a_i b_i, \sum_{i=1}^n b_i e_i)$ since $a_i \in M$ for each $i = 1, \dots, n$. Further, the minimality of S yields $\text{Ker}(u) = \{(b_i, f_i)_{1 \leq i \leq n} \in R^n \mid \sum_{i=1}^n a_i b_i = 0\}$. Let $I := \sum_{i=1}^n Aa_i$ and consider the surjective A -module homomorphism v defined above. Then $\text{Ker}(v)$ is a finitely generated A -module since A is coherent. Consequently, $\text{Ker}(u) = \text{Ker}(v) \otimes E^n$ is a finitely generated R -module. Therefore, J is finitely presented and hence R is coherent.

Now, assume that R is quasi-coherent. We only need show that $\bigcap_{1 \leq i \leq n} Ra_i$ is a finitely generated ideal of A for each $a_i \in M$. This is straightforward since $\bigcap_{1 \leq i \leq n} R(a_i, 0) = (\bigcap_{1 \leq i \leq n} Aa_i) \otimes 0$ is a finitely generated ideal of R .

Conversely, let $J = \bigcap_{1 \leq i \leq n} R(a_i, e_i)$, where $a_i \in M$ and $e_i \in E$. We may suppose that $J \not\subseteq R(a_i, e_i)$ for each $i = 1, \dots, n$. Let $(a, e) \in J$. Then, there exist $b_i \in A$ and $f_i \in E$ such that $(a, e) = (b_i, f_i)(a_i, e_i) = (a_i b_i, b_i e_i)$ for each $i = 1, \dots, n$. We claim that $b_i \in M$ for each $i = 1, \dots, n$. Deny. Then, there exists j such that b_j is invertible in A and so is (b_j, f_j) in R . Hence $(a_j, e_j) = (b_j, f_j)^{-1}(a, e) \in J$ yielding $J = R(a_j, e_j)$, a contradiction. Therefore, $(a, e) = (a_i b_i, 0)$. It follows that $J = \bigcap_{1 \leq i \leq n} (Aa_i \times 0) = (\bigcap_{1 \leq i \leq n} Aa_i) \times 0$ is finitely generated in R since $\bigcap_{1 \leq i \leq n} Aa_i$ is by hypothesis finitely generated in A . Thus R is quasi-coherent.

Finally, similar arguments as above with $n = 2$ lead to the conclusion for the finite conductor property, to complete the proof of Theorem 2.2.6. \square

Next, we explore a different context, namely, the trivial ring extension of a domain by its quotient field.

Theorem 2.2.8 *Let A be a domain which is not a field, $K = \text{qf}(A)$, and $R := A \times K$ be the trivial ring extension of A by K . Then:*

- (1) *R is not a finite conductor ring. In particular, R is neither quasi-coherent nor coherent.*
- (2) *R is a v -coherent ring if and only if so is A .*

Proof. (1) Let $x := (0, 1) \in R$. Then $(0 : x) = 0 \times K$ is not a finitely generated ideal of R . Therefore, R is not a finite conductor ring, as asserted.

(2) Remark first that $(a, e) \in R$ is regular if and only if $a \neq 0$, and that $(0 : c) = 0 \times K$ for any $c := (0, e) \neq 0 \in R$. Further, [22, Theorem 25.1(4)] yields $\bigcap_{1 \leq i \leq n} R(a_i, e_i) = \bigcap_{1 \leq i \leq n} (Ra_i \times K) = (\bigcap_{1 \leq i \leq n} Ra_i) \times K$, for every finite set of elements $(a_i, e_i)_{1 \leq i \leq n}$ of R (with $a_i \neq 0$ for each i). Now, Let I be any ideal of A and E any A -submodule of K with $IK \subseteq E$ and let $J := I \times E$. By [22, Theorem 25.10], we have $J^{-1} = (I \times E)^{-1} = I^{-1} \times K = (I \times IK)^{-1}$. Finally, since $I \times IK$ is finitely generated if I is, the conclusion is straightforward. \square

New examples of original coherent-like rings with zerodivisors with arbitrary Krull dimensions may stem from Theorems 2.2.6 & 2.2.8, as shown by the following constructions.

Example 2.2.9 Let K be any field and X_1, X_2, \dots be indeterminates over K . Let n be any integer ≥ 1 , $A = K[[X_1, \dots, X_n]]$ the power series ring in n variables over K , and $R := A \rtimes K$. Then, by Theorem 2.2.6, R is an n -dimensional coherent ring that is not G-GCD. \square

Example 2.2.10 Let A be as in the above example and $R := A \rtimes K[Y]$, where Y is another indeterminate over K . Then, by Theorem 2.2.6, R is an n -dimensional v -coherent ring that is not finite conductor. \square

Example 2.2.11 Let $R := \mathbb{Z}[X_1, \dots, X_{n-1}] \rtimes \mathbb{Q}(X_1, \dots, X_{n-1})$, where n is any integer ≥ 1 , \mathbb{Z} the ring of integers, and \mathbb{Q} the field of rational numbers. Then, by Theorem 2.2.8, R is an n -dimensional v -coherent ring that is not finite conductor. \square

More examples are provided in the next section, denying any possible interplay between some of these coherent-like conditions and n -coherence.

2.3 Homological Approach

For a nonnegative integer n , an R -module E is n -presented if there is an exact sequence $F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_0 \rightarrow E \rightarrow 0$ in which each F_i is a finitely generated free R -module [6]. In particular, “0-presented” means finitely generated and “1-presented” means finitely presented. Throughout, $\text{pd}_R(E)$ and $\text{fd}_R(E)$ will denote the projective dimension and the flat dimension of E as an R -module, respectively. Also $\text{w.dim}(R)$ will denote the weak dimension of R .

In 1994, Costa introduced a doubly filtered set of classes of rings, called the (n, d) -rings, with the aim of obtaining a good understanding of the structures of some non-Noetherian rings [8]. The Noetherianity forces the regularity of the (n, d) -rings. However, outside Noetherian settings, the richness of this classification resides in its ability to unify classic concepts such as von Neumann regular, hereditary, Dedekind, semihereditary, and Prüfer rings.

Given nonnegative integers n and d , a ring R is called an (n, d) -ring if every n -presented R -module has projective dimension $\leq d$; and a weak (n, d) -ring if

every n -presented cyclic R -module has projective dimension $\leq d$ (equivalently, if every $(n - 1)$ -presented ideal of R has projective dimension $\leq d - 1$). For instance, the $(0, 1)$ -domains are the Dedekind domains, the $(1, 1)$ -domains are the Prüfer domains, and the $(1, 0)$ -rings are the von Neumann regular rings [8]. Costa's paper concludes with a number of open problems, including his second conjecture sustaining that the (n, d) - and weak (n, d) -properties are equivalent. This conjecture is valid in the class of coherent rings [9].

The first part of this section studies the transfer of the n -coherence properties (see definitions below) to trivial ring extensions. Due to reciprocal effects [8, Section 2], results of transfer for the (n, d) -property are also provided. Our purpose, in the second part, is mainly to test the validity of Costa's second conjecture for non-coherent contexts.

2.3.1 n -Coherence and Strong n -Coherence

Recall from [12, 13], for $n \geq 1$, that R is n -coherent if each $(n - 1)$ -presented ideal of R is n -presented; and that R is strong n -coherent if each n -presented R -module is $(n + 1)$ -presented (This terminology is not the same as that of [8], where Costa's " n -coherence" is our "strong n -coherence"). In particular, "1-coherence" coincides with coherence, and one may view "0-coherence" as Noetherianity. Any strong n -coherent ring is n -coherent, and the converse holds for $n = 1$ or for coherent rings [13, Proposition 3.3]. Strong n -coherence arose naturally in Costa's study [8] of the (n, d) -rings. As a matter of fact, every (n, d) -ring is strong $\max\{n, d\}$ -coherent [8, Theorem 2.2]; and an (n, d) -ring is strong r -coherent ($r < n$) only if it is an (r, d) -ring [8, Theorem 2.4].

Our main result examines the context of trivial ring extensions of domains by their respective quotient fields.

Theorem 2.3.1 *Let A be a domain which is not a field, $K = \text{qf}(A)$, and $R := A \times K$ be the trivial ring extension of A by K . Let $n \geq 2$ and $d \geq 1$ be integers. Then the following hold:*

- (1) R is not coherent.

- (2) R is strong n -coherent (resp., n -coherent) if and only if so is A .
 (3) R is an (n, d) -ring (resp., a weak (n, d) -ring) if and only if so is A .

The proof of this theorem lies mainly on the following two lemmas which characterize, respectively, finitely generated and n -presented R -submodules of free R -modules.

Let us fix the notation for the next two results. Let R be as in Theorem 3.1 and let H be an R -submodule of R^m , where m is an arbitrary positive integer. Set $U = \{x \in A^m / (x, e) \in H \text{ for some } e \in K^m\}$ and $E = \{e \in K^m / (x, e) \in H \text{ for some } x \in A^m\}$.

Lemma 2.3.2 *Under the above notation, the following statements are equivalent:*

- (i) H is finitely generated and E is a K -vector space;
 (ii) U is finitely generated and $H = U \rtimes KU$.

Proof. $i) \implies ii)$ Let $H = \sum_{i=1}^p R(x_i, e_i) (\subseteq U \rtimes E)$, where p is a positive integer, $x_i \in A^m$, and $e_i \in K^m$ for each $i = 1, \dots, p$. Necessarily, $U = \sum_{i=1}^p Ax_i$ and $E = \sum_{i=1}^p Ae_i + KU$. Next assume $KU \subsetneq E$. Then there exists a nonzero K -vector space F with finite rank such that $F \oplus KU = E$. Write $e_i = y_i + z_i$, where $y_i \in F$ and $z_i \in KU$ for each $i = 1, \dots, p$. From above, it follows that $E = \sum_{i=1}^p Ay_i \oplus KU$ and thus $F = \sum_{i=1}^p Ay_i$. Consequently, F (and hence K) is a finitely generated A -module, the desired contradiction. Hence, $E = KU$. Now let $y \in E = KU$. Then $y = \sum_{i=1}^p b_i x_i$, where $b_i \in K$ for each $i = 1, \dots, p$. So $(0, y) = \sum_{i=1}^p (0, b_i)(x_i, e_i) \in H$. It follows that $0 \rtimes E \subseteq H$; equivalently, $H = U \rtimes KU$.

$ii) \implies i)$ Straightforward. \square

Lemma 2.3.3 *Let n be an integer ≥ 1 . Under the above notation, the following statements are equivalent:*

- (i) H is n -presented;
 (ii) U is n -presented and $H = U \rtimes KU$.

Proof. $i) \implies ii)$ By induction on n . Assume $n = 1$. As above, write $H = \sum_{i=1}^p R(x_i, f_i)$, where p is a positive integer, $x_i \in A^m$, and $f_i \in K^m$ for each

$i = 1, \dots, p$. We have $U = \sum_{i=1}^p Ax_i$ and $E = (\sum_{i=1}^p Af_i) + KU$. Let F' be a K -vector space such that $F' \oplus KU = K^m$. Hence, $f_i = g_i + h_i$, where $g_i \in F'$ and $h_i \in KU$ for each $i = 1, \dots, p$. So that $E = (\sum_{i=1}^p Ag_i) \oplus KU$. Further, it easily can be seen that $H = \sum_{i=1}^p R(x_i, g_i)$. Consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(w) \rightarrow R^p \xrightarrow{w} H \rightarrow 0$$

where $w((a_i, e_i)_{1, \dots, p}) = \sum_{i=1}^p (a_i, e_i)(x_i, g_i) = (\sum_{i=1}^p a_i x_i, \sum_{i=1}^p a_i g_i + \sum_{i=1}^p e_i x_i)$. Set $W := \{(a_i)_{i=1, \dots, p} \in A^p / \sum_{i=1}^p a_i x_i = 0 \text{ and } \sum_{i=1}^p a_i g_i = 0\}$ and $E' := \{(e_i)_{i=1, \dots, p} \in K^p / \sum_{i=1}^p e_i x_i = 0\}$. Clearly, E' is a K -vector space and $\text{Ker}(w)$ is a finitely generated R -submodule of R^p (since H is finitely presented). By Lemma 3.2, W is finitely generated and $\text{Ker}(w) = W \times KW$. Moreover, let $(a_i)_{i=1, \dots, p} \neq 0 \in A^p$ such that $\sum_{i=1}^p a_i x_i = 0$. Then, $(a_i)_{i=1, \dots, p} \in E' = KW$. So there exists $z \neq 0 \in K$ and $(b_i)_{i=1, \dots, p} \in W$ such that $(a_i)_{i=1, \dots, p} = z(b_i)_{i=1, \dots, p}$. Hence $\sum_{i=1}^p a_i g_i = z \sum_{i=1}^p b_i g_i = 0$, whence $(a_i)_{i=1, \dots, p} \in W$. Consequently, $W = \{(a_i)_{i=1, \dots, p} \in A^p / \sum_{i=1}^p a_i x_i = 0\}$. Therefore, the exact sequence of A -modules of natural homomorphisms

$$0 \rightarrow W \rightarrow A^p \rightarrow U \rightarrow 0,$$

upon tensoring by the flat A -module R , yields the exact sequence of R -modules

$$0 \rightarrow W \otimes_A R \cong W \times KW = \text{Ker}(w) \rightarrow R^p \rightarrow U \otimes_A R \cong U \times KU \rightarrow 0.$$

It follows that U is finitely presented and $H = U \times KU$.

The inductive step is carried out just as we did for the case $n = 1$ above, provided one substitutes the induction hypothesis for Lemma 2.3.2.

ii) \implies i) Straightforward. \square

Proof of Theorem 2.3.1. (1) is already handled by Theorem 2.2.8(1). Specifically, $R(0, 1) = 0 \times A$ is a finitely generated ideal of R which is not finitely presented (by Lemma 2.3.3).

(2) and (3) follow readily from a combination of Lemma 2.3.3 with the next three facts:

- (a) R is a faithfully flat A -module.
- (b) For $n \geq 2$, a ring B is n -coherent if and only if every $(n - 1)$ -presented

submodule of a finitely generated free B -module is n -presented.

(c) For $n \geq 2$ and $d \geq 1$, a ring B is an (n, d) -ring if and only if every $(n - 1)$ -presented submodule of a finitely generated free B -module has projective dimension $\leq d - 1$. \square

For $n \leq 1$ or $d = 0$, the (n, d) -property may not survive, in general, in the trivial extension R (even under strong assumption on A). This is illustrated by the next example.

Example 2.3.4 Let A be any arbitrary Prüfer domain (i.e., $(1, 1)$ -domain) and let R be the trivial ring extension of A by its quotient field. Then R is a $(2, 1)$ -ring which is neither a semihereditary ring (i.e., $(1, 1)$ -ring) nor a 2-von Neumann regular ring (i.e., $(2, 0)$ -ring).

Proof. That R is a $(2, 1)$ -ring which is not a $(1, 1)$ -ring, this is ensured by Theorem 2.3.1(3)&(1), respectively. Now, let $J := R(a, 0)$, where a is a non-zero non-invertible element of A . Since $(a, 0)$ is a regular element of R , then the ideal J of R has no non-zero annihilator. By [26, Theorem 2.1], R is not a 2-von Neumann regular ring. \square

The Bézout property, however, does transfer reciprocally from A to R , as shown by the next result.

Proposition 2.3.5 *Let R be as in Theorem 2.3.1. Then R is a Bézout ring if and only if A is a Bézout domain.*

Proof. Assume R is a Bézout ring and let I be a finitely generated proper ideal of A . Then $J := I \times IK = I \times K$ is a finitely generated ideal of R . So $J = R(a, e)$ for some $a \in A$ and $e \in E$. Therefore, $I = Aa$ and hence A is a Bézout domain.

Conversely, let J be a finitely generated proper ideal of R . Set $I := \{a \in A / (a, e) \in J \text{ for some } e \in K\}$. We next envisage two cases. CASE 1: $I = 0$. Necessarily, $J = 0 \times (1/b)L$ for some $b \neq 0 \in A$ and some finitely generated proper ideal L of A . Further, $L = Aa$ since A is a Bézout domain. Hence $J = 0 \times A(a/b) = R(0, a/b)$, as desired. CASE 2: $I \neq 0$. Let $(a, e) \in J$ such

that $a \neq 0$. Then, $(a, e)(0 \times K) = 0 \times K \subseteq J$; equivalently, $J = I \times IK = I \times K$. But $I = Aa$ for some $a \in A$ since A is a Bézout domain. Therefore, $J = I \times K = R(a, 0)$, completing the proof. \square

Noteworthy is that new families of examples of non-semihereditary Bézout rings stem from the combination of Example 2.3.4 and Proposition 2.3.5.

At this point, for the convenience of the reader, we recall from [23] the main result that establishes the transfer of the (n, d) -property to trivial ring extensions of local rings by their respective residue fields.

Theorem 2.3.6 [23, Theorem 1.1] *Let (A, M) be a local ring and let $R := A \times A/M$ be the trivial ring extension of A by A/M . Then:*

- (1) *R is a $(3, 0)$ -ring provided M is not finitely generated.*
- (2) *R is not a $(2, d)$ -ring, for each integer $d \geq 0$, provided M contains a regular element.* \square

Clearly, Theorems 2.3.1 and 2.3.6 generate original examples of n -coherent rings which, moreover, reflect no obvious correlation between (strong) n -coherence and the large class of finite conductor rings.

Example 2.3.7 Let \mathbb{Z} be the ring of integers and $\mathbb{Q} = \text{qf}(\mathbb{Z})$. Then $R := \mathbb{Z} \times \mathbb{Q}$ is a strong 2-coherent ring which is not a finite conductor ring.

Proof. Straightforward via Theorem 2.3.1 and Theorem 2.2.8. \square

Example 2.3.8 Let (V, M) be a nondiscrete valuation domain. Then $R := V \times V/M$ is a 3-coherent ring which is neither 2-coherent nor a finite conductor ring.

Proof. R is a $(3, 0)$ -ring by Theorem 2.3.6(1) (since M is not a finitely generated ideal of A). Hence R is 3-coherent. Further R is not a $(2, 0)$ -ring by Theorem 2.3.6(2). So by [8, Theorem 2.4] R is not 2-coherent. Finally, Theorem 2.2.6(2) ensures that R is not a finite conductor ring. \square

2.3.2 Costa's Second Conjecture

A well-known fact about semihereditary rings is that a ring R is a $(1, 1)$ -ring if and only if it is a weak $(1, 1)$ -ring. In this vein, Costa's second conjecture wonders whether the (n, d) -property and the weak (n, d) -property are equivalent for any non-negative integers n and d . So far, it has been shown that this conjecture is valid under the coherence assumption [9]. It remains however elusively open, in general.

Our modest objective in this subsection is mainly to test its validity beyond the class of coherent rings. In this line, two results are stated generating two new contexts of validity for this conjecture. The first of these involves trivial ring extensions issued from coherent domains.

Theorem 2.3.9 *Let A be a non-trivial coherent domain, $K = \text{qf}(A)$, and $R := A \times K$. Then, for any integers $n \geq 2$ and $d \geq 1$, R is a non-coherent ring such that the following statements are equivalent:*

- (i) R is an (n, d) -ring;
- (ii) R is a weak (n, d) -ring;
- (iii) $\text{w.dim}(A) \leq d$.

Proof. (i) \iff (ii) is a straightforward application of Theorem 2.3.1(1)&(3) combined to [9, Proposition 2].

(i) \implies (iii) Assume R is an (n, d) -ring. By Theorem 2.3.1(3) and [8, Theorem 2.4], A is a $(1, d)$ -domain, hence $\text{w.dim}(A) \leq d$ by [21, Theorem 1.3.9].

(iii) \implies (i) Assume that $\text{w.dim}(A) \leq d$. Let $M := I \times K$ be any arbitrary maximal ideal of R , where I is a maximal ideal of A (Cf. [22, Theorem 25.1]). We have $\text{fd}_A(I) \leq d - 1$ and so $\text{fd}_R(M) \leq d - 1$ (since $M \cong I \otimes R$ and R is A -flat). By [8, Theorem 4.1], R is a $(d + 1, d)$ -ring. Further, by Theorem 2.3.1(2), R is strong 2-coherent. It follows that R is a $(2, d)$ -ring (by [8, Theorem 2.4]) and hence a (n, d) -ring, as desired. \square

Recall that a ring R is finite conductor if any ideal I of R with $\mu(I) \leq 2$ is finitely presented, where $\mu(I)$ denotes the cardinality of a minimal set of generators of I [20, Proposition 2.1]. Our next (and last) theorem tests Costa's second conjecture in the class of finite conductor rings. As might be expected,

the “ $\mu(I) \leq 2$ ” assumption (in the above definition) restricts the scope of this result to $n = 2$ and $d \leq 1$.

Theorem 2.3.10 *Let R be a finite conductor ring. Then:*

(1) *The following statements are equivalent:*

- (i) *R is a Prüfer domain;*
- (ii) *R is a $(2, 1)$ -domain;*
- (iii) *R is a weak $(2, 1)$ -domain.*

(2) *The following statements are equivalent:*

- (i) *R is a von Neumann regular ring;*
- (ii) *R is a $(2, 0)$ -ring;*
- (iii) *R is a weak $(2, 0)$ -ring.*

Proof. (1) We need only prove (iii) \implies (i). Suppose that (iii) holds. Let I be an arbitrary 2-generated ideal of R (i.e., $\mu(I) \leq 2$). Then I is finitely presented, and hence projective by (iii). Therefore R is a Prüfer domain by [21, Theorem 1.4.1].

(2) We need only prove (iii) \implies (i). Suppose that (iii) holds. It suffices to show that each principal ideal of R is a direct summand of R . Let I be a principal ideal of R . Then I is finitely presented by hypothesis, so that R/I is a 2-presented cyclic R -module, and hence projective by (iii). Therefore the following exact sequence splits:

$$0 \rightarrow I \rightarrow R \rightarrow R/I \rightarrow 0$$

leading to the conclusion. \square

Remark 2.3.11 Assertion (1) of Theorem 2.3.10 cannot extend to rings with zerodivisors. Indeed, let $R := \mathbb{Z} \times E$ as in Example 2.2.3. Then R is a finite conductor ring which is not a semihereditary ring (since it is not coherent). On the other hand, by [8, Theorem 4.5], R is a $(2, 1)$ -ring since $w\dim(R) = 1$. \square

Bibliography

- [1] D.D. Anderson, D.F. Anderson, Generalized GCD domains, *Comment. Math. Univ. St. Paul.* 28 (1980), no. 2, 215–221.
- [2] V. Barucci, On a class of Mori domains, *Comm. Algebra* 11 (1983), no. 17, 1989–2001.
- [3] V. Barucci, D.F. Anderson, D.E. Dobbs, Coherent Mori domains and the principal ideal theorem, *Comm. Algebra* 15 (1987), no. 6, 1119–1156.
- [4] V. Barucci, S. Gabelli, How far is a Mori domain from being a Krull domain?, *J. Pure Appl. Algebra* 45 (1987), no. 2, 101–112.
- [5] N. Bourbaki, *Algèbre homologique*, Chapitre 10, Masson, Paris, 1980.
- [6] N. Bourbaki, *Commutative algebra*, Chapters 1-7, Springer, Berlin, 1998.
- [7] W. Brewer, E. Rutter, $D+M$ constructions with general overrings, *Michigan Math. J.* 23 (1976), no. 1, 33–42.
- [8] D. Costa, Parameterizing families of non-Noetherian rings, *Comm. Algebra* 22 (1994), no. 10, 3997–4011.
- [9] D. Costa, S. Kabbaj, Classes of $D + M$ rings defined by homological conditions, *Comm. Algebra* 24 (1996), no. 3, 891–906.
- [10] R. Damiano, J. Shapiro, Commutative torsion stable rings, *J. Pure Appl. Algebra* 32 (1984) 21–32.
- [11] F. Dischinger, W. Müller, Left PF is not right PF, *Comm. Algebra* 14 (1986) 1223-1227.

- [12] D.E. Dobbs, S. Kabbaj, N. Mahdou, n -coherent rings and modules, Lecture Notes in Pure and Appl. Math., Dekker, 185 (1997) 269–281.
- [13] D.E. Dobbs, S. Kabbaj, N. Mahdou, M. Sobrani, When is $D + M$ n -coherent and an (n, d) -domain?, Lecture Notes in Pure and Appl. Math., Dekker, 205 (1999) 257–270.
- [14] D.E. Dobbs, I. Papick, When is $D + M$ coherent? Proc. Amer. Math. Soc. 56 (1976) 51–54.
- [15] M. Fontana, S. Gabelli, On the class group and the local class group of a pullback, J. Algebra 181 (1996), no. 3, 803–835.
- [16] R. Fossum, Commutative extensions by canonical modules are Gorenstein rings, Proc. Amer. Math. Soc. 40 (1973) 395–400.
- [17] S. Gabelli, On Nagata’s theorem for the class group, II, Lecture Notes in Pure and Appl. Math., Dekker, 206 (1999) 117–142.
- [18] S. Gabelli, E. Houston, Coherentlike conditions in pullbacks, Michigan Math. J. 44 (1997), no. 1, 99–123.
- [19] R. Gilmer, Multiplicative ideal theory, Pure and Applied Mathematics, No. 12. Marcel Dekker, Inc., New York, 1972.
- [20] S. Glaz, Finite conductor rings, Proc. Amer. Math. Soc. 129 (10)(2000), no. 1, 2833–2843.
- [21] S. Glaz, Commutative coherent rings, Lecture Notes in Mathematics, 1371, Springer-Verlag, Berlin, 1989.
- [22] J.A. Huckaba, Commutative rings with zero divisors, Marcel Dekker, New York-Basel, 1988.
- [23] S. Kabbaj, N. Mahdou, Trivial extensions of local rings and a conjecture of Costa, Lecture Notes in Pure and Appl. Math., Dekker, 231 (2003) 301–311.

- [24] G. Levin, Modules and Golod homomorphisms, *J. Pure Appl. Algebra* 38 (1985) 299–304.
- [25] C. Löfwall, On the homotopy Lie algebra of a local ring, *J. Pure Appl. Algebra* 38 (1985) 305–312.
- [26] N. Mahdou, On Costa’s conjecture, *Comm. Algebra* 29 (7) (2001) 2775–2785.
- [27] N. Mahdou, Steinitz properties in trivial extensions of commutative rings, *Arab. J. Sci. Eng. Sect. C Theme Issues* 26 (2001), no. 1, 119–125.
- [28] D. Nour El Abidine, Sur le groupe des classes d’un anneau intègre, *C. R. Math. Rep. Acad. Sci. Canada* 13 (1991), no. 2-3, 69–74.
- [29] D. Nour El Abidine, Sur un théorème de Nagata, *Comm. Algebra* 20 (1992), no. 7, 2127–2138.
- [30] I. Palmér, J.-E. Roos, Explicit formulae for the global homological dimensions of trivial extensions of rings, *J. Algebra* 27 (1973) 380–413.
- [31] D. Popescu, General Néron desingularization, *Nagoya Math. J.* 100 (1985) 97–126.
- [32] J. E. Roos, Finiteness conditions in commutative algebra and solution of a problem of Vasconcelos, In: *London Math. Soc. Lectures Notes in Mathematics*, 72 (1981), 179–204.
- [33] W.V. Vasconcelos, *The Rings of Dimension Two*, Marcel Dekker, New York, 1976.
- [34] M. Zafrullah, On finite conductor domains, *Manuscripta Math.* 24 (1978), no. 2, 191–204.

List of Papers

- [1] S. Kabbaj, N. Mahdou, Trivial extensions of local rings and a conjecture of Costa, Lecture Notes in Pure and Appl. Math., Dekker, 231 (2003) 301–311.
- [2] S. Kabbaj, N. Mahdou, Trivial extensions defined by coherent-like conditions (Submitted for publication).

Acknowledgments

This work has been funded by King Fahd University of Petroleum and Minerals under Project # MS/TRIVIAL/224.