



King Fahd University of Petroleum & Minerals

**DEPARTMENT OF MATHEMATICAL SCIENCES**

---

Technical Report Series

TR 267

June 2001

**Decay of solutions of a nonlinear hyperbolic system  
describing heat propagation by second sound**

Messaoudi A. Salim

# Decay of solutions of a nonlinear hyperbolic system describing heat propagation by second sound.

Messaoudi A. Salim

Mathematical Sciences Department,

KFUPM, Dhahran 31261,

Saudi Arabia.

E-mail : messaoud@kfupm.edu.sa

## Abstract

In this work we establish a decay result for the solutions of a nonlinear hyperbolic system describing heat propagation, where the heat flux is given by Cattaneo's law.

**Keywords** heat, second sound, nonlinear, hyperbolic, exponential decay.

**AMS Subject Classification** 35L45 - 35K05 - 35K65.

## 1 Introduction

In the absence of deformation and external sources, the evolution of the heat flux and the absolute temperature is given by the system

$$\begin{aligned}q + \kappa(\theta)\theta_x &= 0 \\ q_x + c(\theta)\theta_t &= 0,\end{aligned}\tag{1.1}$$

where  $\kappa$  and  $c$  are strictly positive functions characterizing the material in consideration. In the case where  $c$  and  $\kappa$  are independent of  $\theta$ , we get the familiar linear heat equation

$$\theta_t = k\theta_{xx}, \quad k = \frac{\kappa}{c}.$$

This equation provides a useful description of heat conduction under a large range of conditions and predicts an infinite speed of propagation; that is, any thermal disturbance at one point has an instantaneous effect elsewhere in the body. This is not always the case. In fact, experiments showed that heat conduction in some dielectric crystals at low temperatures is free of this paradox (infinite speed propagation) and

disturbances which are almost entirely thermal may propagate in a finite speed. This phenomenon in dielectric crystals is called second sound.

These observations go back to 1948, when Cattaneo [1] proposed, instead of Fourier's law, a new constitutive relation

$$\tau(\theta)q_t + q = -\kappa(\theta)\theta_x, \quad (1.2)$$

where  $\tau$  and  $\kappa$  are strictly positive functions depending on the absolute temperature. Coleman, Fabrizio, and Owen [2] showed in 1982 that, if (1.2) is adopted then compatibility with thermodynamics requires that the internal energy be given by

$$e = \tilde{e}(\theta, q) = a(\theta) + b(\theta)q^2, \quad (1.3)$$

where  $b$  is a function determined by  $\tau$  and  $\kappa$ . In particular  $b(\theta) > 0$ . In this case the system governing the evolution of  $\theta$  and  $q$  takes the form

$$\begin{aligned} q_x + (a'(\theta) + b'(\theta)q^2)\theta_t + 2b(\theta)qq_t &= 0 \\ \tau(\theta)q_t + q + \kappa(\theta)\theta_x &= 0. \end{aligned} \quad (1.4)$$

Global existence and decay of classical solutions to the Cauchy problem, as well as to some initial boundary value problems, have been established by Coleman, Hrusa, and Owen [3]. They also showed that  $(\theta, q)$  tends to the equilibrium state, however no rate of decay has been discussed. Concerning the formation of singularities, Messaoudi [4] studied the following system

$$\begin{aligned} \tau(\theta)q_t + q + \kappa(\theta)\theta_x &= 0 \\ c(\theta)\theta_t + q_x &= 0 \end{aligned}$$

and showed, under the same restrictions on  $\tau$ ,  $c$  and  $\kappa$ , that classical solutions to the Cauchy problem break down in finite time if the initial data are chosen small in the  $L^\infty$  norm with large enough derivatives. This result has been improved later and established by the author [5] for an equivalent system of the form

$$\begin{aligned} \sigma(e, q)q_t + \mu(e, q)q &= -e_x + \lambda(e, q)qq_x \\ e_t &= -q_x, \end{aligned} \quad (1.5)$$

where  $\sigma, \mu$  satisfy

$$\sigma(\xi, \eta) \geq \underline{\sigma} > 0, \quad \mu(\xi, \eta) \geq \underline{\mu} > 0, \quad \forall(\xi, \eta) \in \mathbb{R}^2. \quad (1.6)$$

For the derivation of (1.5) from (1.4) and the proof of the global existence, we refer the reader to [5] and [6].

In this work we consider (1.5) together with the initial and boundary conditions

$$\begin{aligned} e(x, 0) &= e_0(x), \quad q(x, 0) = q_0(x), \quad x \in I = (0, 1) \\ q(0, t) &= q(1, t) = 0, \quad t \geq 0 \end{aligned}$$

and show that global classical solutions decay exponentially if the initial data are sufficiently small.

## 2 Exponential Decay

In this section we state and prove our main result. For this purpose we set

$$\hat{e} = e - e_1, \quad e_1 = \int_0^1 e_0(x) dx.$$

It is clear that  $(\hat{e}, q)$  satisfies the following problem

$$\sigma(\hat{e}, q)q_t + \mu(\hat{e}, q)q = -\hat{e}_x + \lambda(\hat{e}, q)qq_x \quad (2.1)$$

$$\hat{e}_t = -q_x, \quad (2.2)$$

$$\hat{e}(x, 0) = e_0(x), \quad q(x, 0) = q_0(x), \quad x \in I \quad (2.3)$$

$$q(0, t) = q(1, t) = 0, \quad t \geq 0, \quad (2.4)$$

where

$$\sigma(\hat{e}, q) = \sigma(\hat{e} + e_1, q), \mu(\hat{e}, q) = \mu(\hat{e} + e_1, q), \lambda(\hat{e}, q) = \lambda(\hat{e} + e_1, q).$$

**Remark 2.1 :** In the sequel we go back to the notation  $e$  instead of  $\hat{e}$ .

**Remark 2.2 :** By using (2.2), we easily see that  $\hat{e}$  satisfies Poincaré's inequality.

**Theorem.** *Assume that  $\sigma, \mu, \lambda$  are  $C^2$  functions satisfying (1.6). Then there exists a small positive constant  $\delta$  such that for any  $e_0$  in  $H^2(I)$  and  $q_0$  in  $H^2(I) \cap H_0^1(I)$  satisfying*

$$\|e_0\|_2^2 + \|q_0\|_2^2 < \delta^2, \quad (2.5)$$

*the solution of (2.1) - (2.4) decays exponentially as  $t \rightarrow +\infty$ .*

In order to carry out the proof, we consider another problem which agrees with (3.1) - (3.4) when  $(e, q)$  are close enough to the equilibrium state  $(0, 0)$ . For this purpose, we introduce the functions  $A, B, C$  satisfying the following hypotheses

h1)  $A, B, C \in C_b^2(\mathbb{R}^2)$

h2)  $A(\xi, \eta) = \sigma(\xi, \eta), \quad B(\xi, \eta) = \mu(\xi, \eta), \quad C(\xi, \eta) = \lambda(\xi, \eta),$   
 $\forall (\xi, \eta) \in \mathcal{V}$  a neighborhood of  $(0, 0)$

h3)  $A(\xi, \eta) \geq \underline{A} > 0, \quad B(\xi, \eta) \geq \underline{B} > 0.$

Here  $C_b^2$  denotes the space of continuous and bounded functions, as well as, their first and second order derivatives. We note that functions with these properties can be constructed by virtue of (1.6). Therefore, instead of (3.1) - (3.4) we consider the following problem

$$A(e, q)q_t + B(e, q)q = -e_x + qC(e, q)q_x \quad (2.6)$$

$$e_t = -q_x \quad x \in I, \quad t \geq 0 \quad (2.7)$$

$$e(x, 0) = e_0(x), \quad q(x, 0) = q_0(x), \quad x \in I. \quad (2.8)$$

$$q(0, t) = q(1, t) = 0, \quad t \geq 0, \quad (2.9)$$

**Remark 2.3.** Any solution  $(e, q)$  to (2.6) – (2.9) satisfying  $(e, q) \in \mathcal{V}$  is also a solution to (2.1) – (2.4) by virtue of (h2).

We also set

$$\begin{aligned} \mathcal{E}(t) : &= \int_0^1 (e^2 + e_t^2 + e_x^2 + e_{xt}^2 + e_{xx}^2 + e_{tt}^2 + q^2 + q_t^2 \\ &\quad + q_x^2 + q_{xt}^2 + q_{xx}^2 + q_{tt}^2)(x, t) dx \end{aligned} \quad (2.10)$$

$$\begin{aligned} \Lambda(t) : &= \int_0^1 [e^2 + e_t^2 + e_x^2 + e_{xt}^2 + e_{tt}^2](x, t) dx \\ &\quad + \int_0^1 A(e, q)[q^2 + q_t^2 q_x^2 + q_{xt}^2 + q_{tt}^2](x, t) dx \end{aligned} \quad (2.11)$$

$$\alpha(t) := \sup_{0 \leq x \leq 1} [(|e| + |e_x| + |e_t| + |q| + |q_t| + |q_x|)(x, t)] \quad (2.12)$$

**Proof.**

We multiply (2.6) by  $q$  and (2.7) by  $e$ , integrate over  $I$ , use integration by parts, and add equalities, to obtain

$$\frac{1}{2} \frac{d}{dt} \int_0^1 [Aq^2 + e^2](x, t) dx \leq - \int_0^1 Bq^2(x, t) dx + \Gamma \alpha(t) \mathcal{E}(t), \quad (2.13)$$

where  $\Gamma$  denotes a positive (possibly large) generic constant independent of  $e, q, t$ .

To get the next estimates, we differentiate (2.6), (2.7) with respect to  $t$

$$Aq_{tt} + A_t q_t + Bq_t + B_t q = -e_{xt} + C_t q q_x + C q q_{xt} + C q_t q_x \quad (2.14)$$

$$e_{tt} = -q_{xt} \quad (2.15)$$

and multiply (2.14) by  $q_t$  and (2.15) by  $e_t$ . Similar computations as above then yield

$$\frac{1}{2} \frac{d}{dt} \int_0^1 [Aq_t^2 + e_t^2](x, t) dx \leq - \int_0^1 Bq_t^2(x, t) dx + \Gamma (\alpha(t) + \alpha^2(t)) \mathcal{E}(t). \quad (2.16)$$

To establish bounds on terms involving  $e_{tt}$  and  $q_{tt}$ , we introduce the difference operator as follows : for  $h > 0$ , we set

$$\Delta_h W(x, t) = W(x, t+h) - W(x, t), \quad x \in \mathbb{R}, \quad t \geq 0 \quad (2.17)$$

We apply the above operator to equations (2.14), (2.15), multiply the resulting equalities by  $\Delta_h q_t$  and  $\Delta_h e_t$  respectively, integrate over  $I$ , and add the inequalities. After a number of integrations, using integration by parts, we divide by  $h^2$  and let  $h$  go to zero. Thus we get

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_0^1 [Aq_{tt}^2 + e_{tt}^2](x, t) dx &\leq - \int_0^1 Bq_{tt}^2(x, t) dx \\ &\quad + \Gamma (\alpha(t) + \alpha^2(t) + \alpha^3(t)) \mathcal{E}(t). \end{aligned} \quad (2.18)$$

For additional estimates, we differentiate (2.6), (2.7) with respect to  $x$  to get

$$Aq_{xt} + A_x q_t + Bq_x + B_x q = -e_{xx} + C_x q q_x + C q q_{xx} + C q_x^2 \quad (2.19)$$

$$e_{xt} = -q_{xx}. \quad (2.20)$$

We then multiply (2.19) by  $q_x$  and (2.20) by  $e_x$  to obtain, by similar calculations,

$$\frac{1}{2} \frac{d}{dt} \int_0^1 [Aq_x^2 + e_x^2](x, t) dx \leq - \int_0^1 Bq_x^2(x, t) dx + \Gamma (\alpha(t) + \alpha^2(t)) \mathcal{E}(t). \quad (2.21)$$

Again we apply the operator (2.17) to equations (2.19), (2.20), multiply the resulting equalities by  $\Delta_h q_x$  and  $\Delta_h e_x$  respectively, integrate over  $I$ , and add the inequalities. After a number of integrations, using integration by parts, we divide by  $h^2$  and let  $h$  go to zero to arrive at

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_0^1 [Aq_{xt}^2 + e_{xt}^2](x, t) dx &\leq - \int_0^1 Bq_{xt}^2(x, t) dx \\ &+ \Gamma (\alpha(t) + \alpha^2(t) + \alpha^3(t)) \mathcal{E}(t). \end{aligned} \quad (2.22)$$

By combining (2.13), (2.16), (2.18), (2.21), (2.22), we get

$$\begin{aligned} \Lambda'(t) &\leq - \int_0^1 B (q^2 + q_t^2 + q_x^2 + q_{tt}^2 + q_{xt}^2) (x, t) dx \\ &+ \Gamma (\alpha(t) + \alpha^2(t) + \alpha^3(t)) \mathcal{E}(t). \end{aligned} \quad (2.23)$$

Next we show that, for  $(e, q)$ ,  $\Lambda$  is equivalent to  $\mathcal{E}$ . For this purpose we use equations (2.6), (2.7), (2.14), (2.15), (2.19), (2.20) and the hypotheses (h1) - (h3). Thus, we obtain

$$c_1 (1 + \alpha^2(t) + \alpha^4(t)) \mathcal{E}(t) \leq \Lambda(t) \leq c_2 \mathcal{E}(t) \quad (2.24)$$

where  $c_1, c_2$  are constants. A combination of (2.23) and (2.24) then yields

$$\Lambda'(t) \leq - \int_0^1 B (q^2 + q_t^2 + q_x^2 + q_{tt}^2 + q_{xt}^2) (x, t) dx + \Gamma \alpha(t) \Lambda(t). \quad (2.25)$$

Next we exploit (2.6), (2.7), (2.14), (2.15), (2.19), (2.20) for further estimates

$$\int_0^1 (e_t^2 + e_{tt}^2) (x, t) dx = \int_0^1 (q_x^2 + q_{xt}^2) (x, t) dx \quad (2.26)$$

and

$$\int_0^1 (e_x^2 + e_{xt}^2) (x, t) dx \leq c \int_0^1 (q^2 + q_t^2 + q_{tt}^2) (x, t) dx + \Gamma \alpha(t) \Lambda(t). \quad (2.27)$$

We then use Poincaré's inequality and (2.6) to obtain

$$\int_0^1 e^2(x, t) dx \leq c \int_0^1 (q^2 + q_t^2) (x, t) dx + \Gamma \alpha(t) \Lambda(t). \quad (2.28)$$

By combining (2.25) - (2.28) and using (h3) we conclude

$$\Lambda'(t) \leq -a\Lambda(t) + \Gamma\alpha(t)\Lambda(t). \quad (2.29)$$

where  $a$  is a constant depending only on the upper and lower bounds of  $A, B, C$ . We also note that, by the standard Sobolev embedding inequalities we have  $\alpha(t) \leq \sqrt{2\mathcal{E}(t)}$ . So by choosing  $\delta$  in (2.5) so small that  $\Gamma\alpha(0) < a/2$  and  $(e, q) \in \mathcal{V}$ , the relation (2.29) yields

$$\Lambda'(t) < -\frac{a}{2}\Lambda(t), \quad \forall t \in [0, \varepsilon) \quad (2.30)$$

Direct integration then leads to

$$\Lambda(t) \leq \Lambda(0)e^{-at/2}, \quad \forall t \in [0, \varepsilon). \quad (2.31)$$

Since  $\Lambda(t) \leq \Lambda(0)$  we extend (2.31) beyond  $\varepsilon$ . By repeating the same procedure and using the continuity of  $\Lambda$ , (2.31) is established for all  $t \geq 0$ . This completes the proof.

**Acknowledgement** The author would like to thank KFUPM for its sincere support.

#### References

1. Cattaneo, C., Sulla conduzione del calore, *Atti Sem. Math. Fis Univ. Modena* **3** (1948), 83-101.
2. Coleman, B.D., M. Fabrizio, and D.R. Owen, On the thermodynamics of second sound in dielectric crystals, *Arch. Rational Mech. Analysis* **80** (1982), 135-158.
3. Coleman, B.D., W.J. Hrusa, and D.R. Owen, Stability of equilibrium for a nonlinear hyperbolic system describing heat propagation by second sound in solids, *Arch. Rational Mech. Anal.* **94** (1986), 267-289.
4. Messaoudi, S.A., Formation of singularities in heat propagation guided by second sound, *J. Diff. Eqns.* **130** (1996), 92-99.
5. Messaoudi S. A., On the existence and nonexistence of solutions of a nonlinear hyperbolic system describing heat propagation by second sound, *Applicable Analysis* **73** (1999) 485 - 496
6. Messaoudi Salim, On the Solution of a Hyperbolic Heat System, *Proceedings of the First Saudi Science Conference*, Vol. **3**, (2001), 547 - 556.