

Problem 2. (17pts)

6 pts a) Find an equation of the plane that contains the line

$$x = 10 + t, \quad y = 9 - 5t, \quad z = t$$

and is perpendicular to the plane $3x + 2y - 2z = 7$.

Let $\vec{u} = \langle 1, -5, 1 \rangle$ and $\vec{n} = \langle 3, 2, -2 \rangle$.

A normal vector to the plane is given by

$$\vec{u} \times \vec{n} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -5 & 1 \\ 3 & 2 & -2 \end{vmatrix} = \langle 8, 5, 17 \rangle.$$

3 pts

$(10, 9, 0)$ is a point on that plane. Hence,

the equation of the plane is

$$8(x-10) + 5(y-9) + 17(z-0) = 0 \Leftrightarrow$$

$$\underline{8x + 5y + 17z = 125}$$

2 pts

11 pts b) Identify and sketch the surface given by


$$-4x^2 + 8x + y^2 + 2y - \frac{4z^2}{9} = 7$$

4 pts The standard form of the equation is (after completing the squares)

$$-\frac{(x-1)^2}{1^2} + \frac{(y+1)^2}{2^2} - \frac{z^2}{3^2} = 1$$

3 pts This is a hyperboloid of two sheets. (Like $-\frac{x^2}{1^2} + \frac{y^2}{2^2} - \frac{z^2}{3^2} = 1$, except y is shifted by -1 and x is shifted by 1).

graph.



4 pts

R₂ * general shape = 2 pts
* shifting = 1 pt
* axe = 1 pt

Problem 3. (16pts)

8pts a) Show that the function

$$f(x, y) = \begin{cases} \frac{\sin(xy)}{x^2+y^2} & \text{if } (x, y) \neq (0, 0) \\ 1 & \text{if } (x, y) = (0, 0) \end{cases}$$

is not continuous at $(0, 0)$

$$f(0, 0) = 1.$$

4pts * Along the x-axis ($y=0, x \neq 0$), $f(x, y) = 0$.
 so $f(x, y) \rightarrow 0$ as $(x, y) \rightarrow (0, 0)$ along the x-axis.

* Along the line $y=x$ ($x \neq 0, y \neq 0$), $f(x, y) = \frac{\sin x^2}{2x^2}$
 $f(x, y) \rightarrow \frac{1}{2}$ as $(x, y) \rightarrow (0, 0)$ along the curve $y=x$

Thus $\lim_{(x, y) \rightarrow (0, 0)} f(x, y)$ does not exist.

f is not continuous at $(0, 0)$.

Rk.

* choice of a curve and value of limit along that curve is ~~is~~ 4pts
 * limit $\neq f(0, 0)$ 4pts

8pts b) Evaluate

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{3 - \sqrt{x^2 + y^2 + 9}}{x^2 + y^2}$$

Set $x = r \cos \theta$
 $y = r \sin \theta$

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{3 - \sqrt{x^2 + y^2 + 9}}{x^2 + y^2} =$$

Rk. of the conjugate or L'Hospital rule is applied for a specific curve, give only 2pts.

$$= \lim_{r \rightarrow 0} \frac{3 - \sqrt{r^2 + 9}}{r^2}$$

$$= \lim_{r \rightarrow 0^+} \frac{9 - (r^2 + 9)}{r^2 (3 + \sqrt{r^2 + 9})} \quad 4pts$$

$$= \lim_{r \rightarrow 0} \frac{-1}{3 + \sqrt{r^2 + 9}} = \underbrace{-\frac{1}{6}}_{1pt}$$


3pts

Problem 4. (17pts)

7pts a) Find and sketch the domain of the function

$$f(x, y) = \sqrt{2x + y} \ln(y - x).$$

The domain $D = \left\{ (x, y) \mid \underbrace{2x + y \geq 0}_{2pts}, \underbrace{y > x}_{2pts} \right\}$



10pts b) Use implicit differentiation to find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ if

$$yz = \ln(x + z).$$

1pt $\frac{\partial z}{\partial x} = ?$
 $\rightarrow \frac{\partial}{\partial x} (yz) = \frac{\partial}{\partial x} [\ln(x+z)] \Leftrightarrow$

2pts $\rightarrow y \frac{\partial z}{\partial x} = \frac{1 + \frac{\partial z}{\partial x}}{x+z} \Leftrightarrow$

$y(x+z) \frac{\partial z}{\partial x} = 1 + \frac{\partial z}{\partial x} \Leftrightarrow$

1pt $\rightarrow \frac{\partial z}{\partial x} [y(x+z) - 1] = 1 \Leftrightarrow$

1pt $\rightarrow \frac{\partial z}{\partial x} = \frac{1}{xy + yz - 1}$

$\frac{\partial z}{\partial y} = ?$
 1pt $\frac{\partial}{\partial y} (yz) = \frac{\partial}{\partial y} (\ln(x+z)) \Leftrightarrow$

$z + y \frac{\partial z}{\partial y} = \frac{\frac{\partial z}{\partial y}}{x+z} \Leftrightarrow$

$z(x+z) + y(x+z) \frac{\partial z}{\partial y} = \frac{\partial z}{\partial y} \Leftrightarrow$

$\frac{\partial z}{\partial y} (1 - yx - yz) = z(x+z) \Leftrightarrow$

$\frac{\partial z}{\partial y} = \frac{z(x+z)}{1 - yx - yz}$

Problem 5. (16pts)

8pts a) Find an equation of the tangent plane to the surface

$$z = xye^{-xy^2}$$

at the point $(1, 1, e^{-1})$.

$$\begin{aligned} \frac{\partial z}{\partial x} &= y e^{-xy^2} + (xy)(-y^2) e^{-xy^2} \\ &= y e^{-xy^2} - xy^3 e^{-xy^2} \end{aligned} \quad \left. \vphantom{\frac{\partial z}{\partial x}} \right\} 2 \text{pts}$$

$$\frac{\partial z}{\partial x} \Big|_{(1,1)} = e^{-1} - e^{-1} = 0 \quad 1 \text{pt}$$

$$\frac{\partial z}{\partial y} = x e^{-xy^2} + (xy)(-2xy) e^{-xy^2} = (x - 2xy^2) e^{-xy^2} \quad 2 \text{pts}$$

$$\frac{\partial z}{\partial y} \Big|_{(1,1)} = (1 - 2) e^{-1} = -e^{-1} \quad 1 \text{pt}$$

An equation of the tangent line is thus given by

$$z = e^{-1} + 0(x-1) + (-e^{-1})(y-1) \quad (\Rightarrow)$$

$$z = \frac{1}{e} - \frac{1}{e}(y-1) \quad (\Rightarrow) \quad ez + y = 2 \quad 1 \text{pt}$$

8pts b) Use the linearization $L(x,y)$ of the function

$$f(x,y) = \sqrt{9 - x^2 - 4y^2}$$

at $(-1, 1)$ to find the best estimate for $f(-0.9, 1.1)$.

$$f_x = \frac{-2x}{2\sqrt{9-x^2-4y^2}} = \frac{-x}{\sqrt{9-x^2-4y^2}}, \quad f_x(-1,1) = \frac{1}{2} \quad 1 \text{pt}$$

$$f_y = \frac{-8y}{2\sqrt{9-x^2-4y^2}} = \frac{-4y}{\sqrt{9-x^2-4y^2}}, \quad f_y(-1,1) = -2 \quad 1 \text{pt}$$

$$f(-1,1) = 2 \quad L(x,y) = 2 + \frac{1}{2}(x+1) - 2(y-1) \quad 2 \text{pts}$$

$(-0.9, 1.1)$ is close to $(-1, 1) \Rightarrow$

$$f(-0.9, 1.1) \approx L(-0.9, 1.1) = 2 + \frac{1}{2}(0.1) - 2(0.1)$$

$$= 1.85 \quad 1 \text{pt}$$

Rk. give 2pts for the general formula of $L(x,y)$.

Problem 6. (17pts)

8pts a) Find all second partial derivatives of

$$f(x, y) = xe^{-2y}$$

$$\underline{f_x = e^{-2y}} \quad \text{2pts}, \quad \underline{f_y = -2xe^{-2y}} \quad \text{2pts}$$

$$\underline{f_{xx} = 0} \quad \text{1pt}, \quad \underline{f_{yy} = 4xe^{-2y}} \quad \text{1pt}$$

$$\underline{f_{xy} = -2e^{-2y} = f_{yx}} \quad \text{2pts}$$

8pts b) If $z = y + \cos(x^2 - y^2)$, show that

$$y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y} = x$$

$$\underline{\frac{\partial z}{\partial x} = -2x \sin(x^2 - y^2)} \quad \text{3pts}, \quad \underline{\frac{\partial z}{\partial y} = 1 + 2y \sin(x^2 - y^2)} \quad \text{3pts}$$

$$\underline{y \frac{\partial z}{\partial x} = -2xy \sin(x^2 - y^2)} \quad \text{1pt}, \quad \underline{x \frac{\partial z}{\partial y} = x + 2yx \sin(x^2 - y^2)} \quad \text{1pt}$$

$$y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y} = \cancel{-2xy \sin(x^2 - y^2)} + \cancel{x + 2yx \sin(x^2 - y^2)}$$

$$= x \quad \text{1pt}$$