

MATH 302-2
Quiz 1, Term 101

NAME:

ID :

Instructor: Prof.Dr. Othman Echi

Exercise 1. Let $U = (1, 1, 0)$ and $V = (1, 1, \sqrt{6})$.

- (1) Evaluate the angle between the two vectors U and V .
- (2) Evaluate the area of the parallelogram defined by the vectors U and V .

Exercise 2. Let $U = (1, 1, -1)$, $V = (-1, 1, 2)$ and

$$S = \{W = (x, y, z) \in \mathbb{R}^3 \mid U \cdot W = 0 \text{ and } V \cdot W = 0\}.$$

- (1) Show that S is a subspace of \mathbb{R}^3 .
- (2) Find a basis of S .

SOLUTIONS

Ex. 1. The angle between two nonzero vectors U, V is the real number $\theta \in [0, \pi]$ such that

$$\cos(\theta) = \frac{U \cdot V}{\|U\| \times \|V\|}.$$

The area of the parallelogram defined by the two vectors is

$$\mathcal{A}(U, V) = \|U\| \times \|V\| \sin(\theta) \text{ (unit}^2\text{)}$$

In our case, $\cos(\theta) = \frac{1}{2}$. Hence $\theta = \frac{\pi}{3}$ and consequently,

$$\mathcal{A}(U, V) = \sqrt{2} \times \sqrt{8} \times \frac{\sqrt{3}}{2} = 2\sqrt{3} \text{ (unit}^2\text{)}.$$

□

Ex. 2.

1.

– As $U \cdot 0_{\mathbb{R}^3} = V \cdot 0_{\mathbb{R}^3} = 0$, we get $0_{\mathbb{R}^3} \in S$.

– Let W_1, W_2 be two elements of S . Then

$$U \cdot W_1 = V \cdot W_1 = U \cdot W_2 = V \cdot W_2 = 0.$$

This leads to

$$U \cdot (W_1 + W_2) = U \cdot W_1 + U \cdot W_2 = 0 + 0 = 0,$$

and

$$V \cdot (W_1 + W_2) = V \cdot W_1 + V \cdot W_2 = 0 + 0 = 0.$$

It follows that $W_1 + W_2 \in S$.

– Now, let $W \in S$ and $\alpha \in \mathbb{R}$. Then

$$U \cdot (\alpha W) = \alpha U \cdot W = \alpha \times 0 = 0,$$

and

$$V \cdot (\alpha W) = \alpha V \cdot W = \alpha \times 0 = 0.$$

This implies that $\alpha W \in S$.

Therefore, S is a subspace of \mathbb{R}^3 .

2. Let $W = (x, y, z) \in \mathbb{R}^3$. Then $W \in S$ if and only if the following equations are satisfied

$$x + y - z = 0 \text{ and } -x + y + 2z = 0.$$

This gives $z = -y$ and $x = -3y$. Thus, $W = (x, y, z) = (-3y, y, -2y) = (-y)(3, -1, 2)$. We conclude that $\mathcal{B} = \{(3, -1, 2)\}$ is a basis of S . □