

KING FAHD UNIVERSITY OF PETROLEUM AND  
MINERALS  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
MATH 102 (123)  
FINAL EXAM

Name:.....ID:.....Section:....

**Exercise 1:** The integral

$$\int_{-2}^2 (x^5 + 3)\sqrt{4 - x^2} dx$$

is equal to

- (a)  $6\pi$
- (b)  $3\pi$
- (c)  $\frac{\pi}{2}$
- (d)  $3 + 2\pi$
- (e)  $5 + 2\pi$

**Exercise 2:** If

$$F(x) = \int_1^{x^2} \frac{1}{2\sqrt{t}} \tan^{-1}(\sqrt{t}) dt, \quad x > 0$$

then  $F(1) + F'(1) + F''(1)$  is equal to

- (a)  $\frac{\pi+2}{4}$
- (b)  $\frac{\pi-2}{4}$
- (c)  $\frac{2\pi-1}{2}$
- (d)  $\frac{\pi+1}{2}$
- (e) 0

**Exercise 3:** The integral

$$\int \frac{1}{x^2} \sqrt{2 - \frac{1}{x}} dx$$

is equal to

(a)  $\frac{2}{3} \left[2 - \frac{1}{x}\right]^{3/2} + C$

(b)  $3 \left[2 - \frac{1}{x}\right]^{1/3} + C$

(c)  $2 \left[2 - \frac{1}{x}\right]^{1/2} + C$

(d)  $\frac{1}{3} \left[2 - \frac{1}{x}\right]^{3/2} + C$

(e)  $\left[2 - \frac{1}{x}\right]^{3/2} + C$

**Exercise 4:** The area of the region below the line  $y = 1$  and between the curves  $y = \tan x$  and  $x = 0$  is equal to

(a)  $\frac{1}{4}(\pi - 2 \ln 2)$

(b)  $\frac{1}{4}(\pi + \ln 2)$

(c)  $\frac{\pi}{4} - 1$

(d)  $\frac{\pi}{2} + \ln 2$

(e)  $\frac{\pi}{4} - \ln 2$

**Exercise 5:** The volume of the solid generated by rotating the region bounded by the curves  $y = x$  and  $y = \sqrt{x}$  about the line  $x = -1$  is given by

(a)  $\int_0^1 \pi [(y + 1)^2 - (y^2 + 1)^2] dy$

(b)  $\int_0^1 \pi [(y - 1)^2 - (y^2 + 2)^2] dy$

(c)  $\int_0^1 \pi [y^2 - y^4] dy$

(d)  $\int_0^1 \pi [(y - 1)^2 - (y^2 - 1)^2] dy$

(e)  $\int_0^1 \pi [y^4 - y^2 + 2y - 1] dy$

**Exercise 6:** The region bounded by the curve  $y = 2\sqrt{x}$ , the  $x$ -axis, and the line  $x = 4$  is rotated about the  $y$ -axis. The volume of the solid generated is equal to

(a)  $\frac{2^8}{5}\pi$

(b)  $\frac{2^6}{3}\pi$

(c)  $\frac{2^8}{3}\pi$

(d)  $\frac{2^6}{5}\pi$

(e)  $\frac{2^6}{4}\pi$

**Exercise 7:** The sum of the series

$$\frac{(\ln 3)^2}{2!} + \frac{(\ln 3)^3}{3!} + \frac{(\ln 3)^4}{4!} + \dots$$

is equal to

- (a)  $2 - \ln 3$
- (b)  $3$
- (c)  $\ln 3$
- (d)  $1 + \ln 3$
- (e)  $2$

**Exercise 8:** The length of the curve

$$y = \int_0^x \sqrt{\sec^4 t - 1} dt, \quad 0 \leq x \leq \frac{\pi}{4}$$

is equal to

- (a)  $1$
- (b)  $\sqrt{2}$
- (c)  $3 - \sqrt{2}$
- (d)  $2$
- (e)  $1 + \sqrt{3}$

**Exercise 9:** The area of the surface obtained by rotating the curve  $y = \cosh x$ ,  $0 \leq x \leq 1$  about the  $y$ -axis is given by

(a)  $\int_0^1 2\pi x \cosh x dx$

(b)  $\int_0^1 2\pi x \sinh x dx$

(c)  $\int_0^1 2\pi \cosh x \sinh x dx$

(d)  $\int_0^1 2\pi x \cosh x \sinh x dx$

(e)  $\int_0^1 2\pi x \operatorname{sech} x dx$

**Exercise 10:** The integral

$$\int \frac{dx}{2\sqrt{x} + 2x}$$

is equal to

(a)  $\ln(1 + \sqrt{x}) + C$

(b)  $\ln(\sqrt{x}) + C$

(c)  $2 \ln(1 + \sqrt{x}) + C$

(d)  $\frac{1}{2} \ln(1 + \sqrt{x}) + C$

(e)  $\ln(x + \sqrt{x}) + C$

**Exercise 11:** The integral

$$\int_0^1 e^t \cosh t dt$$

is equal to

- (a)  $\frac{1}{4}(e^2 + 1)$
- (b)  $\frac{1}{2}(e^2 + 1)$
- (c)  $e^2 - 1$
- (d)  $\frac{1}{2}(e - 2)$
- (e)  $\frac{1}{3}(e + 2)$

**Exercise 12:** The integral

$$\int_0^{\frac{\pi}{2}} e^{\cos x} \sin(2x) dx$$

is equal to

- (a) 2
- (b)  $-\frac{1}{2}$
- (c) 0
- (d) 3
- (e) -4

**Exercise 13:** The integral

$$\int \frac{\sin^3 x}{\cos^4 x} dx$$

is equal to

- (a)  $\frac{1}{3} \sec^3 x - \sec x + C$
- (b)  $3 \sec x - \frac{1}{3} \sec^3 x + C$
- (c)  $\sec^3 x + \sec x + C$
- (d)  $\sec^2 x + \frac{1}{3} \sec^3 x + C$
- (e)  $-\sec^3 x + \frac{1}{3} \sec x + C$

**Exercise 14:** If  $x > 2$ , the integral

$$\int \frac{\sqrt{x^2 - 4}}{x} dx$$

is equal to

- (a)  $\sqrt{x^2 - 4} - 2 \sec^{-1} \left( \frac{x}{2} \right) + C$
- (b)  $\frac{2\sqrt{x^2-4}}{x} - 2 \sec^{-1} \left( \frac{x}{2} \right) + C$
- (c)  $2\sqrt{x^2 - 4} + 2 \tan^{-1} \left( \frac{x}{2} \right) + C$
- (d)  $x - 2 \sec^{-1} \left( \frac{x}{2} \right) + C$
- (e)  $2x\sqrt{x^2 - 4} - 2 \sec^{-1} \left( \frac{x}{2} \right) + C$



**Exercise 15:** If

$$\frac{3x^2 + 2x + 1}{(x - 1)(x^2 + 2x + 5)} = \frac{A}{x - 1} + \frac{Bx + C}{x^2 + 2x + 5}$$

then  $A + B + C$  is equal to

- (a)  $\frac{23}{4}$
- (b)  $\frac{12}{5}$
- (c)  $\frac{21}{10}$
- (d)  $\frac{4}{5}$
- (e) 0

**Exercise 16:** The integral

$$\int \frac{x^2 + 2x - 1}{2x^3 + 3x^2 - 2x} dx$$

is equal to

- (a)  $\frac{1}{2} \ln |x| + \frac{1}{10} \ln |2x - 1| - \frac{1}{10} \ln |x + 2| + C$
- (b)  $\ln |x| + \ln |2x - 1| - \ln |x + 2| + C$
- (c)  $\frac{1}{2} \ln |x| - \ln |x + 2| + C$
- (d)  $\frac{1}{2} \ln |x| + \frac{1}{5} \ln |2x - 1| + 3 \ln |x + 2| + C$
- (e)  $2 \ln |x| + 3 \ln |2x - 1| - 3 \ln |x + 2| + C$

**Exercise 17:** The integral

$$\int_2^{\infty} \frac{dx}{x(\ln x)^p}$$

converges for

- (a)  $p > 1$
- (b)  $p = 0$
- (c)  $p = -1$
- (d)  $p = 1$
- (e)  $p < 1$

**Exercise 18:** The sequence

$$\left\{ \left( 1 - \frac{2}{5n} \right)^{5n} \right\}_{n \geq 1}$$

is

- (a) convergent and its limit is  $e^{-2}$
- (b) convergent and its limit is  $e^{-5}$
- (c) convergent and its limit is  $e^{-2/5}$
- (d) convergent and its limit is  $e^3$
- (e) divergent

**Exercise 19:** The series

$$\sum_{n \geq 1} \frac{\sin(n\pi) + 2^n}{3^n}$$

is equal to

- (a) 2
- (b) 3
- (c) 33
- (d)  $\frac{2}{33}$
- (e) 6

**Exercise 20:** Which of the following proposition is True about the series

$$\sum_{n \geq 1} \frac{n}{n^2 + 1}$$

- (a) Diverges by the integral test
- (b) Converges by the integral test
- (c) Converges by the  $n^{\text{th}}$  term test
- (d) Diverges by the  $n^{\text{th}}$  term test
- (e) The integral test cannot be applied

**Exercise 21:** The series

$$\sum_{n \geq 1} \frac{4}{n(n+2)}$$

is equal to

- (a) 3
- (b) 4
- (c) 5
- (d) 7
- (e) 1024

**Exercise 22:** The series

$$\sum_{n \geq 1} \left( \frac{3n}{4n+1} \right)^n$$

is

- (a) Convergent by the root test
- (b) Divergent by the root test
- (c) A series for which the root test is inconclusive
- (d) Divergent by the  $n^{\text{th}}$ -term test of Divergence
- (e) Divergent by the limit comparison test

**Exercise 23:** The series

$$\sum_{n \geq 1}^n \frac{3^{n+2}}{\ln n}$$

is

- (a) Divergent by the ratio test
- (b) Convergent by the ratio test
- (c) A series for which the ratio test is inconclusive
- (d) Convergent by the  $n^{\text{th}}$  root test
- (e) The ratio test cannot be applied

**Exercise 24:** Which of the following proposition is False about the series

$$\sum_{n \geq 1} (-1)^{n+1} \frac{n}{n^3 + 1}$$

- (a) Diverges
- (b) Converges absolutely
- (c) Converges
- (d) Converges by the alternating series test
- (e) converges with the absolute convergence test and the alternating series test

**Exercise 25:** Which of the following proposition is True about the series

$$\sum_{n \geq 1} (-1)^{n+1} \frac{n!}{2^n}$$

- (a) Diverges
- (b) Converges absolutely
- (c) Converges by the ratio test
- (d) Converges by the integral test
- (e) converges absolutely by the ratio test

**Exercise 26:** The interval of convergence of the power series

$$\sum_{n \geq 1} \left(1 + \frac{1}{n}\right)^n (x - 1)^n$$

is

- (a)  $(0, 2)$
- (b)  $[0, 2]$
- (c)  $\left(\frac{e-1}{e}, \frac{e+1}{e}\right)$
- (d)  $[-1, 3)$
- (e)  $[0, 2)$

**Exercise 27:** The interval of convergence of the power series

$$\sum_{n \geq 1} \frac{5^n}{n} (x + 1)^n$$

is

- (a)  $[-\frac{6}{5}, -\frac{4}{5})$
- (b)  $(-\frac{6}{5}, -\frac{4}{5})$
- (c)  $[-\frac{6}{5}, -\frac{4}{5}]$
- (d)  $(-2, 1)$
- (e)  $[-2, 1)$

**Exercise 28:** The Taylor series of  $e^{2x} \sin x$  is

- (a)  $x + 2x^2 + \frac{11}{6}x^3 + \dots$
- (b)  $x - 3x^2 + \frac{1}{6}x^3 + \dots$
- (c)  $x + 2x^2 + \frac{11}{5}x^3 + \dots$
- (d)  $x - 3x^2 + \frac{11}{6}x^3 + \dots$
- (e)  $x + 2x^2 - \frac{11}{5}x^3 + \dots$