

King Fahd University of Petroleum and Minerals
Department of Mathematics and Statistics

Math 102
Final Exam
Term 152
Tuesday 17/05/2016

EXAM COVER

Number of versions: 4
Number of questions: 28
Number of Answers: 5 per question

This exam was prepared using mcqs
For questions send an email to Dr. Ibrahim Al-Lehyani (iallehyani@kaau.edu.sa)

King Fahd University of Petroleum and Minerals
Department of Mathematics and Statistics

Math 102
Final Exam
Term 152
Tuesday 17/05/2016
Net Time Allowed: 180 minutes

MASTER VERSION

1. The definite integral $\int_{-1}^5 3x + 2 dx$ is equal to

(a) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{6}{n} \left(\frac{18i}{n} - 1 \right)$

(b) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{6}{n} \left(\frac{6i}{n} + 1 \right)$

(c) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left(\frac{18i}{n} - 4 \right)$

(d) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left(\frac{6i}{n} - 3 \right)$

(e) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{6}{n} \left(\frac{18i}{n} + 5 \right)$

2. Let A be the exact area below a curve over the interval $[0, 1]$ and let B be an estimation of the same area using left endpoints of 10 subintervals. For which one of the following curves B is greater than A ?

(a) $\cos x$

(b) $\sin x$

(c) e^x

(d) x

(e) \sqrt{x}

3. $\int \sin^3 x \cos^2 x dx =$

(a) $\frac{1}{5} \cos^5 x - \frac{1}{3} \cos^3 x + C$

(b) $\frac{1}{4} \cos^4 x - \frac{1}{3} \sin^3 x + C$

(c) $\frac{1}{4} \cos^4 x - \frac{1}{6} \cos^6 x + C$

(d) $\frac{1}{3} \cos^3 x + \frac{1}{2} \sin^2 x + C$

(e) $-\cos^3 x \sin^2 x + C$

4. $\int_0^{\pi/4} \tan x \ln(\cos x) dx =$

(a) $-\frac{1}{8}(\ln 2)^2$

(b) $\sqrt{2} \ln 2$

(c) $2\sqrt{2}$

(d) -1

(e) 0

5. If g is a continuous function so that

$$\int_{\pi}^{2x} \cos\left(\frac{t}{2}\right) g(t) dt = \frac{x}{2} \sin x - \frac{\pi}{4}, \text{ then } g(2\pi) =$$

- (a) $\frac{\pi}{4}$
 - (b) $\frac{\pi}{2}$
 - (c) $-\pi$
 - (d) $\frac{1 + \pi}{2}$
 - (e) $\frac{-1 + \pi}{4}$
6. The area enclosed by the curves $x = 2y^2$ and $x = 4 + y^2$ is equal to

- (a) $\frac{32}{3}$
- (b) $\frac{31}{3}$
- (c) $\frac{29}{3}$
- (d) $\frac{28}{3}$
- (e) $\frac{26}{3}$

7. The improper integral $\int_1^{\infty} \frac{e^{1/x}}{x^2} dx$

- (a) converges to $e - 1$.
- (b) converges to $1 - \frac{1}{e}$.
- (c) converges to e .
- (d) converges to 1.
- (e) diverges.

8. The area of the surface obtained by revolving the curve $y = \ln(\sec x)$, $0 \leq x \leq \pi/3$ about the y -axis is

- (a) $2\pi \int_0^{\pi/3} x \sec x dx$
- (b) $2\pi \int_0^{\pi/3} \ln(\sec x) \sec x dx$
- (c) $2\pi \int_0^{\pi/3} \sec x dx$
- (d) $2\pi \int_0^{\pi/3} x \tan x dx$
- (e) $2\pi \int_0^{\pi/3} \ln(\sec x) \tan x dx$

9. If the velocity of a moving particle is $v(t) = t^2 + 5t - 6$ in m/s , then the total distance travelled by the particle during the time interval $0 \leq t \leq 4$ is

(a) $\int_1^4 t^2 + 5t - 6 dt - \int_0^1 t^2 + 5t - 6 dt$

(b) $\int_0^1 t^2 + 5t - 6 dt - \int_1^4 t^2 + 5t - 6 dt$

(c) $\int_1^4 t^2 + 5t - 6 dt$

(d) $\int_0^4 t^2 + 5t - 6 dt$

(e) $\int_0^1 t^2 + 5t - 6 dt$

10. The volume generated by rotating the region bounded by $y = \ln x$, $x = e$, and $y = 0$ about the y -axis is

(a) $\frac{\pi}{2}(e^2 + 1)$

(b) $\frac{\pi}{18}(2e^3 + 1)$

(c) $\frac{\pi}{2}(2e^2 - 1)$

(d) $\frac{\pi}{18}(e^3 - 1)$

(e) $\pi(2e^2 + 1)$

11. $\int_0^{\pi/2} \sinh x \sin x \, dx =$

(a) $\frac{1}{2} \cosh\left(\frac{\pi}{2}\right)$

(b) $2 \sinh\left(\frac{\pi}{2}\right)$

(c) $\frac{1}{2} \left[\cosh\left(\frac{\pi}{2}\right) - \sinh\left(\frac{\pi}{2}\right) \right]$

(d) $2 \left[\cosh\left(\frac{\pi}{2}\right) + \sinh\left(\frac{\pi}{2}\right) \right]$

(e) 0

12. $\int_1^{\sqrt{2}} \frac{\sqrt{x^2 - 1}}{x^2} \, dx =$

(a) $\ln(\sqrt{2} + 1) - \frac{\sqrt{2}}{2}$

(b) $\ln(\sqrt{3} + 2) - \frac{\sqrt{3}}{2}$

(c) $\ln(\sqrt{2}) + \frac{\sqrt{2}}{2}$

(d) $\ln(\sqrt{3}) + \frac{\sqrt{3}}{2}$

(e) $\ln(\sqrt{2} + 2) - \frac{\sqrt{2}}{2}$

13. $\int \frac{x+2}{x^2+4} dx =$

(a) $\ln \sqrt{x^2+4} + \tan^{-1} \left(\frac{x}{2} \right) + C$

(b) $\ln |x-2| + C$

(c) $\ln(x^2+4) + 2 \tan^{-1} x + C$

(d) $\ln \sqrt{x^2+4} + C$

(e) $\ln(x^2+4) + \frac{1}{2} \tan^{-1} \left(\frac{x}{2} \right) + C$

14. The length of the curve $y = \frac{1}{3} + \frac{4}{3}x^{3/2}$, $0 \leq x \leq 2$ is

(a) $\frac{13}{3}$

(b) $\frac{13}{2}$

(c) $\frac{52}{3}$

(d) $\frac{26}{3}$

(e) $\frac{15}{2}$

15. $\lim_{n \rightarrow \infty} \frac{1^3 + 2^3 + \dots + n^3}{n^4 - 5n} =$

- (a) $\frac{1}{4}$
- (b) $\frac{1}{5}$
- (c) 1
- (d) 0
- (e) does not exist

16. The series $\sum_{n=1}^{\infty} \frac{6}{9n^2 - 3n - 2}$ is

- (a) convergent and its sum is 2.
- (b) convergent and its sum is 1.
- (c) convergent and its sum is $2/3$.
- (d) convergent and its sum is 6.
- (e) divergent.

17. If $s_n = n \sin(1/n)$ is the sequence of partial sums of the series $\sum_{n=1}^{\infty} a_n$, then

(a) $\lim_{n \rightarrow \infty} a_n = 0$.

(b) the series $\sum_{n=1}^{\infty} a_n$ is divergent.

(c) the series $\sum_{n=1}^{\infty} a_n$ is convergent and its sum is 0.

(d) $\lim_{n \rightarrow \infty} s_n = 0$.

(e) $\lim_{n \rightarrow \infty} a_n$ does not exist.

18. The series $\sum_{n=1}^{\infty} \frac{n^3}{(n^2 + n)^q}$ is convergent for

(a) $q > 2$

(b) $q \geq 2$

(c) $q < 2$

(d) $q \leq 2$

(e) $q = 2$

19. Which one of the following statements is TRUE for the series $\sum_{n=1}^{\infty} \frac{\cos^2 n}{1+n^2}$, where $f(x) = \frac{\cos^2 x}{1+x^2}$?
- (a) The integral test is not applicable because f is not decreasing on $[1, \infty)$.
 - (b) The series converges by the integral test.
 - (c) The series diverges by the integral test.
 - (d) The integral test is not applicable because f is not positive on $[1, \infty)$.
 - (e) The integral test is not applicable because f is discontinuous on $[1, \infty)$.
20. Let $\sum_{n=1}^{\infty} (-1)^n b_n$, where $b_n > 0$, be an alternating series. If the sequence $\{b_n\}$ converges to a non zero number, then
- (a) the series diverges by the test for divergence.
 - (b) the series diverges by the alternating series test.
 - (c) the series conditionally converges.
 - (d) the series absolutely converges.
 - (e) the series diverges by the root test.

21. The series $\sum_{n=0}^{\infty} \frac{(n!)^2}{(2n+1)!}$

- (a) converges by the ratio test.
- (b) diverges by the alternating series test.
- (c) conditionally converges.
- (d) converges by the integral test.
- (e) diverges by the test for divergence.

22. The series $\sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi}{2}\right)}{n^{2/3}}$

- (a) converges conditionally.
- (b) diverges.
- (c) converges absolutely.
- (d) is not an alternating series.
- (e) diverges by the ratio test.

both choices are accepted
as correct answers.

23. The interval of convergence of the series $\sum_{n=1}^{\infty} \frac{3^n}{n} (2x - 1)^n$ is

(a) $\left[\frac{1}{3}, \frac{2}{3}\right)$

(b) $\left(\frac{1}{3}, \frac{2}{3}\right)$

(c) $\left(-\frac{1}{3}, \frac{1}{3}\right)$

(d) $\left(-\frac{1}{3}, \frac{1}{3}\right]$

(e) $(-\infty, \infty)$

24. For $|x| < 1$, a power series representation of $f(x) = x \tan^{-1} x$ is

(a) $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+2}}{2n+1}$

(b) $\sum_{n=0}^{\infty} \frac{x^{2n+2}}{2n+1}$

(c) $\sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1}$

(d) $\sum_{n=1}^{\infty} (-1)^n \frac{x^{2n+2}}{2n}$

(e) $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n+1}}{2n}$

25. The radius of convergence of the series $\sum_{n=1}^{\infty} \left(1 + \frac{1}{n}\right)^{-n^2} x^n$ is

(a) e

(b) $\frac{1}{e}$

(c) ∞

(d) e^2

(e) $\frac{1}{e^2}$

26. If $f(x) = \frac{1}{1+x}$ has a power series expansion at $x = 2$, then its Taylor series centered at $x = 2$ is

(a) $\sum_{n=0}^{\infty} \frac{(-1)^n}{3^{n+1}} (x-2)^n$

(b) $\sum_{n=0}^{\infty} (-1)^n (x-2)^n$

(c) $\sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} (x-2)^n$

(d) $\sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} (x-2)^n$

(e) $\sum_{n=0}^{\infty} \frac{(-1)^n}{3^n} (x-2)^n$

27. If $P(x)$ is the sum of the first three non zero terms of the Maclaurin series of $f(x) = (1+x)^{-1/2} \cos x$, then $P(1/2) =$ (Hint: You may use the product of the Maclaurin series of $\cos x$ and $(1+x)^{-1/2}$.)

(a) $\frac{23}{32}$

(b) $\frac{16}{17}$

(c) $\frac{25}{32}$

(d) $\frac{33}{34}$

(e) $\frac{9}{4}$

28. The sum of the series $\sum_{n=0}^{\infty} (-1)^n \frac{\pi^{2n+1}}{4^n (2n+1)!}$ is

(a) 2

(b) -2

(c) 1

(d) -1

(e) 0

Q	MM	V1	V2	V3	V4
1	a	a	d	b	a
2	a	e	e	d	a
3	a	d	b	a	b
4	a	e	b	a	d
5	a	b	e	c	c
6	a	b	c	b	b
7	a	d	d	e	d
8	a	b	b	e	c
9	a	b	d	a	a
10	a	b	b	e	e
11	a	b	a	d	c
12	a	d	d	b	a
13	a	d	c	e	a
14	a	e	d	c	a
15	a	d	a	a	e
16	a	a	a	c	b
17	a	c	a	a	b
18	a	c	a	a	a
19	a	c	c	c	b
20	a	c	a	a	d
21	a	e	c	e	c
22	a	d	c	b	e
23	a	d	b	e	d
24	a	d	a	e	d
25	a	e	b	a	d
26	a	a	a	c	a
27	a	a	b	a	b
28	a	b	c	b	e