

$$\bar{x} \pm z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \text{ or } \bar{x} \pm t_{\frac{\alpha}{2}, n-1} \frac{s}{\sqrt{n}}$$

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\frac{\alpha}{2}, df} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} \text{ where}$$

$$df = \frac{\left(s_1^2/n_1 + s_2^2/n_2\right)^2}{\frac{(s_1^2/n_1)^2}{n_1-1} + \frac{(s_2^2/n_2)^2}{n_2-1}} \text{ or}$$

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\frac{\alpha}{2}, n_1+n_2-2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \text{ where}$$

$$s_p = \sqrt{\frac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{n_1+n_2-2}}$$

## Chapter 9

$$z_0 = \frac{\bar{x} - \mu_0}{\frac{\sigma}{\sqrt{n}}} \text{ or } t_0 = \frac{\bar{x} - \mu_0}{\frac{s}{\sqrt{n}}}$$

$$z_0 = \frac{p - \pi_0}{\sqrt{\frac{\pi_0(1-\pi_0)}{n}}} \text{ or } z_0 = \frac{x - n\pi_0}{\sqrt{n\pi_0(1-\pi_0)}}$$

## Chapter 10

$$t_0 = \frac{\bar{D} - \mu_D}{\frac{s_D}{\sqrt{n}}} \text{ where } \bar{D} = \frac{\sum_i^n d_i}{n} \text{ and}$$

$$s_D = \sqrt{\frac{\sum_{i=1}^n (d_i - \bar{D})^2}{n-1}}$$

$$z_0 = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

$$t_0 = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$t_0 = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

If  $\bar{p} = \frac{n_1 p_1 + n_2 p_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$  then

$$z = \frac{(p_1 - p_2) - (\pi_1 - \pi_2)}{\sqrt{\bar{p}(1-\bar{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$$

$$F_{STAT} = \frac{s_1^2}{s_2^2} \text{ with}$$

$$df_1 = n_1 - 1 \text{ and } df_2 = n_2 - 1.$$

## Chapter 12

$$\chi_{stat}^2 = \sum_{j=1}^r \sum_{i=1}^c \frac{(O_{ij} - E_{ij})^2}{E_{ij}} \quad df = (c-1)(r-1)$$

$$|p_i - p_j| > \sqrt{\chi_{\alpha}^2} \sqrt{\frac{p_i(1-p_i)}{n_i} + \frac{p_j(1-p_j)}{n_j}}$$

$$Z_0 = \frac{B-C}{\sqrt{B+C}}$$