

EE 340


Electromagnetics

Lab

Problem Session #1


Part 1: Visualization of surfaces in 3D coordinate systems:

(a)




Part 1: Visualization of surfaces in 3D coordinate systems:

(b)




Part 1: Visualization of surfaces in 3D coordinate systems:

(c)




Part 1: Visualization of surfaces in 3D coordinate systems:

(d)




Part 1: Visualization of surfaces in 3D coordinate systems:

(e)




Part 1: Visualization of surfaces in 3D coordinate systems:

(f)




Part 1: Visualization of surfaces in 3D coordinate systems:

(g)




Part 2: Visualization of surfaces in 3D coordinate systems:

(a)




Part 2: Visualization of surfaces in 3D coordinate systems:

(b)




Part 2: Visualization of surfaces in 3D coordinate systems:

(c)



Part 2: Visualization of surfaces in 3D coordinate systems:

(d)



Part(3): Vector Algebra

Problem 1.5

For $\bar{U} = Ux\alpha x + 5Ay - Az$, $\bar{V} = 2Ax - VyAy + 3Az$ and $\bar{W} = 6Ax + Ay + WzAz$, obtain Ux , Vy and Wz such that U , V and W are mutually orthogonal ?

$$\begin{aligned} \bar{U}, \bar{V} &= (uXax + 5ay - az), (2ax - vy, ay + 3az) \\ &= 2Ux - 5Vy - 3 = 0 \quad \dots \dots \dots (1) \\ \bar{U}, \bar{W} &= (Ux, Ax + 5ay - az), (6ax + ay + Wz, az) \\ &= -6Ux + 5Wz = 0 \quad \dots \dots \dots (2) \\ \bar{V}, \bar{W} &= (2ax - Vy, ay + 3az), (6ax + ay + Wz, az) \\ &= 12 - Vy + 3Wz = 0 \quad \dots \dots \dots (3) \end{aligned} \quad \left. \begin{array}{l} \text{eq.1} * (-3) + \text{eq.2} \Rightarrow 15Vy - 6Wz + 15 = 0 \\ \text{eq.3} * (5) + \text{eq.4} \Rightarrow Wz = -\frac{25}{22} \end{array} \right\} \begin{array}{l} -4 \\ -22 \end{array}$$

Problem 1.10

$$(a) \quad \tilde{A} \cdot (\tilde{A} \tilde{X} \tilde{B}) = 0 = \tilde{B} \cdot (\tilde{A} \tilde{X} \tilde{B})$$

$$(AX\vec{B}) = \begin{vmatrix} ax & ay & az \\ Ax & Ay & Az \\ Bx & By & Bz \end{vmatrix}$$

$$-ax(AyBz - AzBy) - ay(AxBz - AzBx) + az(AxBy - AyBx)$$

$$A \cdot (AXB) = AxAyBz - AxAzBy - AyAxBz + AyAzBxAzAxBy - AyAzBx = 0$$

$$B. \quad (AXB) = AyBxBz - AzBxBz - AxByBz + AzBxBz + AxByBz - AyBxRz = 0$$

So this statement is true.

Remember (det phras.)

$$\alpha_x \cdot \alpha_y = \alpha_y \cdot \alpha_x = \alpha_2 \cdot \alpha_2 = 1$$

$$a_x \cdot a_y = a_y \cdot a_z = a_z \cdot a_x = 0$$

Part 3: Vector Algebra

Problem 1.5

$$(b) (\overline{A} \cdot \overline{B})^2 + |\overline{AXB}|^2 = (AB)^2$$

$|\overline{AXB}|^2 - (AB)^2 \sin^2 \theta \quad \xrightarrow{\text{direction ignored.}}$

$$(A \cdot B)^2 + |\overline{AXB}|^2 = (AB \cos \theta)^2 + (AB \sin \theta)^2 = (AB)^2 \cos^2 \theta + (AB)^2 \sin^2 \theta$$

$$= (AB)^2 (\cos^2 \theta + \sin^2 \theta) = (AB)^2 * I = RHS$$

$$(c) \quad \bar{A} = Ax \bar{ax} + Ay \bar{ay} + Az \bar{az}$$

Because $A \cdot Ax \in Ax$

$$A \cdot Av \in Av$$

$$A \cap A_7 \in A_7$$

[View Details](#)

7 cube Ax, Ay or Az loads;

$$\begin{array}{l} A \cdot Ay \in Ay \\ A \cdot Az \in Az \end{array} \quad \left\{ \begin{array}{l} \text{Satz 1.1.1} \\ \text{Satz 1.1.2} \end{array} \right.$$

$$= (\bar{z} \bar{z}) = -(\bar{z} \bar{z}) \in \langle \sqrt{5} \rangle$$

$$\bar{A} = (A \cdot \alpha_x) \alpha_x + (A \cdot \alpha_y) \alpha_y + (A \cdot \alpha_z) \alpha_z$$

Part 4: Coordinate Transformations

Problem 2.1 convert the following points to Cartesian coordinates:

$$X = \rho \cos\theta \quad Y = \rho \sin\theta \quad Z = Z$$

(a) $P(5, 120^\circ, 0)$

$$X = 5 \cos 120^\circ = -2.5 \quad Y = 5 \sin 120^\circ = 4.33 \quad Z = 0$$

$$P(-2.5, 4.33, 0)$$

(b) $P(1, 30^\circ, -10)$

$$X = 1 \cos 30^\circ = 0.866 \quad Y = 1 \sin 30^\circ = 0.5 \quad Z = -10$$

$$P(0.866, 0.5, -10)$$

(c) $P(10, 3\pi/4, \pi/2)$

$$X = r \sin\theta \cos\phi \quad Y = r \sin\theta \sin\phi \quad Z = r \cos\theta$$

$$X = 10 \sin 3\pi/4 \cos \pi/2 = 0$$

$$Y = 10 \sin 3\pi/4 \sin \pi/2 = 7.071$$

$$Z = 10 \cos 3\pi/4 = -7.071$$

$$P(0, 7.071, -7.071)$$

Part 4: Coordinate Transformations

Problem 2.2 Express in Cylindrical and spherical coordinates

(a) $P(1, -4, -3)$

$$\rho = \sqrt{1+16} = \sqrt{17}, \quad \phi = \tan^{-1}(-4/1) = -75.94^\circ = 284.06^\circ, \quad \& z = -3$$

$$P(\sqrt{17}, -75.94^\circ, -3) \xrightarrow{\text{Cylindrical}}$$

$$r = \sqrt{1+16+9} = \sqrt{26} \quad \text{as } r = \sqrt{x^2 + y^2 + z^2}$$

$$\theta = \tan^{-1}(\sqrt{17}/-3) = -53.96^\circ$$

$$P(\sqrt{26}, -53.96^\circ, -75.94^\circ) \xrightarrow{\text{Spherical}}$$

(b) $Q(3, 0, 5)$

$$\rho = \sqrt{9+0} = 3, \quad \phi = \tan^{-1}(0/3) = 0^\circ \quad \& z = 5$$

$$Q(3, 0, 5) \xrightarrow{\text{Cylindrical}}$$

$$Q(\sqrt{34}, 30.96^\circ, 0) \xrightarrow{\text{Spherical}}$$

(c) $R(-2, 6, 0)$

$$\rho = \sqrt{4+36} = \sqrt{40}, \quad \phi = \tan^{-1}(6/-2) = -71.565^\circ \quad \& z = 0$$

$$P(\sqrt{40}, -71.565^\circ, 0)$$

Part 4: Coordinate Transformations

2.3(a)
$$\begin{pmatrix} P_r \\ P_\theta \\ P_z \end{pmatrix} = \begin{pmatrix} \cos\phi & \sin\phi & 0 \\ -\sin\phi & \cos\phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y+z \\ 0 \\ 0 \end{pmatrix} \quad (\text{Eq. 2.13})$$

Cylindrical

$$P_r = \cos\phi (y+z) = \cos\phi (\rho \sin\phi + z)$$

$$P_\theta = -\sin\phi (y+z) = -\sin\phi (\rho \sin\phi + z)$$

$$P_z = 0$$

$$\rightarrow \vec{P} = (\rho \sin\phi + z) (\cos\phi \vec{a}_r - \sin\phi \vec{a}_\theta)$$

Spherical

$$P_r = (y+z) (\sin\phi \cos\theta) = (r \sin\theta \sin\phi + r \cos\theta) (\sin\phi \cos\theta)$$

$$P_\theta = (y+z) (\cos\theta \cos\phi) \Rightarrow$$

$$P_\phi = (y+z) (-\sin\phi) \Rightarrow$$

$$\rightarrow \vec{P} = r (\sin\theta \sin\phi + \cos\theta) (\sin\phi \cos\phi \vec{a}_r + \cos\theta \cos\phi \vec{a}_\theta - \sin\phi \vec{a}_\phi)$$

Part 4: Coordinate Transformations

2.3(b) Cylindrical

$$\vec{Q} = (y \cos\phi + xz \sin\phi) \vec{a}_r + (-y \sin\phi + xz \cos\phi) \vec{a}_\theta + (x+y) \vec{a}_z$$

$$\text{sub. } x = \rho \cos\phi \quad y = \rho \sin\phi$$

$$\Rightarrow \vec{Q} = \frac{1}{2} \rho \sin 2\phi (1+z) \vec{a}_r + \rho (z \cos^2\phi - \sin^2\phi) \vec{a}_\theta + \rho (\cos\phi + \sin\phi)$$

Spherical

$$\vec{Q} = \left[\frac{1}{2} r \sin 2\phi \sin^2\theta + \frac{1}{2} r^2 \sin^2\theta \cos\theta \sin^2\phi + \frac{1}{2} r (\cos\phi + \sin\phi) \cos 2\theta \right]$$

$$+ \left[\frac{1}{4} r \sin 2\theta \sin 2\phi + \frac{1}{2} r^2 \cos^2\theta \sin\theta \sin 2\phi - \frac{r}{2} (\cos\phi + \sin\phi) \sin^2\theta \right]$$

$$+ \left(-r \sin\theta \sin^2\phi + \frac{1}{2} r^2 \sin 2\theta \cos^2\phi \right) \vec{a}_\phi$$

Part 4: Coordinate Transformations (Cylindrical case)

$$\frac{2.3}{(c)} T = \left[\frac{x^2}{x^2 + y^2} - y^2 \right] ax + \left[\frac{xy}{x^2 + y^2} + xy \right] ay + az$$

$$\begin{bmatrix} A\rho \\ A\theta \\ Az \end{bmatrix} = \begin{bmatrix} \cos\theta & \sin\theta & 0 \\ -\sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} ax \\ ay \\ az \end{bmatrix}$$

$$\begin{bmatrix} Ar \\ A\theta \\ A\phi \end{bmatrix} = \begin{bmatrix} \sin\theta\cos\phi & \sin\theta\sin\phi & \cos\theta \\ \cos\theta\cos\phi & \cos\theta\sin\phi & -\sin\theta \\ -\sin\phi & \cos\phi & 0 \end{bmatrix} \begin{bmatrix} Ax \\ Ay \\ Az \end{bmatrix}$$

$$\boxed{A\rho} = \cos\phi \left[\frac{x^2}{x^2 + y^2} - y^2 \right] + \sin\phi \left[\frac{xy}{x^2 + y^2} + xy \right] + 0 \\ = \cos^3\phi - \rho^2 \sin^2\phi \cos\phi + \sin^2\phi \cos\phi + \rho^2 \sin^2\phi \cos\phi = \cos\phi$$

Remember: $x = \rho \cos\phi$; $y = \rho \sin\phi$; $z = z$

$$\boxed{A\phi} = \left[\frac{x^2}{x^2 + y^2} - y^2 \right] (-\sin\phi) + \left[\frac{xy}{x^2 + y^2} + xy \right] (\cos\phi) + 0 \\ = -\sin\phi \cos^2\phi + \rho^2 \sin^3\phi + \sin\phi \cos^2\phi + \rho^2 \sin\phi \cos^2\phi = \rho^2 \sin\phi$$

$$\boxed{Az=1} \quad \text{So, } \boxed{T = \cos\phi \bar{a}_\rho + \rho^2 \sin\phi \bar{a}_\phi + \bar{a}_z}$$

Part 4: Coordinate Transformations (Spherical case)

$$Ar = \left[\frac{x^2}{x^2 + y^2} - y^2 \right] \sin\theta \cos\phi + \left[\frac{xy}{x^2 + y^2} + xy \right] \sin\theta \sin\phi + \cos\theta \\ = \sin\theta \cos^3\phi - r^2 \sin^3\theta \sin^2\phi \cos\phi + \cos\phi \sin^2\phi \sin\theta + r^2 \sin^3\theta \cos\phi \sin^2\phi \\ + \cos\theta \\ = \sin\theta \cos\phi (\cos^2\phi + \sin^2\phi + r^2 \sin^2\theta \sin^2\phi - r^2 \sin^2\phi \sin^2\theta) + \cos\theta \\ = \sin\theta \cos\phi + \cos\theta$$

Remember: $x = r \sin\theta \cos\phi$

$$AO = \left[\frac{x^2}{x^2 + y^2} - y^2 \right] \cos\theta \cos\phi + \left[\frac{xy}{x^2 + y^2} + xy \right] \cos\theta \sin\phi - \sin\theta \\ = (\cos^2\phi - r^2 \sin^2\theta \cos^2\phi) (\cos\theta \cos\phi) + (\cos\phi \sin\phi + r^2 \sin^2\theta \cos\phi \sin\phi) (\cos\theta \sin\phi) + \cos\theta \\ = r^2 \sin^2\theta \cos\phi \cos\theta (\cos^2\phi + \sin^2\phi) + \cos\theta \cos\phi (\sin^2\phi - \cos^2\phi) + \cos\theta \\ = \cos\theta \cos^3\phi - r^2 \sin^2\theta \sin^2\phi \cos\theta \cos\phi + \cos\theta \sin^2\phi \cos\phi + r^2 \sin^2\theta \cos\phi \sin^2\phi \cos\theta - \sin\theta \\ = \cos\theta \cos\phi \sin\theta$$

Part 4: Coordinate Transformations (Spherical case)

$$\begin{aligned} A\phi &= \left[\frac{x^2}{x^2 + y^2} - y^2 \right] (-\sin\phi) + \left[\frac{xy}{x^2 + y^2} + xy \right] \cos\phi + 0 \\ &= (\cos^2\phi - r^2 \sin^2\theta \cos^2\phi)(-\sin\phi) + (\cos\phi \sin\phi + r^2 \sin^2\theta \cos\phi \sin\phi)\cos\phi + 0 \\ &= -\sin\phi \cos^2\theta + r^2 \sin^2\theta \sin^3\phi + \cos^2\theta \sin\phi + r^2 \sin^2\theta \cos^2\theta \sin\phi \\ &= r^2 \sin^2\theta \sin\phi \end{aligned}$$

$$T = (\sin\theta \cos\phi + \cos\theta) \cancel{\hat{a}_r} + (\cos\theta \cos\phi - \sin\theta) \cancel{\hat{a}_\theta} + r^2 \sin^2\theta \sin\phi \cancel{\hat{a}_\phi}$$