


Q1.

A sinusoidal travelling wave in a string is given by equation $y = y_m \sin(kx + \omega t)$ and its snapshot at an instant is shown in **FIGURE 1**. Three string elements are indicated by the lettered points. Which of the following is correct about the direction of motion of the string elements **a**, **b**, and **c** at the instant of the snapshot.

Fig#



- A) a-up, b-down, c-up
- B) a-up, b-up, c-down
- C) a-down, b-down, c-up
- D) a-down, b-up, c-down
- E) a-up, b-down, c-down

Q2.

Two sinusoidal waves, identical except for phase, travel in the same direction along a string are given by

$$y_1 = 0.025 \sin(15.0x - 90.0t)$$

$$y_2 = 0.025 \sin(15.0x - 90.0t + \pi/3)$$


Where x, y are in m and t is in s. At what average rate does the resultant wave transport the energy? ($\mu = 500 \text{ g/m}$ for the string)

- A) 22.8 W
- B) 17.5 W
- C) 12.3 W
- D) 7.58 W
- E) 9.37 W

Q3.

A sinusoidal wave travels along a string under tension. **FIGURE 2** gives the slopes (of string elements) along the string at time $t = 0$. The scale of the x -axis is set by $x_s = 0.80 \text{ m}$. what is the amplitude of the wave?

Fig#



- A) 1.3 cm
- B) 0.20 cm
- C) 0.40 cm
- D) 2.1 cm
- E) 1.8 cm

Q4.

A rope, under a tension of 200 N and fixed at both ends, oscillates in a second-harmonic standing wave pattern. The displacement of the rope is given by:

$$y = (0.10 \text{ m}) (\sin \pi x/2) (\cos 12\pi t)$$

where $x = 0$ at one end of the rope, x is in meters, and t is in seconds. What is the speed of the waves on the rope?

- A) 24 m/s
- B) 12 m/s
- C) 6.0 m/s
- D) 36 m/s
- E) 18 m/s

Q5.

In an experiment on standing waves, a string 90 cm long is attached to an oscillator that oscillates at a frequency of 80 Hz. The mass of the string is 0.044 kg. Find the tension in the string if it is oscillating in four loops.

- A) 63 N
- B) 1.8 N
- C) 34 N
- D) 27 N
- E) 51 N

Q6.


A man strikes one end of a rod with a hammer. The speed of sound in the rod is 15.0 times the speed of sound in air. A man, at the other end with his ear close to the rod, hears the sound of the blow twice with a 0.120 s interval between them; one sound comes through the rod and the other comes through the air along-side of the rod. If the speed of sound in air is 343 m/s, what is the length of the rod?

- A) 44.1 m
 - B) 617 m
 - C) 41.2 m
 - D) 246 m
 - E) 17.3 m
-

Q7.

Two sound sources S_1 and S_2 , shown in **FIGURE 3** are driven by the same oscillator whose frequency is 686 Hz. They are located at distance of 4.0 m on a vertical line. If you slide a microphone from point P to S_2 along the horizontal line PS_2 , how many time(s) the microphone will detect minimum in sound intensity along this line. (speed of sound 343 m/s)

Fig#



- A) Four
- B) One
- C) Three
- D) None
- E) Two

Q8.


Suppose that the sound level of a sound is initially at 70.0 dB and then drops to 50.0 dB. Assuming that the frequency of the sound is 600 Hz, determine the ratio between initial sound wave amplitude S_{m1} to final sound wave amplitude S_{m2} (i.e. S_{m1}/S_{m2}).

- A) 10.0
- B) 100
- C) 1.40
- D) 1.96
- E) 0.0150

Q9.

FIGURE 4 shows a moving sound source S that emits at a certain frequency, and four stationary sound detectors (located at points 1, 2, 3, and 4). Rank the detectors according to the frequency of the sound they detect from the source, **GREATEST FIRST**. (The source is moving towards the detector **1**).

Fig#



- A) 1, 4, 3, 2
- B) 1 and 2 tie, 3, 4
- C) 3, 4, then 1 and 2 tie
- D) 3, 4, 1, 2
- E) 4, 3, 2, 1

Q10.

Suppose that on a linear temperature scale X, water boils at -53.5°X and freezes at -170°X . What is a temperature of 340 K on the X scale? ($0^\circ\text{C} = 273\text{ K}$)

- A) -91.9°X
- B) -211°X
- C) -58.7°X
- D) $+50.2^\circ\text{X}$
- E) $+83.7^\circ\text{X}$

Q11.

What is the change in density of an aluminum cube of mass 200 g and of edge length 5.0 cm when heated from 10°C to 80°C (coefficient of linear expansion of aluminum $23 \times 10^{-6}/^\circ\text{C}$).

- A) $7.7 \times 10^{-3}\text{ g/cm}^3$
- B) $5.0 \times 10^{-1}\text{ g/cm}^3$
- C) 3.3 g/cm^3
- D) $2.4 \times 10^{-3}\text{ g/cm}^3$
- E) $9.7 \times 10^{-2}\text{ g/cm}^3$

Q12.


Materials A, B, and C are solids that are at their melting temperatures. Material A requires 200 J to melt 4 kg, material B requires 300 J to melt 5 kg, and material C requires 300 J to melt 6 kg. Rank the materials according to their heats of fusion, **GREATEST FIRST**.

- A) B, then A and C tie
- B) B and C tie, A
- C) C, B, A
- D) A, B, C
- E) A, B and C all tie

Q13.

FIGURE 5 displays a closed cycle for a gas. The change in internal energy along path **ca** is -160 J . The energy transferred to the gas as heat is 200 J along **ab**, and 40.0 J along path **bc**. How much work is done by the gas along path **abc**?

Fig#



- A) 80.0 J
- B) 400 J
- C) 0.00 J
- D) 200 J
- E) 40.0 J

Q14.

Suppose 1.80 mol of an ideal gas is taken from a volume of 3.00 cm^3 to a volume of 1.50 cm^3 via an isothermal compression at $30.0 \text{ }^\circ\text{C}$. How much energy is transferred as heat during the compression.

- A) $-3.14 \times 10^3 \text{ J}$
- B) $+3.14 \times 10^3 \text{ J}$
- C) $+2.95 \times 10^2 \text{ J}$
- D) $-2.95 \times 10^2 \text{ J}$
- E) $+5.63 \times 10^3 \text{ J}$


Q15.

A cylinder contains a mixture of helium (He) and argon (Ar) gas in equilibrium at $150 \text{ }^\circ\text{C}$. Find the ratio of root-mean-square speed of helium to that of argon ($V_{\text{He}}:V_{\text{Ar}}$). (molar masses of helium and argon are 4.00 g/mole and 40.0 g/mole , respectively).

- A) 3.16
 - B) 0.172
 - C) 4.00
 - D) 0.250
 - E) 6.32
-

Q16.
Rank the four paths of **FIGURE 6** according to the change in the internal energy of the gas, **MOST POSITIVE FIRST** and most negative last.

Fig#




- A) 1, 2, then 3 and 4 tie
- B) 1 and 2 tie, then 3 and 4 tie
- C) 2, 1, 3, 4
- D) 1 and 4 tie, 2, 3
- E) 4, 3, 2, 1

Q17.

One mole of an ideal monoatomic gas passes through a cycle as shown in **FIGURE 7**. The temperatures $T_1 = 300.0$ K, $T_2 = 600.0$ K, and $T_3 = 455.0$ K. What is the work done for path $2 \rightarrow 3$?

Fig#



- A) +1807 J
- B) -1807 J
- C) +1353 J
- D) -1353 J
- E) +9752 J

Q18.


0.300 kg of water at 90.0 °C and 0.700 kg of water at 10.0 °C are mixed together in an isolated container and come to equilibrium. Determine the change in entropy of the system of 1.00 kg of water.

- A) +28.1 J/K
- B) +239 J/K
- C) -210 J/K
- D) -13.2 J/K
- E) +550 J/K

Q19.

Which one is/are **NOT** a Carnot heat engine in **FIGURE 8**?

Fig#



- A) B only
- B) C only
- C) A and B
- D) A and C
- E) B and C

Q20.

An ideal refrigerator utilizes a Carnot cycle operating between 0 °C and 25 °C. To turn 10 kg of liquid water at 0 °C into 10 kg of ice at 0 °C how much energy must be supplied to the refrigerator?

- A) 3.0×10^5 J
- B) 3.6×10^6 J
- C) 4.3×10^6 J
- D) 1.7×10^5 J
- E) 5.9×10^5 J