



Suggested problems: Chapter 19- HRW-Principles of Physics- ISV 10th Edition.

14. One mole of an ideal diatomic gas goes from a to c along the diagonal path in Fig. 19-18. The scale of the vertical axis is set by $p_{ab} = 5.0$ kPa and $p_c = 0.5$ kPa, and the scale of the horizontal axis is set by $V_{bc} = 4.0$ m³ and $V_a = 2.0$ m³. During the transition, (a) what is the change in internal energy of the gas, and (b) how much energy is added to the gas as heat? (c) How much heat is required if the gas goes from a to c along the indirect path abc?

**Fig. 19-18, Problem 14****Answer:**(a) -20 kJ; (b) -15 kJ; (c) -10 kJ

23. Figure 19-21 shows a cycle undergone by 2.00 mol of an ideal monatomic gas. The temperatures are $T_1 = 300$ K, $T_2 = 600$ K, and $T_3 = 455$ K. For $1 \rightarrow 2$, what are (a) heat Q , (b) the change in internal energy ΔE_{int} , and (c) the work done W ? For $2 \rightarrow 3$, what are (d) Q , (e) ΔE_{int} , and (f) W ? For $3 \rightarrow 1$, what are (g) Q , (h) ΔE_{int} , and (i) W ? For the full cycle, what are (j) Q , (k) ΔE_{int} , and (l) W ? The initial pressure at point 1 is 1.00 atm ($= 1.013 \times 10^5$ Pa). What are the (m) volume and (n) pressure at point 2 and the (o) volume and (p) pressure at point 3?

**Fig. 19-21 Problem 23**

Answer: (a) +3.74 kJ; (b) +3.74 kJ; (c) zero; (d) zero; (e) -1.81 kJ; (f) +1.81 kJ; (g) -3.22 kJ; (h) -1.93 kJ; (i) -1.29 kJ; (j) +0.520 kJ; (k) zero; (l) +0.520 kJ; (m) 0.0246 m³; (n) 2.026×10^5 Pa = 2.00 atm; (o) 0.0373 m³; (p) 1.013×10^5 Pa = 1.00 atm

29. The volume of an ideal gas is adiabatically reduced from 350 L to 130 L. The initial pressure and temperature are 2.00 atm and 380 K. The final pressure is 8.00 atm. (a) Is the gas monatomic, diatomic, or polyatomic? (b) What is the final temperature? (c) How many moles are in the gas?

Answer: (a) diatomic; (b) 565 K; (c) 22.4 mol.

30. Under constant pressure, the temperature of 3.00 mol of an ideal monatomic gas is raised by 15.0 K. What are (a) the work W done by the gas, (b) the energy transferred as heat Q , (c) the change ΔE_{int} in the internal energy of the gas, and (d) the change ΔK in the average kinetic energy per atom?

Answer:(a) +374 J; (b) +935 J; (c) +561 J; (d) $+3.11 \times 10^{-22}$ J

62. Compute (a) the number of moles and (b) the number of molecules in 1.00 cm³ of an ideal gas at a pressure of 75.0 Pa and a temperature of 285 K

Answer:(a) 3.17×10^{-8} mol; (b) 1.19×10^{16} molecules

63. (a) Compute the rms speed of a nitrogen molecule at 80.0 °C. The molar mass of nitrogen molecules (N_2) is given in Table 19-1. At what temperatures will the rms speed be (b) half that value and (c) twice that value?

Answer: (a) 561 m/s; (b) 88.25 K = -185 °C; (c) 1.412×10^3 K = 1.14×10^3 °C