
QUESTION NO: 1

0

A 0.3-kg mass, attached to the end of a string, swings in a vertical circle, as shown in Figure 1. At the instant when theta equals 50 degrees, the tension in the string is 8.0 N. What is the magnitude of the resultant force on the mass at this instant?

- A. 6.5 N
- B. 4.7 N
- C. 8.4 N
- D. 1.4 N
- E. 7.9 N

_*****

QUESTION NO: 2

0

In Figure 2, the coefficient of kinetic friction between m_1 and the table is 0.39. The mass $m_1=37$ kg and the mass $m_2=55$ kg. Assume the pulley is massless and frictionless. The magnitude of the acceleration is:

- A. 4.32 m/s^2 .
- B. 5.07 m/s^2 .
- C. 4.68 m/s^2 .
- D. 3.51 m/s^2 .
- E. 5.63 m/s^2 .

_*****

QUESTION NO: 3

0

A 1 kg mass is attached to a light string of length 2 m to form a simple pendulum. The mass is released from rest at $\theta = 45$ degrees. Find the tension in the string at the lowest point.

- A. 15.54 N
- B. 13.34 N
- C. 18.16 N
- D. 16.37 N
- E. 12.67 N

1 TEST CODE: PAGE: 002

_*****

QUESTION NO: 4

0

A force F , shown as a function of x in Figure 3, acts on a 2 kg mass. If the particle starts with an initial speed of 8 m/s, find the speed of the particle at $x=6$ m.

- A. 10.2 m/s.
- B. 9.2 m/s.
- C. 8.8 m/s.
- D. 6.2 m/s.
- E. 12.4 m/s.

QUESTION NO: 5

0

A 20 kg mass is fastened to a light spring ($k=300 \text{ N/m}$) that passes over a pulley as shown in Figure 4. The pulley is frictionless, and the mass is released from rest when the spring is unstretched. After the mass has moved downwards 0.40 m, the speed of the 20 kg mass is:

- A. 2.33 m/s.
- B. 1.82 m/s.
- C. 2.12 m/s.
- D. 4.11 m/s.
- E. 3.65 m/s.

QUESTION NO: 6

0

A 3 kg object moving with 8 m/s in the positive x direction has a one dimensional elastic collision with an object of mass M initially at rest. After the collision the object of the unknown mass M has a velocity of 6 m/s in the positive x direction. The mass M is:

- A. 5.0 kg.
- B. 9.0 kg.
- C. 6.6 kg.
- D. 4.0 kg.
- E. 8.1 kg.

1

TEST CODE: PAGE: 003

QUESTION NO: 7

0

Three particles are located in the xy plane. The 40 g particle is located at (4, 3) m, and the 50 g particle is located at (-2,-2) m. Where must the 20 g particle be placed so that the center of mass of this three-particle system is at the origin?

- A. (-3.0, -1.0) m.
- B. (-2.0, -0.67) m.
- C. (-6.0, -2.1) m.
- D. (-1.0, 1.3) m.
- E. (2.1, -1.7) m.

QUESTION NO: 8

0

A mass ($m_1=5.0 \text{ kg}$) is connected by a light cord that passes over a pulley, to a mass ($m_2=4.0 \text{ kg}$) which slides on a smooth surface as shown in Figure 5. The pulley (radius 0.2 m) rotates about a frictionless axle. If the acceleration of m_2 is 3.5 m/s^2 , then the moment of inertia of the pulley is:

- A. $0.20 \text{ kg}\cdot\text{m}^2$.
- B. $0.42 \text{ kg}\cdot\text{m}^2$.
- C. $0.08 \text{ kg}\cdot\text{m}^2$.

D. $0.16 \text{ kg}\cdot\text{m}^2$.

E. $0.33 \text{ kg}\cdot\text{m}^2$.

QUESTION NO: 9

0

A thin rod of mass M and length L is free to rotate about A , the midpoint of the rod. The rod is struck at one end by a ball of clay of mass m moving with speed v , as shown in figure 6. The ball sticks to the rod. After collision, the angular momentum of the clay-rod system about A is:

A. $mvL/2$.

B. mvL .

C. $3mvL/2$.

D. $2mvL/5$.

E. $5mvL/2$.

1

TEST CODE: PAGE: 004

QUESTION NO: 10

0

A column of fluid, is open to the atmosphere at the top, and is 9.5 m high. If the density of the fluid is $1680 \text{ kg}/\text{m}^3$, what is the total pressure at the bottom of this column?

A. $2.58 \times 10^5 \text{ Pa}$.

B. $1.75 \times 10^5 \text{ Pa}$.

C. $2.25 \times 10^5 \text{ Pa}$.

D. $1.25 \times 10^5 \text{ Pa}$.

E. $3.65 \times 10^5 \text{ Pa}$.

QUESTION NO: 11

0

The velocity of the flow of water in a pipe is 4.5 m/s . If the pipe has a diameter of 8.4 cm , what is the mass of water coming out of the pipe per second?

A. 24.9 kg/s .

B. 14.5 kg/s .

C. 29.9 kg/s .

D. 18.7 kg/s .

E. 11.3 kg/s .

QUESTION NO: 12

0

A pipe carrying water from the ground floor to the fourth floor of a building which is 13 m high. At the fourth floor the pipe has a cross-sectional area of $4.1 \times 10^{-4} \text{ m}^2$, a pressure of $1.66 \times 10^5 \text{ Pa}$ and the velocity of water flow is 8.4 m/s . At the ground floor, the cross-sectional area of the pipe is $9.3 \times 10^{-4} \text{ m}^2$. What is the pressure in the pipe at the ground floor?

A. $3.22 \times 10^5 \text{ Pa}$.

B. $2.92 \times 10^5 \text{ Pa}$.

- C. 3.41×10^5 Pa.
- D. 2.44×10^5 Pa.
- E. 4.12×10^5 Pa.

1 TEST CODE: PAGE: 005

QUESTION NO: 13

0

The mass of a planet is 6.00×10^{24} kg. The gravitational acceleration on the surface of this planet is 12.0 m/s^2 . The gravitational constant is G is $6.672 \times 10^{-11} \text{ N.m}^2/\text{kg}^2$. Calculate the radius of this planet.

- A. 5776 km.
- B. 5347 km.
- C. 5002 km.
- D. 5112 km.
- E. 5883 km.

QUESTION NO: 14

0

Find the distance from center of the earth to the center of the moon, using the fact that the moon completes an orbit in 27.3 days. ($G=6.672 \times 10^{-11} \text{ N.m}^2/\text{kg}^2$, mass of the earth 5.98×10^{24} kg).

- A. 3.83×10^8 m.
- B. 6.38×10^6 m.
- C. 5.42×10^8 m.
- D. 2.43×10^6 m.
- E. 4.41×10^7 m.

QUESTION NO: 15

0

A particle at the end of a spring executes simple harmonic motion with an amplitude of 4.0 cm. At what displacement (x) will its speed be equal to one half its maximum speed?

- A. 3.46 cm.
- B. 5.20 cm.
- C. 6.93 cm.
- D. 7.12 cm.
- E. 4.13 cm.

1 TEST CODE: PAGE: 006

QUESTION NO: 16

0

A particle of mass $m=0.14$ kg at the end of a spring executes a simple harmonic motion according to the equation:

$$x=0.2 \cos(10t + \pi/2)$$

Find the maximum potential energy of the spring.?

- A. 0.28 J.
- B. 0.36 J.
- C. 0.44 J.

- D. 0.56 J.
- E. 0.62 J.

QUESTION NO: 17

0

Consider a horizontal spring-mass system. The force constant of the spring is $k=360$ N/m and $M=1$ kg is initially at rest. A bullet of mass 100 g is fired with initial speed $v_i= 100$ m/s at the mass M , and embedded in it. Find the maximum amplitude of oscillation of the mass bullet system. (see figure 7)

- A. 0.50 m.
- B. 0.55 m.
- C. 0.67 m.
- D. 0.43 m.
- E. 0.38 m.

QUESTION NO: 18

0

A uniform bar of length 1.2 m and weight 120 N is supported by two ropes (see figure 8). two 400-N weights are suspended at $L/3$ and $2L/3$ from the left end. Find the tension T_1 in the right hand rope.

- A. 531.2 N.
- B. 300.2 N.
- C. 415.7 N.
- D. 117.8 N.
- E. 422.7 N

1 TEST CODE: PAGE: 007

QUESTION NO: 19

0

A uniform circular disc of mass 4 kg is rolling without slipping along a horizontal surface. The velocity of its center of mass is 5 m/s. Its total kinetic energy is:

- A. 75 J.
- B. 48 J.
- C. 108 J.
- D. 50 J
- E. 15 J

QUESTION NO: 20

0

A wheel (radius = 12 cm) is mounted on a frictionless, horizontal axle that is perpendicular to the wheel and passes through the center of mass of the wheel. A light cord wrapped around the wheel supports a mass of 0.40 kg. The mass is released from rest and the cord remains stretched. The mass is observed to fall with a downward acceleration of 3.0 m/s². What is the moment of inertia of the wheel?

- A. 0.013 kg.m²

- B. $0.022 \text{ kg}\cdot\text{m}^2$
- C. $0.008 \text{ kg}\cdot\text{m}^2$
- D. $0.416 \text{ kg}\cdot\text{m}^2$
- E. $0.335 \text{ kg}\cdot\text{m}^2$

QUESTION NO: 21

0

A 10 g bullet is fired into a 990 g wooden block at rest on a horizontal surface that has coefficient of friction equal to 0.5. The bullet remains stuck in the wood, which slides 0.4 m before coming to rest.

The speed of the bullet just before it strikes the block is:

- A. 198 m/s.
- B. 140 m/s.
- C. 171 m/s.
- D. 182 m/s.
- E. 163 m/s.

1

TEST CODE: PAGE: 008

QUESTION NO: 22

0

How large a force is required to accelerate a 1500 kg car that is originally at rest to a speed of 20 m/s in a distance of 80 m.

- A. 3750 N.
- B. 8438 N.
- C. 5859 N.
- D. 3999 N.
- E. 6865 N.

QUESTION NO: 23

0

A 1000 kg car is moving with a constant velocity of 3 m/s.

A constant frictional force of 400 N acts on the car.

What is the power delivered by the motor of the car?

- A. 1200 W.
- B. 1500 W.
- C. 1800 W.
- D. 1600 W.
- E. 600 W.

QUESTION NO: 24

0

A football player on another planet can have a maximum horizontal range of 20 meters if he jumps with an initial speed of 10 m/s.

Find the acceleration due to gravity "g" on this planet:

- A. 5.0 m/s^2 .
- B. 3.2 m/s^2 .
- C. 7.2 m/s^2 .
- D. 9.8 m/s^2 .

E. 1.3 m/s^2 .

1 TEST CODE: PAGE: 009

QUESTION NO: 25

0

An object moves with a constant speed in a horizontal circle of radius R . Its acceleration is 32 m/sec^2 . What would its acceleration have been if it had the same speed but the circle's radius is increased to $4R$?

- A. 8 m/s^2 .
- B. 16 m/s^2 .
- C. 4 m/s^2 .
- D. 12 m/s^2 .
- E. 6 m/s^2 .

QUESTION NO: 26

0

A student jumps vertically upwards. It takes him 0.6 seconds to jump up and come down to his initial position. His initial velocity and the maximum height he reached are, respectively:

- A. 2.94 m/s, 0.44 m.
- B. 1.96 m/s, 0.20 m.
- C. 2.45 m/s, 0.31 m.
- D. 3.35 m/s, 0.66 m.
- E. 4.41, m/s 0.24 m.

QUESTION NO: 27

0

A stone is dropped from the roof of a 60 m high building. At the same time a second stone is thrown vertically upward from the bottom of this building with an initial speed of 20 m/s. Where will the two stones meet? (Hint: they will be at the same height

- A. 15.9 m from the ground.
- B. 19.4 m from the ground.
- C. 17.9 m from the ground.
- D. 21.3 m from the ground.
- E. 11.4 m from the ground.

1 TEST CODE: PAGE: 010

QUESTION NO: 28

0

Two points in a plane have polar coordinates (2.5m, 30 degrees) and (3.8m, 120 degrees). Find the distance between them.

- A. 4.55 m.
- B. 5.17 m.
- C. 5.89 m.
- D. 3.79 m.
- E. 4.94 m

QUESTION NO: 29

0

A motorboat is heading north at 30 km/h relative to the water in a place where the water current has a velocity of 10 km/h in a direction 6 degrees south of east. Find the resultant velocity of the boat.

- A. 21.9 km/h, 76.8 degrees north of east.
- B. 20.5 km/h, 73.0 degrees north of east.
- C. 23.4 km/h, 80.2 degrees north of east.
- D. 17.1 km/h, 66.1 degrees north of east.
- E. 26.4 km/h, 60.0 degrees north of east.

QUESTION NO: 30

0

The mass of a hollow spherical shell of inner radius 5 cm and outer radius 15, cm and of density 25 g/cm^3 is equal to:

- A. 340 kg.
- B. 347 kg.
- C. 331 kg.
- D. 362 kg.
- E. 353 kg.

