

- Q1 Q0 An empty fuel tank of a car needs 50 liters of gasoline  
 Q0 to fill up. Find the volume of the fuel tank in  $\text{m}^3$ .  
 Q0 (1 milliliter =  $1 \text{ cm}^3$ )  
 Q0  
 A1 0.050  
 A2 50 000  
 A3 50  
 A4 500  
 A5 0.50  
 Q0
- Q2 Q0 Fig. 1 shows a graph of position versus time for a particle  
 Q0 moving along the x axis. What is the total distance travelled  
 Q0 by the particle in 15 s?  
 Q0  
 A1 12.5 m  
 A2 7.5 m  
 A3 10 m  
 A4 5.0 m  
 A5 22.5 m  
 Q0
- Q3 Q0 An object starts from rest at the origin and moves along the  
 Q0 x-axis with a constant acceleration of  $5.0 \text{ m/s}^2$ . Find its  
 Q0 average velocity as it goes from  $x = 0 \text{ m}$  to  $x = 10 \text{ m}$ .  
 Q0  
 A1 5.0 m/s  
 A2 10 m/s  
 A3 17 m/s  
 A4 3.0 m/s  
 A5 8.0 m/s  
 Q0
- Q4. Q0 Starting at time  $t = 0$ , an object moves along a straight line  
 Q0 with a velocity in m/s given by  $v = 72 - 2t^2$ ,  
 Q0 where  $t$  is in seconds. Find its acceleration when it stops  
 Q0 momentarily.  
 Q0  
 A1  $-24 \text{ m/s}^2$   
 A2 0  
 A3  $-4.0 \text{ m/s}^2$   
 A4  $-9.8 \text{ m/s}^2$   
 A5  $-4.9 \text{ m/s}^2$   
 Q0
- Q5 Q0 A stone is thrown vertically upward with an initial speed of  
 Q0 15 m/s. What is its speed at a height of 10 m from its release  
 Q0 point?  
 Q0  
 A1 5.4 m/s  
 A2 0  
 A3 It will not reach the height of 10 m.  
 A4 9.8 m/s  
 A5 12 m/s  
 Q0
- Q6 Q0 The angle between the two vectors  $A = 2i + 4j$  and  
 Q0  $B = 4i - 2j$  is:  
 Q0  
 A1 90 degrees  
 A2 27 degrees  
 A3 39 degrees  
 A4 180 degrees  
 A5 0 degrees  
 Q0
- Q7 Q0 As shown in Fig. 3, a block moves down on a 45-degree inclined

- Q0 plane of 2.5 m length, then horizontally for another 2.5 m, and  
 Q0 then falls down vertically a height of 2.5 m. Find the magnitude  
 Q0 and direction of the resultant displacement vector of the block.  
 Q0
- A1 6.0 m and 45 degrees below horizontal axis  
 A2 3.5 m and 30 degrees below horizontal axis  
 A3 6.0 m and 30 degrees below horizontal axis  
 A4 3.5 m and 45 degrees below horizontal axis  
 A5 5.5 m and 60 degrees below horizontal axis  
 Q0
- Q8 Q0 Given the vectors  $A = 3j + 6k$ ,  $B = 15i + 21k$ . Find the  
 Q0 magnitude of vector C that satisfies equation  $2A + 3C - B = 0$ .  
 Q0
- A1 6.16  
 A2 5.48  
 A3 18.5  
 A4 6.71  
 A5 8.60  
 Q0
- 9 Q0 At  $t=0$ , a particle moving in the xy plane with a constant  
 Q0 acceleration of  $a=(2i + 4j)$  m/s\*\*2 has a velocity  $V_0=(-4j)$  m/s  
 Q0 at the origin. Find the speed of the particle at  $t=3$  s.  
 Q0
- A1 10 m/s  
 A2 0  
 A3 4 m/s  
 A4 24 m/s  
 A5 20 m/s  
 Q0
- 10 Q0 A ball is projected from the ground into the air with velocity  
 Q0  $V_0$ . At a height of 10.0 m the velocity is observed to be  
 Q0  $V = 8.5i + 9.1j$  in m/s. Find  $V_0$ .  
 Q0
- A1  $(8.5i + 16.7j)$  m/s  
 A2  $(16.7i + 9.1j)$  m/s  
 A3  $(8.5i + 9.1j)$  m/s  
 A4  $(2.5i + 3.1j)$  m/s  
 A5  $(6.2i + 1.1j)$  m/s  
 Q0
- 11 Q0 Rain is falling vertically at constant speed of 6.0 m/s.  
 Q0 At what angle from the vertical do the rain appear to be falling  
 Q0 as viewed by the driver of a car traveling on a straight, level  
 Q0 road with a speed of 8.0 m/s?  
 Q0
- A1 53 degrees  
 A2 37 degrees  
 A3 49 degrees  
 A4 41 degrees  
 A5 0 degree  
 Q0
- 12 Q0 The speed of a particle moving in uniform circular motion is  
 Q0 doubled while the radius of the path of the particle is  
 Q0 increased by a factor of 4. The new centripetal force needed  
 Q0 will be :  
 Q0
- A1 the same as before  
 A2 half as great as before  
 A3 twice as great as before  
 A4 1/4 of its original value  
 A5 four times as great as before  
 Q0

- 13 Q0 A ball is thrown horizontally with speed  $V_0$  from the edge of  
 Q0 a cliff 35 m high. The ball strikes the ground at a point 80 m  
 Q0 from the base of the cliff. Find  $V_0$ .  
 Q0  
 A1 30 m/s  
 A2 9.8 m/s  
 A3 2.5 m/s  
 A4 22 m/s  
 A5 45 m/s  
 Q0
- 14 Q0 As shown in Fig. 7, a 25-kg box is pushed across a frictionless  
 Q0 horizontal floor with a force of 20 N, directed at an angle of  
 Q0 20 degrees below the horizontal. The magnitude of the  
 Q0 acceleration of the box is:  
 Q0  
 A1 0.75 m/s\*\*2  
 A2 0.27 m/s\*\*2  
 A3 17 m/s\*\*2  
 A4 21 m/s\*\*2  
 A5 0.82 m/s\*\*2  
 Q0
- 15 Q0 An object of mass  $M = 10$  kg moving on frictionless horizontal  
 Q0 surface is subjected to two applied forces as shown in Fig. 2.  
 Q0 In which situation is the object accelerating to the right?  
 Q0  
 A1 (d)  
 A2 (a)  
 A3 (c)  
 A4 (b)  
 A5 (e)  
 Q0
- 16 Q0 Two blocks A ( $M_A = 4$  kg) and B ( $M_B = 20$  kg) are in contact with  
 Q0 each other and are placed on a horizontal frictionless surface.  
 Q0 A 36-N constant force is applied to A as shown in Fig. 4. The  
 Q0 magnitude of the force exerted on A by B is  
 Q0  
 A1 30 N  
 A2 0 N  
 A3 36 N  
 A4 15 N  
 A5 3.6 N  
 Q0
- 17 Q0 Two masses  $m_1 = 2$  kg,  $m_2 = 4$  kg are connected by a light string  
 Q0 that passes over a frictionless and massless pulley (see Fig. 5).  
 Q0 Find the magnitude of the acceleration of the masses.  
 Q0  
 A1 3.27 m/s\*\*2  
 A2 2.15 m/s\*\*2  
 A3 10.5 m/s\*\*2  
 A4 0.75 m/s\*\*2  
 A5 1.23 m/s\*\*2  
 Q0
- 18 Q0 A stone, of mass  $m$ , is attached to a strong string and rotates  
 Q0 in a vertical circle of radius  $R$ . At the bottom of the path the  
 Q0 tension in the string is 3 times the weight of the stone. The  
 Q0 speed of the stone at this point is given by .  
 Q0  
 A1  $\sqrt{2gR}$ .  
 A2  $2\sqrt{gR}$   
 A3  $2gR$   
 A4  $\sqrt{3gR}$

A5  $\sqrt{gR/2}$

Q0

- 19 Q0 A block attached to a string, rotates counter-clockwise in a circle on a smooth horizontal surface. The string breaks at point P (Fig. 6). What path will the block follow?

Q0

A1 path B

A2 path A

A3 path C

A4 path D

A5 path E

Q0

- 20 Q0 A box slides down a 30 degree incline with an acceleration =  $3.2 \text{ m/s}^2$ . Find the coefficient of kinetic friction between the box and the incline.

Q0

A1 0.20

A2 0.25

A3 0.15

A4 0.30

A5 0.62

**PHYS101 - FIRST MAJOR EXAM – FIGURES Term-031**

