




$$(1) \frac{s^2 + s + 1}{s(s^2 + 2s + 1)} = \frac{1}{s} (s^2 + s + 1) \cdot \frac{1}{s^2 + 2s + 1}$$



∴ total Bode magnitude plot is



$$(2) \quad G(s) = \frac{k}{s(s^2 + s + 1)} = \frac{k}{s'(s + 0.5 + j \cdot 0.866)(s + 0.5 - j \cdot 0.866)}$$



Notice that $G(s)$ is stable, so we can use the special case of Nyquist criterion to determine system's stability.

$$\therefore \frac{k}{j\omega((j\omega)^2 + j\omega + 1)} = \frac{k}{-j\omega^3 + j\omega - \omega^2} = \frac{k}{-\omega^2 + j\omega(1 - \omega^2)}$$

$$= \frac{-k\omega^2}{A(\omega)} - j \frac{k\omega(1 - \omega^2)}{A(\omega)}$$

$$A(\omega) = \sqrt{\omega^4 + \omega^2(1 - \omega^2)^2}$$

Intersection with real-axis (Imaginary = 0)

$$1 - \omega^2 = 0 \quad \therefore \omega = 1$$

The real part $-k \frac{1}{\sqrt{1+0}} = -k$.

\therefore Stable value for system is.

$$1 > k > 0$$