

**King Fahd University of Petroleum and Minerals**  
**Department of Mathematical Sciences**  
**Math 101**  
**Final Exam**  
**Semester I, 2001–2002 (011)**  
**Dr. Faisal Fairag**

---

Name:	
-------	--

ID:	
-----	--

Sec:	<b>7      11</b>
------	------------------

Serial No:	
------------	--

---

**Form (1)**

Question #		Points
1		10
2		27
3		34
4		27
5		27
6		9
7-23		8 each
24		3 each
Total:		300

1. Let  $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2$

(a) Find the interval on which  $f$  is decreasing

(b) Find the open interval on which  $f$  is concave up.

2. The equation  $x^3 - x^2 - 2x + 1 = 0$  has one real solution for  $0 < x < 1$ . Approximate it by Newton's Method. If  $x_1 = 1$ , then find  $x_6$ .

3. Find the extreme values for  $f(x) = x^{4/3} - 3x^{1/3}$  on the interval  $[-1, 8]$  and determine where those values occur.

4. Let  $f(x) = x^{2/3}(x + 5)$


(a) Find all relative max. and all relative min.

(b) Find all inflection points

(c) Find all cusp which may or may not exist.

5. A rectangular field is to be bounded by a fence on three sides and by a wall on the fourth one. Find the dimensions of the field with maximum area that can be enclosed with 1000 meters of fence ?

6. The lower corner of a page of width  $k$  is folded over so as just to reach the inner edge of the page ( see Figure ). Find the width of the part folded over (the value of  $x$  ) when the area of the triangle floded over is a minimum.



7. If  $f(x) = x^4 - x$  on  $[-1, 1]$ , find the value  $c$  that satisfies the Mean value Theorem.

- (a) 1
- (b) 0
- (c) 2
- (d) 3
- (e) 4

8.  $\lim_{x \rightarrow 9} \frac{x + 5}{\sqrt{x} - 3} =$

- (a) it does not exist
- (b)  $+\infty$
- (c)  $-\infty$
- (d) 84
- (e) -84

9.  $\lim_{x \rightarrow -\infty} \sqrt{\frac{20x^{10} - 2x^5 + 2}{5x^{10} + x^5 - 3}} =$

- (a)  $+\infty$
- (b) 2
- (c)  $-\infty$
- (d) -2
- (e) it does not exist



10. To prove that  $\lim_{x \rightarrow 5} (x - 2) = 3$  a reasonable relationship between  $\delta$  and  $\epsilon$  would be

(a)  $\delta = \epsilon$

(b)  $\delta = 5\epsilon$

(c)  $\delta = \sqrt{\epsilon}$

(d)  $\delta = \frac{1}{\epsilon}$

(e)  $\delta = \frac{1}{\sqrt{\epsilon}}$

11. Find the value of  $k$ , if possible, that will make the function continuous

$$f(x) = \begin{cases} x + 2k & x \leq 1 \\ kx^2 + x + 1 & x > 1 \end{cases}$$

(a)  $-1$

(b)  $2$

(c)  $1$

(d)  $-2$

(e) none exists

12. find the limit  $\lim_{x \rightarrow +\infty} \left( \cos \left( \frac{4}{x} \right) \cdot \sin \left( \frac{5}{x} \right) \right) =$

(a)  $1$

(b)  $-1$

(c)  $+\infty$

(d)  $-\infty$

(e)  $0$

13. Find an equation for the tangent line to the curve  $y = x^7 - 5$  at  $(1, -4)$ .

(a)  $y = 7x$

(b)  $y = 7x + 5$

(c)  $y = 7x - 3$

(d)  $y = 7x - 11$

(e)  $y = 7x - 5$

14. If  $y = \frac{2}{x+3}$ , then  $y'(0) =$

(a)  $\frac{-2}{9}$

(b) 0

(c)  $\frac{4}{9}$

(d)  $\frac{2}{9}$

(e)  $\frac{-4}{9}$

15.  $g(x) = x^3 f(x)$ . Find  $g'(2)$ , given that  $f(2) = 6$  and  $f'(2) = 3$

(a) 48

(b) -60

(c) 96

(d) 60

(e) -48

16.  $y = x^{-3} + x$ . Find  $y'''$

(a) -6

(b)  $-60x^{-6} + x^{-2}$

(c)  $-60x^{-6} - x^{-2}$

(d)  $-3x^{-2} + 1$

(e)  $-60x^{-6}$

17. If  $y = x^5 \cos x$ , find  $d^2y/dx^2$

- (a)  $20x^3 \cos x$
- (b)  $20x^3 \cos x - x^5 \cos x$
- (c)  $20x^3 \cos x + x^5 \cos x$
- (d)  $20x^3 \cos x - 10x^4 \sin x - x^5 \cos x$
- (e)  $-x^5 \cos x$

18. Use  $dy$  to approximate  $\sqrt{4.04}$  starting at  $x = 4$

- (a) 2.01
- (b) 1.99
- (c) 4.01
- (d) 3.99
- (e) 1.59

19. Find  $\frac{dy}{dx}$  if  $x^2 + y^2 = 49$ .

- (a)  $\frac{49x}{y}$
- (b)  $\frac{x}{y}$
- (c)  $-\frac{x}{y}$
- (d)  $-\frac{49x}{y}$
- (e)  $\frac{y}{x}$

20. The number of critical points for  $f(x) = |(x-1)(x-2)(x-3)|$  is

- (a) 5
- (b) 2
- (c) 3
- (d) 4
- (e) 1

21.  $y = \cot^{-1} \sqrt{x}$ . Find  $dy/dx$ .

- (a)  $-\frac{x}{1+x}$
- (b)  $-\frac{1}{2\sqrt{x}(1+x)}$
- (c)  $-\frac{x}{(1+x^2)}$
- (d)  $-\sqrt{\frac{1}{1+x^2}}$
- (e)  $\frac{x}{1+x}$

22.  $\lim_{x \rightarrow 0^+} \frac{\sin x}{\ln(x^2 + 1)} =$

- (a)  $-\infty$
- (b) 10
- (c) 0
- (d)  $+\infty$
- (e) -10

23.  $\lim_{x \rightarrow 0^+} (1 - \ln(2x))^{2x} =$

- (a) 0
- (b) 1
- (c)  $+\infty$
- (d)  $-\infty$
- (e) -1

24. True (T) or False (F)

(a)  $f(x) = \frac{1}{x^3}$  on  $[-1, 1]$  satisfies the hypotheses of Rolle's Theorem ( )

(b) The Mean Value Theorem can be used on  $f(x) = |x - 1|$  on  $[-2, 1]$  ( )

(c)  $\tan x$  has a point of inflection on  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  ( )

(d)  $f(x) = x^{3/5}$  has a critical point ( )

(e) All relative extrema occur at critical points ( )

(f)  $f(x) = |\tan^2 x|$  has no relative extrema on  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  ( )

(g)  $f(x) = \frac{x^3}{x^5 - 2}$  has no horizontal asymptote ( )

(h)  $f(x) = |x^2 - 4|$  has points of discontinuity at  $x = 2$  and  $x = -2$ . ( )

(i) The function  $f(x) = \frac{x+5}{x-1}$  has a removable discontinuity at  $x = 1$  ( )

(j) The slope of the tangent line to the graph of  $f(x) = x^3 - 5$  at  $x_0 = 3$  is 22. ( )